



Health
Canada

Santé
Canada

Statistics
Canada

Statistique
Canada

Canadian Community Health Survey Cycle 2.2, Nutrition (2004)

Nutrient Intakes from Food

Provincial, Regional and National Summary Data Tables
Volume 2

Revised February 2009



Canada 

Health Canada is the federal department responsible for helping the people of Canada maintain and improve their health.

We assess the safety of drugs and many consumer products, help improve the safety of food, and provide information to Canadians to help them make healthy decisions. We provide health services to First Nations people and to Inuit communities. We work with the provinces to ensure our health care system serves the needs of Canadians.

Published by authority of the Minister of Health.

**Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) —
Nutrient Intakes from Food**

Provincial, Regional and National Summary Data Tables, Volume 2

is available on Internet at the following address:

www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/index-eng.php

Également disponible en français sous le titre :

Enquête sur la santé dans les collectivités canadiennes, cycle 2.2, Nutrition (2004) —

Apports nutritionnels provenant des aliments

Tableaux sommaires provinciaux, régionaux et nationaux, Volume 2

This publication can be made available on request on
diskette, large print, audio-cassette and braille.

For further information or to obtain additional copies, please contact:

Publications

Health Canada

Ottawa, Ontario K1A 0K9

Tel.: (613) 954-5995

Fax: (613) 941-5366

E-Mail: info@hc-sc.gc.ca

© Her Majesty the Queen in Right of Canada, represented by
the Minister of Health Canada, 2008

This publication may be reproduced without permission
provided the source is fully acknowledged.

Cat.: H164-45/2-2008E-PDF

ISBN: 978-0-662-48761-6

Acknowledgements

Health Canada would like to acknowledge and thank the individuals who have contributed to this report. This publication was produced as a joint venture between Health Canada and Statistics Canada. Technical experts from Health Canada's Food Directorate and Office of Nutrition Policy and Promotion and from Statistics Canada's Health Statistics Division developed the report. Project guidance was provided by Health Canada's Food and Nutrition Surveillance System Working Group, which consists of individuals from Health Canada, the Public Health Agency of Canada and the Federal/Provincial/Territorial Group on Nutrition. External experts have advised both departments, through the CCHS 2.2 Users Group, on the need for a consistent approach to analyzing and reporting these data. The draft report was reviewed by internal experts from both departments.

Table of Contents

	Acknowledgements	i
	List of Tables.....	iii
	List of Appendices	viii
	List of Abbreviations.....	ix
I	Introduction.....	1
II	Summary Data Tables	3
	<i>(table numbering continued from Volume 1)</i>	
	14. Folate (DFE/d): Usual intakes from food	3
	15. Iron (mg/d): Usual intakes from food	17
	16. Linoleic acid (g/d): Usual intakes from food	31
	17. Percentage of total energy intake from linoleic acid	45
	18. Magnesium (mg/d): Usual intakes from food	59
	19. Niacin (NE/d): Usual intakes from food	73
	20. Phosphorus (mg/d): Usual intakes from food.....	87
	21. Potassium (mg/d): Usual intakes from food.....	101
	22. Riboflavin (mg/d): Usual intakes from food	115
	23. Thiamin (mg/d): Usual intakes from food	129
	24. Vitamin B ₆ (mg/d): Usual intakes from food.....	143
	25. Vitamin B ₁₂ (µg/d): Usual intakes from food.....	157
	26. Vitamin C (mg/d): Usual intakes from food (by smoking status)....	171
	27. Vitamin D (µg/d): Usual intakes from food	175
	28. Zinc (mg/d): Usual intakes from food	189

List of Tables

14. Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population	3
Table 14.1 Newfoundland and Labrador, 2004	4
Table 14.2 Prince Edward Island, 2004	5
Table 14.3 Nova Scotia, 2004	6
Table 14.4 New Brunswick, 2004	7
Table 14.5 Quebec, 2004	8
Table 14.6 Ontario, 2004	9
Table 14.7 Manitoba, 2004	10
Table 14.8 Saskatchewan, 2004	11
Table 14.9 Alberta, 2004	12
Table 14.10 British Columbia, 2004	13
Table 14.11 Atlantic Region, 2004	14
Table 14.12 Prairie Region, 2004	15
Table 14.13 Canada excluding territories, 2004	16
15. Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population	17
Table 15.1 Newfoundland and Labrador, 2004	18
Table 15.2 Prince Edward Island, 2004	19
Table 15.3 Nova Scotia, 2004	20
Table 15.4 New Brunswick, 2004	21
Table 15.5 Quebec, 2004	22
Table 15.6 Ontario, 2004	23
Table 15.7 Manitoba, 2004	24
Table 15.8 Saskatchewan, 2004	25
Table 15.9 Alberta, 2004	26
Table 15.10 British Columbia, 2004	27
Table 15.11 Atlantic Region, 2004	28
Table 15.12 Prairie Region, 2004	29
Table 15.13 Canada excluding territories, 2004	30
16. Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population	31
Table 16.1 Newfoundland and Labrador, 2004	32
Table 16.2 Prince Edward Island, 2004	33
Table 16.3 Nova Scotia, 2004	34
Table 16.4 New Brunswick, 2004	35
Table 16.5 Quebec, 2004	36
Table 16.6 Ontario, 2004	37
Table 16.7 Manitoba, 2004	38
Table 16.8 Saskatchewan, 2004	39
Table 16.9 Alberta, 2004	40
Table 16.10 British Columbia, 2004	41
Table 16.11 Atlantic Region, 2004	42
Table 16.12 Prairie Region, 2004	43
Table 16.13 Canada excluding territories, 2004	44

17. Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population.....	45
Table 17.1 Newfoundland and Labrador, 2004.....	46
Table 17.2 Prince Edward Island, 2004.....	47
Table 17.3 Nova Scotia, 2004.....	48
Table 17.4 New Brunswick, 2004	49
Table 17.5 Quebec, 2004.....	50
Table 17.6 Ontario, 2004	51
Table 17.7 Manitoba, 2004	52
Table 17.8 Saskatchewan, 2004	53
Table 17.9 Alberta, 2004.....	54
Table 17.10 British Columbia, 2004.....	55
Table 17.11 Atlantic Region, 2004	56
Table 17.12 Prairie Region, 2004.....	57
Table 17.13 Canada excluding territories, 2004.....	58
18. Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population.....	59
Table 18.1 Newfoundland and Labrador, 2004.....	60
Table 18.2 Prince Edward Island, 2004.....	61
Table 18.3 Nova Scotia, 2004.....	62
Table 18.4 New Brunswick, 2004	63
Table 18.5 Quebec, 2004.....	64
Table 18.6 Ontario, 2004	65
Table 18.7 Manitoba, 2004	66
Table 18.8 Saskatchewan, 2004	67
Table 18.9 Alberta, 2004.....	68
Table 18.10 British Columbia, 2004.....	69
Table 18.11 Atlantic Region, 2004	70
Table 18.12 Prairie Region, 2004.....	71
Table 18.13 Canada excluding territories, 2004.....	72
19. Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population.....	73
Table 19.1 Newfoundland and Labrador, 2004.....	74
Table 19.2 Prince Edward Island, 2004.....	75
Table 19.3 Nova Scotia, 2004.....	76
Table 19.4 New Brunswick, 2004	77
Table 19.5 Quebec, 2004.....	78
Table 19.6 Ontario, 2004	79
Table 19.7 Manitoba, 2004	80
Table 19.8 Saskatchewan, 2004	81
Table 19.9 Alberta, 2004.....	82
Table 19.10 British Columbia, 2004.....	83
Table 19.11 Atlantic Region, 2004	84
Table 19.12 Prairie Region, 2004.....	85
Table 19.13 Canada excluding territories, 2004.....	86

20. Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population	87
Table 20.1 Newfoundland and Labrador, 2004	88
Table 20.2 Prince Edward Island, 2004	89
Table 20.3 Nova Scotia, 2004	90
Table 20.4 New Brunswick, 2004	91
Table 20.5 Quebec, 2004	92
Table 20.6 Ontario, 2004	93
Table 20.7 Manitoba, 2004	94
Table 20.8 Saskatchewan, 2004	95
Table 20.9 Alberta, 2004	96
Table 20.10 British Columbia, 2004	97
Table 20.11 Atlantic Region, 2004	98
Table 20.12 Prairie Region, 2004	99
Table 20.13 Canada excluding territories, 2004	100
21. Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population	101
Table 21.1 Newfoundland and Labrador, 2004	102
Table 21.2 Prince Edward Island, 2004	103
Table 21.3 Nova Scotia, 2004	104
Table 21.4 New Brunswick, 2004	105
Table 21.5 Quebec, 2004	106
Table 21.6 Ontario, 2004	107
Table 21.7 Manitoba, 2004	108
Table 21.8 Saskatchewan, 2004	109
Table 21.9 Alberta, 2004	110
Table 21.10 British Columbia, 2004	111
Table 21.11 Atlantic Region, 2004	112
Table 21.12 Prairie Region, 2004	113
Table 21.13 Canada excluding territories, 2004	114
22. Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population	115
Table 22.1 Newfoundland and Labrador, 2004	116
Table 22.2 Prince Edward Island, 2004	117
Table 22.3 Nova Scotia, 2004	118
Table 22.4 New Brunswick, 2004	119
Table 22.5 Quebec, 2004	120
Table 22.6 Ontario, 2004	121
Table 22.7 Manitoba, 2004	122
Table 22.8 Saskatchewan, 2004	123
Table 22.9 Alberta, 2004	124
Table 22.10 British Columbia, 2004	125
Table 22.11 Atlantic Region, 2004	126
Table 22.12 Prairie Region, 2004	127
Table 22.13 Canada excluding territories, 2004	128

23. Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population.....	129
Table 23.1 Newfoundland and Labrador, 2004.....	130
Table 23.2 Prince Edward Island, 2004.....	131
Table 23.3 Nova Scotia, 2004.....	132
Table 23.4 New Brunswick, 2004	133
Table 23.5 Quebec, 2004.....	134
Table 23.6 Ontario, 2004	135
Table 23.7 Manitoba, 2004	136
Table 23.8 Saskatchewan, 2004	137
Table 23.9 Alberta, 2004.....	138
Table 23.10 British Columbia, 2004.....	139
Table 23.11 Atlantic Region, 2004	140
Table 23.12 Prairie Region, 2004.....	141
Table 23.13 Canada excluding territories, 2004.....	142
24. Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population.....	143
Table 24.1 Newfoundland and Labrador, 2004.....	144
Table 24.2 Prince Edward Island, 2004.....	145
Table 24.3 Nova Scotia, 2004.....	146
Table 24.4 New Brunswick, 2004	147
Table 24.5 Quebec, 2004.....	148
Table 24.6 Ontario, 2004	149
Table 24.7 Manitoba, 2004	150
Table 24.8 Saskatchewan, 2004	151
Table 24.9 Alberta, 2004.....	152
Table 24.10 British Columbia, 2004.....	153
Table 24.11 Atlantic Region, 2004	154
Table 24.12 Prairie Region, 2004.....	155
Table 24.13 Canada excluding territories, 2004.....	156
25. Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population.....	157
Table 25.1 Newfoundland and Labrador, 2004.....	158
Table 25.2 Prince Edward Island, 2004.....	159
Table 25.3 Nova Scotia, 2004.....	160
Table 25.4 New Brunswick, 2004	161
Table 25.5 Quebec, 2004.....	162
Table 25.6 Ontario, 2004	163
Table 25.7 Manitoba, 2004	164
Table 25.8 Saskatchewan, 2004	165
Table 25.9 Alberta, 2004.....	166
Table 25.10 British Columbia, 2004.....	167
Table 25.11 Atlantic Region, 2004	168
Table 25.12 Prairie Region, 2004.....	169
Table 25.13 Canada excluding territories, 2004.....	170

26. Vitamin C (mg/d): Usual intakes from food, by smoking status, household population.....	171
Table 26.1 By sex and region, aged 19 and older, 2004...	172
Table 26.2 By DRI age–sex group, Canada excluding territories, 2004.....	173
27. Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population	175
Table 27.1 Newfoundland and Labrador, 2004	176
Table 27.2 Prince Edward Island, 2004	177
Table 27.3 Nova Scotia, 2004	178
Table 27.4 New Brunswick, 2004	179
Table 27.5 Quebec, 2004	180
Table 27.6 Ontario, 2004.....	181
Table 27.7 Manitoba, 2004.....	182
Table 27.8 Saskatchewan, 2004.....	183
Table 27.9 Alberta, 2004	184
Table 27.10 British Columbia, 2004.....	185
Table 27.11 Atlantic Region, 2004	186
Table 27.12 Prairie Region, 2004	187
Table 27.13 Canada excluding territories, 2004	188
28. Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population	189
Table 28.1 Newfoundland and Labrador, 2004	190
Table 28.2 Prince Edward Island, 2004	191
Table 28.3 Nova Scotia, 2004	192
Table 28.4 New Brunswick, 2004	193
Table 28.5 Quebec, 2004	194
Table 28.6 Ontario, 2004.....	195
Table 28.7 Manitoba, 2004.....	196
Table 28.8 Saskatchewan, 2004.....	197
Table 28.9 Alberta, 2004	198
Table 28.10 British Columbia, 2004.....	199
Table 28.11 Atlantic Region, 2004	200
Table 28.12 Prairie Region, 2004	201
Table 28.13 Canada excluding territories, 2004	202

List of Appendices

Appendix A: Table Footnotes	203
Appendix B: Iron Estimation.....	207
Appendix C: Justification for Excluding Nutrients from Volume 2 and Volume 3	209
List of Nutrients Included in the Three-Volume Set	210
Appendix D: References	211

List of Abbreviations

Abbreviation	Meaning
AI	Adequate Intake
AMDR	Acceptable Macronutrient Distribution Range
CCHS	Canadian Community Health Survey
CV	coefficient of variation
d	day
DFE	Dietary Folate Equivalent
DRI	Dietary Reference Intake
EAR	Estimated Average Requirement
g	gram
IOM	Institute of Medicine
µg	microgram (sometimes reported as mcg)
mg	milligram
n	sample size
NE	Niacin Equivalent
SE	standard error
SIDE	Software for Intake Distribution Estimation
UL	Tolerable Upper Intake Level

I Introduction

This is Volume 2 of a three-volume set that provides summary data tables about the nutrient intakes from food obtained by Canadians in 2004. The data come from the Canadian Community Health Survey (CCHS), Cycle 2.2, Nutrition (2004). As in Volume 1, data are provided for 14 Dietary Reference Intake (DRI) age–sex groups (with the exception of Vitamin C, which is provided by smoking status). For nutrients that have DRIs, the tables also compare usual intakes of these nutrients to the DRIs. Data used for producing the tables in this report were obtained from the CCHS 2.2 Share File. The nutrient intakes represent food consumption; data on nutrient intakes from vitamin and mineral supplements were not included in the production of these tables.¹

This series of products released by Health Canada is part of its ongoing support to users of the CCHS 2.2 data. It has been undertaken as a joint venture with Statistics Canada. The series is a reference for those who will use the CCHS Cycle 2.2 data and its findings to guide nutrition-related program and policy decisions. It will be of particular benefit to provincial ministries of health, researchers and graduate students, policy makers and analysts, public health professionals, epidemiologists, dietitians, the food industry, and the health media.

All three volumes consist primarily of data tables; they do not provide any interpretation or draw conclusions. To optimize the usage of this volume, we recommend that it be read in concert with Volume 1, as well as with the report, *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data* (available at www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_focus-volet_escc-eng.php), published by Health Canada in 2006. That report includes an overview of the CCHS 2.2, including a description of the survey sample, how the survey was conducted, survey components and an introduction to DRIs, including examples of how to interpret the CCHS 2.2 data.

¹ Because supplements may make meaningful contributions to nutrient intakes, inferences about the prevalence of nutrient excess or inadequacy based on intakes from food alone may respectively underestimate or overestimate the prevalences based on total nutrient intakes from both food and supplements.

Volume 2 includes 15 sets of data tables, on 14 nutrients. Results are presented for 13 geographical areas: the 10 provinces, the Atlantic Region, the Prairie Region, and Canada excluding the territories. Data from the four Atlantic provinces and the three Prairie provinces were combined into the Atlantic Region and the Prairie Region, respectively, given the small sample sizes in these provinces.

A revised list of the data tables that appear in all three volumes is found in **Appendix C**.

Quality Assurance and Quality Control

Data quality processes were established to oversee the data analysis and production of the data tables. As the project developed, these processes also helped to monitor and address methodological issues that arose. The processes followed were jointly developed and agreed to by Health Canada and Statistics Canada. Refer to Volume 1 for a description of the methodology used to produce the tables, including the process used and the means of addressing problems encountered.

II Summary Data Tables

14. Folate (DFE/d): Usual intakes from food

Table 14.1 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	262 (20)	146 (16)	162 (17)	195 (18)	236 (21)	282 (26)	333 (32)	368 (37)	120	<3	
	4-8	127	430 (40)	309 (36)	335 (39)	383 (45)	447 (51)	525 (61)	614 (77)	677 (90)	160	0.0	(0.0)
Male	9-13	111	505 (31)	333 (43)	360 (39)	407 (36)	472 (38)	562 (48)	672 (71)	745 (89)	250	<3	
	14-18	107	537 (43)	395 (66) ^E	433 (60)	501 (52)	584 (52)	677 (67)	772 (95)	835 (117)	330	F	
	19-30	77	546 (57)	376 (59)	417 (61)	490 (65)	575 (70)	661 (74)	739 (77)	784 (79)	320	F	
	31-50	145	428 (26)	243 (42) ^E	271 (38)	325 (33)	395 (30)	483 (37)	583 (56)	653 (73)	320	F	
	51-70	182	431 (24)	201 (32)	235 (32)	304 (30)	397 (29)	512 (34)	637 (48)	722 (62)	320	29.3	(7.9) ^E
	>70	63	451 (34)	331 (46)	354 (44)	397 (43)	451 (48)	513 (59)	576 (74)	616 (86)	320	F	
	19+	467	455 (19)	256 (26)	292 (24)	353 (22)	434 (23)	536 (27)	643 (36)	718 (45)	320	F	
	Female												
Female	9-13	96	382 (25)	223 (33)	251 (31)	303 (29)	372 (31)	454 (41)	540 (54)	597 (65)	250	F	
	14-18	105	397 (30)	220 (59) ^E	249 (54) ^E	303 (46)	375 (41)	459 (51)	543 (73)	597 (89)	330	F	
	19-30	91	345 (24)	249 (39)	267 (36)	297 (31)	332 (31)	368 (37)	403 (48)	423 (55)	320	F	
	31-50	167	374 (25)	238 (38)	267 (37)	321 (35)	389 (36)	471 (44)	558 (58)	617 (69)	320	F	
	51-70	198	388 (28)	197 (51) ^E	228 (46) ^E	286 (36)	366 (30)	473 (45)	617 (93)	728 (142) ^E	320	F	
	>70	74	401 (48)	263 (49) ^E	286 (48) ^E	331 (47)	390 (49)	461 (58)	537 (74)	590 (87)	320	F	
	19+	530	376 (17)	230 (22)	259 (22)	310 (21)	378 (22)	461 (26)	552 (36)	613 (46)	320	28.6	(7.5) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.2 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	58	268 (22)	127 (31) ^E	154 (30) ^E	205 (29)	268 (29)	337 (33)	405 (39)	447 (44)	120	F	
	4-8	110	363 (18)	241 (35)	264 (31)	306 (24)	357 (23)	410 (34)	462 (50)	495 (61)	160	<3	
Male	9-13	95	423 (23)	330 (35)	344 (36)	368 (36)	397 (37)	425 (38)	452 (40)	468 (40)	250	<3	
	14-18	87	523 (38)	339 (65) ^E	382 (60)	460 (53)	553 (55)	655 (69)	753 (89)	815 (104)	330	F	
	19-30	70	557 (51)	274 (71) ^E	322 (67) ^E	414 (61)	536 (61)	687 (83)	857 (126)	978 (162)	320	F	
	31-50	109	475 (30)	408 (65)	419 (56)	437 (43)	459 (41)	481 (60)	502 (89) ^E	515 (109) ^E	320	F	
	51-70	128	419 (20)	304 (37)	328 (33)	370 (28)	423 (27)	486 (33)	550 (43)	591 (52)	320	F	
	>70	65	392 (21)	251 (29)	278 (28)	326 (27)	387 (29)	455 (36)	522 (48)	566 (58)	320	F	
	19+	372	467 (17)	287 (22)	319 (21)	379 (20)	456 (22)	545 (29)	641 (41)	706 (52)	320	F	
	Female	9-13	75	384 (31)	221 (47) ^E	253 (45) ^E	317 (41)	397 (43)	478 (52)	556 (65)	609 (77)	250	F
14-18	81	384 (25)	183 (47) ^E	217 (42) ^E	280 (33)	360 (29)	448 (38)	535 (56)	591 (71)	330	40.4	(11.2) ^E	
19-30	101	401 (28)	269 (40)	299 (38)	352 (36)	415 (37)	482 (42)	547 (53)	587 (61)	320	F		
31-50	116	376 (26)	209 (34)	243 (30)	303 (26)	377 (29)	461 (42)	546 (59)	603 (71)	320	30.3	(9.5) ^E	
51-70	146	363 (19)	286 (39)	301 (35)	326 (29)	355 (26)	386 (29)	415 (38)	433 (45)	320	F		
>70	94	314 (24)	176 (30) ^E	199 (29)	243 (27)	303 (28)	377 (35)	460 (48)	517 (60)	320	56.7	(12.0) ^E	
19+	457	369 (14)	233 (18)	261 (17)	310 (16)	370 (17)	437 (21)	504 (28)	548 (34)	320	28.9	(6.4) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.3 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	112	297 (19)	154 (28) ^E	181 (25)	231 (22)	295 (24)	366 (31)	438 (44)	484 (53)	120	F	
	4-8	177	367 (19)	259 (21)	281 (23)	327 (25)	379 (28)	427 (28)	470 (29)	501 (31)	160	<3	
Male	9-13	111	466 (27)	341 (50)	367 (45)	412 (38)	466 (35)	526 (42)	584 (57)	622 (69)	250	<3	
	14-18	113	502 (40)	302 (73) ^E	345 (66) ^E	426 (53)	509 (51)	594 (65)	690 (89)	759 (111)	330	F	
	19-30	91	523 (39)	357 (62) ^E	384 (57)	432 (49)	488 (44)	548 (48)	605 (62)	641 (74)	320	F	
	31-50	101	521 (36)	329 (56) ^E	365 (50)	430 (43)	506 (40)	592 (48)	680 (64)	736 (78)	320	F	
	51-70	134	445 (33)	296 (55) ^E	321 (48)	366 (38)	429 (35)	508 (49)	589 (77)	644 (101)	320	F	
	>70	56	398 (26)	295 (39)	314 (36)	349 (33)	389 (33)	433 (41)	476 (56)	502 (68)	320	F	
	19+	382	487 (19)	298 (29)	330 (26)	388 (22)	462 (20)	548 (26)	639 (39)	700 (50)	320	F	
	Female	9-13	105	382 (20)	346 (55)	356 (47)	373 (35)	393 (22)	412 (34)	429 (52)	440 (65)	250	F
14-18	120	402 (46)	F	227 (69) ^E	306 (56) ^E	399 (48)	500 (54)	604 (75)	675 (95)	330	F		
19-30	91	342 (31)	241 (26)	256 (28)	284 (32)	318 (37)	358 (43)	401 (48)	430 (51)	320	F		
31-50	159	384 (23)	234 (35)	262 (32)	313 (28)	378 (29)	451 (39)	521 (55)	565 (66)	320	F		
51-70	174	399 (33)	222 (45) ^E	250 (42) ^E	304 (36)	377 (33)	464 (45)	558 (69)	622 (89)	320	F		
>70	80	301 (22)	214 (34)	230 (32)	259 (30)	295 (32)	338 (39)	383 (51)	415 (63)	320	65.7 (19.6) ^E		
19+	504	370 (15)	241 (26)	265 (23)	308 (20)	362 (18)	424 (24)	489 (35)	533 (45)	320	30.3 (9.2) ^E		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.4 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	99	257 (17)	168 (27)	185 (25)	215 (22)	252 (20)	294 (24)	336 (34)	364 (42)	120	<3	
	4-8	140	413 (28)	251 (42) ^E	279 (39)	334 (34)	406 (33)	490 (43)	574 (62)	628 (78)	160	<3	
Male													
	9-13	92	453 (38)	339 (45)	357 (45)	387 (45)	422 (46)	459 (46)	495 (48)	517 (48)	250	<3	
	14-18	107	603 (50)	436 (36)	474 (39)	544 (45)	632 (54)	733 (64)	839 (77)	911 (86)	330	<3	
	19-30	73	673 (68)	423 (87) ^E	473 (83) ^E	565 (80)	681 (87)	814 (112)	947 (151)	1034 (182) ^E	320	F	
	31-50	134	525 (39)	324 (57) ^E	355 (53)	413 (45)	486 (43)	570 (56)	657 (83)	713 (106)	320	F	
	51-70	131	456 (28)	311 (20)	336 (22)	382 (26)	439 (32)	506 (40)	582 (48)	637 (54)	320	F	
	>70	55	413 (32)	297 (49) ^E	317 (46)	352 (42)	397 (39)	448 (46)	502 (79)	540 (133) ^E	320	F	
	19+	393	525 (24)	340 (39)	372 (35)	431 (30)	509 (28)	602 (39)	702 (60)	769 (77)	320	F	
Female													
	9-13	79	397 (31)	268 (50) ^E	292 (46)	335 (42)	386 (40)	446 (47)	514 (63)	562 (79)	250	F	
	14-18	104	414 (32)	369 (80) ^E	380 (68) ^E	398 (50)	418 (39)	439 (46)	458 (68)	470 (86) ^E	330	F	
	19-30	101	380 (37)	210 (67) ^E	242 (65) ^E	303 (60) ^E	384 (54)	480 (54)	580 (65)	647 (79)	320	F	
	31-50	143	408 (33)	214 (55) ^E	244 (50) ^E	304 (42)	388 (38)	486 (53)	585 (79)	649 (99)	320	F	
	51-70	193	392 (20)	230 (43) ^E	259 (38)	309 (30)	372 (25)	446 (30)	528 (47)	587 (64)	320	F	
	>70	94	323 (20)	196 (23)	218 (22)	256 (22)	299 (26)	348 (34)	402 (47)	439 (57)	320	61.8	(13.8) ^E
	19+	531	387 (17)	211 (22)	242 (20)	300 (19)	375 (19)	461 (25)	550 (32)	609 (38)	320	31.5	(6.6) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.5 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	311	304	(15)	142 (22)	172 (21)	227 (19)	297 (19)	375 (24)	450 (31)	498 (37)	120	F
	4-8	485	433	(18)	329 (45)	353 (40)	394 (32)	443 (25)	498 (29)	552 (42)	588 (54)	160	<3
Male													
	9-13	277	556	(29)	392 (53)	426 (49)	491 (42)	573 (40)	670 (52)	769 (75)	833 (93)	250	<3
	14-18	339	608	(26)	373 (44)	418 (42)	504 (37)	615 (34)	740 (58)	870 (91)	966 (107)	330	F
	19-30	237	643	(30)	543 (70)	568 (61)	612 (47)	663 (40)	716 (51)	765 (74)	796 (90)	320	<3
	31-50	423	575	(23)	359 (35)	398 (32)	471 (29)	565 (28)	676 (34)	793 (49)	872 (62)	320	F
	51-70	387	509	(17)	320 (30)	355 (27)	419 (23)	499 (21)	591 (27)	683 (39)	743 (49)	320	F
	>70	132	439	(30)	220 (45) ^E	260 (42)	329 (41)	412 (43)	508 (52)	609 (68)	679 (83)	320	F
	19+	1179	559	(13)	355 (20)	393 (19)	463 (17)	551 (17)	653 (20)	757 (27)	826 (33)	320	F
Female													
	9-13	281	456	(21)	281 (34)	313 (32)	372 (30)	447 (30)	530 (35)	613 (45)	666 (53)	250	F
	14-18	321	488	(20)	309 (30)	345 (28)	408 (25)	487 (25)	578 (32)	673 (44)	736 (54)	330	F
	19-30	249	505	(33)	418 (73) ^E	436 (66)	466 (53)	501 (40)	538 (65)	572 (115) ^E	593 (158) ^E	320	F
	31-50	364	472	(27)	355 (56)	382 (51)	430 (42)	487 (35)	546 (38)	603 (53)	640 (68)	320	F
	51-70	467	435	(14)	273 (26)	304 (23)	362 (18)	434 (16)	512 (23)	589 (33)	637 (41)	320	F
	>70	215	356	(15)	211 (25)	238 (24)	287 (23)	351 (24)	427 (28)	510 (38)	567 (47)	320	37.8 (9.6) ^E
	19+	1295	453	(14)	294 (20)	325 (19)	383 (17)	456 (17)	537 (21)	617 (28)	671 (34)	320	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.6 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)									
Both																				
	1-3	644	278	(9)	146	(19)	171	(16)	216	(13)	273	(10)	338	(13)	405	(22)	451	(29)	120	F
	4-8	956	389	(9)	235	(20)	264	(18)	316	(13)	382	(10)	458	(14)	537	(24)	590	(32)	160	<3
Male																				
	9-13	589	465	(13)	310	(32)	339	(28)	392	(20)	457	(15)	531	(21)	608	(35)	658	(47)	250	<3
	14-18	639	541	(17)	338	(51)	376	(44)	447	(32)	538	(22)	641	(33)	747	(58)	816	(78)	330	F
	19-30	481	552	(24)	373	(52)	407	(45)	467	(32)	541	(25)	623	(42)	706	(72)	760	(96)	320	F
	31-50	709	489	(18)	351	(59) ^E	379	(51)	428	(37)	488	(22)	556	(32)	625	(64)	669	(89)	320	F
	51-70	758	490	(14)	274	(38)	313	(33)	387	(24)	479	(16)	583	(22)	694	(43)	772	(62)	320	F
	>70	734	421	(15)	214	(16)	247	(15)	307	(14)	389	(14)	497	(19)	625	(32)	721	(45)	320	28.8 (4.2)
	19+	2682	497	(10)	288	(25)	325	(22)	393	(18)	482	(13)	588	(17)	701	(31)	779	(42)	320	F
Female																				
	9-13	585	417	(12)	278	(39)	304	(33)	352	(23)	410	(14)	476	(22)	542	(40)	586	(53)	250	F
	14-18	645	442	(14)	283	(37)	314	(33)	370	(24)	440	(17)	517	(24)	596	(41)	647	(54)	330	F
	19-30	514	368	(13)	228	(33)	253	(28)	299	(20)	355	(15)	418	(23)	481	(37)	522	(49)	320	34.0 (9.4) ^E
	31-50	758	409	(13)	239	(29)	267	(26)	320	(20)	390	(16)	475	(21)	564	(36)	623	(48)	320	24.9 (7.2) ^E
	51-70	955	390	(12)	213	(20)	244	(18)	302	(15)	376	(14)	460	(17)	546	(25)	603	(32)	320	30.9 (5.1)
	>70	1345	353	(8)	203	(16)	228	(14)	274	(11)	334	(9)	407	(12)	485	(21)	536	(29)	320	44.4 (4.2)
	19+	3572	389	(7)	226	(11)	254	(10)	306	(9)	373	(8)	453	(11)	536	(17)	592	(22)	320	29.9 (3.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.7 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	(SE)							
Both																				
	1-3	324	273	(16)	119	(17)	144	(17)	192	(16)	256	(16)	328	(22)	407	(31)	463	(40)	120	F
	4-8	425	360	(10)	245	(38)	268	(32)	308	(19)	355	(11)	406	(23)	456	(42)	488	(56)	160	<3
Male																				
	9-13	274	465	(17)	339	(15)	362	(15)	404	(16)	457	(18)	518	(21)	580	(25)	621	(28)	250	<3
	14-18	297	591	(27)	271	(41)	328	(39)	435	(32)	577	(30)	741	(47)	910	(74)	1021	(95)	330	F
	19-30	249	635	(55)	410	(35)	452	(39)	529	(46)	629	(56)	750	(70)	881	(89)	972	(103)	320	<3
	31-50	309	499	(24)	347	(50)	376	(43)	428	(32)	492	(28)	564	(40)	634	(61)	679	(75)	320	F
	51-70	277	466	(22)	266	(46) ^E	300	(42)	365	(34)	451	(27)	554	(34)	663	(57)	736	(78)	320	F
	>70	136	391	(25)	185	(31)	216	(30)	283	(26)	373	(26)	470	(34)	572	(52)	647	(69)	320	34.8 (8.1) ^E
	19+	971	510	(17)	318	(36)	353	(32)	417	(26)	501	(20)	604	(28)	715	(50)	792	(68)	320	F
Female																				
	9-13	265	394	(19)	300	(17)	315	(18)	341	(18)	371	(19)	402	(21)	432	(22)	450	(23)	250	<3
	14-18	290	424	(23)	314	(50)	335	(43)	372	(37)	417	(25)	466	(46)	513	(64)	543	(79)	330	F
	19-30	197	428	(34)	284	(29)	313	(31)	363	(35)	422	(40)	485	(45)	545	(50)	581	(53)	320	F
	31-50	312	408	(19)	308	(41)	326	(35)	357	(27)	394	(22)	435	(30)	474	(45)	498	(57)	320	F
	51-70	312	391	(19)	248	(40)	272	(36)	318	(29)	378	(24)	448	(29)	520	(45)	569	(59)	320	F
	>70	239	312	(17)	253	(45) ^E	266	(39)	290	(29)	319	(17)	349	(40)	379	(96) ^E	F	320	51.1 (13.7) ^E	
	19+	1060	393	(11)	265	(28)	289	(24)	333	(17)	387	(13)	446	(20)	506	(33)	544	(43)	320	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.8 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	249 (13)	135 (20)	154 (18)	188 (16)	233 (17)	285 (22)	341 (33)	379 (43)	120	F	
	4-8	213	395 (19)	251 (32)	278 (28)	327 (22)	387 (20)	455 (28)	522 (43)	565 (54)	160	<3	
Male	9-13	122	506 (60)	302 (54) ^E	342 (52)	416 (52)	509 (59)	622 (80)	746 (110)	829 (131)	250	F	
	14-18	150	624 (44)	400 (69) ^E	440 (64)	515 (55)	615 (53)	733 (71)	855 (108)	935 (137)	330	F	
	19-30	106	595 (64)	282 (79) ^E	334 (73) ^E	432 (64)	560 (59)	717 (75)	896 (114)	1025 (152)	320	F	
	31-50	155	530 (31)	321 (62) ^E	356 (54)	423 (41)	512 (35)	616 (53)	721 (84)	788 (106)	320	F	
	51-70	122	479 (40)	261 (52) ^E	301 (46)	373 (40)	473 (42)	611 (75)	799 (159) ^E	968 (261) ^E	320	F	
	>70	88	405 (25)	244 (35)	271 (33)	323 (30)	391 (30)	472 (37)	559 (52)	618 (65)	320	F	
	19+	471	517 (23)	262 (22)	305 (21)	385 (19)	494 (20)	635 (32)	795 (54)	910 (74)	320	12.3	(3.1) ^E
	Female	9-13	103	456 (32)	311 (54) ^E	339 (50)	391 (45)	456 (44)	526 (52)	596 (68)	641 (81)	250	F
14-18		142	399 (22)	230 (19)	257 (19)	313 (22)	392 (28)	481 (38)	583 (51)	659 (61)	330	30.2	(7.5) ^E
19-30		111	400 (23)	246 (35)	276 (31)	330 (25)	399 (27)	476 (39)	553 (59)	603 (75)	320	F	
31-50		146	395 (21)	247 (52) ^E	274 (46) ^E	324 (37)	387 (29)	461 (32)	538 (48)	590 (63)	320	F	
51-70		184	398 (20)	369 (74) ^E	375 (61)	385 (37)	395 (25)	407 (51)	417 (85) ^E	423 (108) ^E	320	F	
>70		143	328 (19)	230 (36)	250 (33)	287 (27)	334 (24)	386 (30)	439 (42)	474 (51)	320	F	
19+		584	387 (11)	244 (22)	270 (20)	319 (17)	380 (16)	452 (18)	526 (25)	575 (31)	320	25.5	(6.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.9 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	169	246 (17)	105 (32) ^E	130 (29) ^E	177 (24)	238 (20)	311 (24)	389 (42)	443 (58)	120	F	
	4-8	281	367 (14)	237 (39)	260 (34)	302 (24)	355 (16)	415 (25)	474 (44)	511 (57)	160	<3	
Male													
	9-13	183	486 (28)	322 (49)	357 (46)	417 (38)	487 (29)	562 (50)	643 (105)	702 (147) ^E	250	F	
	14-18	187	571 (29)	446 (73)	474 (65)	523 (51)	581 (39)	641 (44)	696 (64)	731 (81)	330	F	
	19-30	223	560 (29)	529 (127) ^E	537 (105) ^E	552 (67)	568 (35)	584 (60)	599 (112) ^E	608 (148) ^E	320	F	
	31-50	229	500 (29)	392 (66) ^E	420 (60)	468 (50)	527 (42)	590 (47)	651 (67)	689 (83)	320	F	
	51-70	197	415 (25)	226 (49) ^E	260 (45) ^E	330 (36)	419 (32)	510 (40)	593 (58)	646 (72)	320	F	
	>70	72	413 (38)	256 (42)	284 (40)	336 (39)	404 (41)	492 (53)	596 (80)	675 (107)	320	F	
	19+	721	488 (15)	325 (41)	359 (36)	423 (27)	502 (19)	591 (26)	680 (43)	738 (57)	320	F	
Female													
	9-13	165	426 (19)	280 (45)	313 (40)	374 (32)	452 (27)	542 (38)	630 (58)	686 (73)	250	F	
	14-18	206	412 (27)	243 (45) ^E	272 (42)	326 (36)	393 (32)	468 (38)	542 (52)	588 (63)	330	F	
	19-30	191	411 (26)	180 (34) ^E	214 (33)	284 (33)	380 (35)	499 (43)	628 (59)	718 (74)	320	34.4 (9.2) ^E	
	31-50	258	370 (19)	279 (21)	299 (22)	335 (24)	376 (26)	419 (29)	464 (32)	493 (34)	320	F	
	51-70	249	375 (18)	239 (38)	264 (34)	309 (26)	366 (21)	432 (27)	500 (43)	544 (55)	320	F	
	>70	128	280 (14)	170 (20)	192 (18)	232 (17)	277 (18)	328 (25)	390 (36)	435 (43)	320	72.0 (9.2)	
	19+	826	371 (10)	210 (20)	237 (19)	290 (16)	362 (14)	446 (17)	535 (25)	595 (32)	320	35.2 (5.8)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.10 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	317 (17)	193 (33) ^E	217 (30)	261 (25)	316 (23)	379 (29)	443 (42)	485 (52)	120	<3	
	4-8	321	387 (14)	244 (22)	270 (20)	316 (16)	375 (16)	443 (23)	516 (37)	566 (49)	160	<3	
Male													
	9-13	226	473 (23)	299 (41)	332 (37)	392 (31)	468 (28)	557 (37)	652 (57)	716 (74)	250	F	
	14-18	262	595 (39)	399 (79) ^E	437 (71)	507 (58)	593 (48)	688 (56)	780 (81)	838 (101)	330	F	
	19-30	197	602 (40)	465 (77)	489 (68)	531 (54)	582 (45)	637 (56)	690 (85)	723 (109)	320	F	
	31-50	282	603 (42)	411 (61)	448 (55)	514 (47)	594 (45)	679 (58)	759 (78)	808 (92)	320	F	
	51-70	234	506 (28)	276 (41)	312 (40)	382 (36)	478 (33)	598 (59)	729 (175) ^E	F	320	F	
	>70	119	438 (35)	231 (41) ^E	268 (39)	338 (37)	428 (40)	530 (54)	634 (74)	701 (88)	320	F	
	19+	832	559 (22)	329 (23)	369 (22)	443 (21)	540 (23)	657 (32)	776 (45)	850 (55)	320	F	
Female													
	9-13	226	417 (29)	254 (33)	286 (30)	339 (28)	413 (31)	502 (45)	598 (68)	663 (84)	250	F	
	14-18	242	450 (24)	240 (38)	276 (35)	343 (29)	429 (28)	522 (38)	616 (57)	680 (70)	330	F	
	19-30	208	423 (27)	247 (37)	280 (35)	342 (32)	420 (33)	509 (42)	601 (58)	662 (70)	320	F	
	31-50	263	447 (23)	241 (42) ^E	275 (37)	337 (31)	418 (29)	520 (41)	630 (67)	702 (87)	320	F	
	51-70	322	382 (15)	219 (26)	249 (23)	306 (20)	378 (19)	461 (24)	546 (35)	601 (45)	320	29.8	(6.9) ^E
	>70	198	326 (12)	184 (16)	208 (16)	252 (15)	308 (17)	377 (22)	454 (32)	511 (43)	320	55.1	(7.4)
	19+	991	409 (11)	221 (12)	253 (12)	313 (12)	392 (14)	487 (18)	589 (26)	659 (32)	320	27.0	(3.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.11 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	(SE)							
Both																				
	1-3	348	274	(10)	150	(14)	171	(13)	211	(12)	263	(13)	324	(16)	386	(22)	428	(27)	120	<3
	4-8	554	395	(15)	276	(31)	300	(28)	345	(24)	401	(21)	465	(24)	531	(34)	575	(43)	160	<3
Male																				
	9-13	409	467	(18)	368	(35)	386	(31)	418	(24)	455	(21)	495	(25)	534	(35)	558	(43)	250	<3
	14-18	414	543	(24)	375	(46)	415	(42)	486	(35)	573	(32)	675	(40)	785	(60)	859	(77)	330	F
	19-30	311	579	(29)	380	(44)	419	(40)	489	(35)	575	(35)	669	(44)	763	(61)	822	(74)	320	F
	31-50	489	498	(20)	289	(22)	324	(21)	384	(20)	469	(22)	567	(28)	669	(38)	737	(46)	320	F
	51-70	575	444	(18)	281	(27)	309	(24)	361	(21)	428	(20)	509	(26)	597	(39)	657	(51)	320	F
	>70	239	414	(17)	274	(28)	300	(28)	344	(26)	402	(25)	473	(30)	548	(80)	597	(168) ^E	320	F
	19+	1614	491	(12)	289	(14)	323	(13)	386	(13)	469	(13)	568	(16)	675	(23)	748	(29)	320	9.5 (2.3) ^E
Female																				
	9-13	355	387	(14)	271	(24)	295	(22)	338	(19)	391	(18)	449	(22)	506	(29)	543	(34)	250	F
	14-18	410	403	(22)	211	(34)	248	(30)	312	(27)	393	(24)	485	(27)	576	(35)	638	(42)	330	30.2 (7.9) ^E
	19-30	384	359	(17)	244	(29)	266	(27)	305	(24)	354	(23)	409	(28)	464	(36)	500	(44)	320	F
	31-50	585	389	(15)	213	(16)	245	(17)	306	(17)	385	(20)	477	(26)	573	(34)	636	(40)	320	29.3 (5.8) ^E
	51-70	711	392	(16)	222	(19)	251	(18)	304	(17)	374	(17)	462	(24)	561	(37)	630	(49)	320	30.6 (6.1) ^E
	>70	342	329	(15)	182	(15)	205	(15)	250	(16)	312	(18)	385	(25)	466	(35)	525	(43)	320	53.2 (8.1)
19+	2022	376	(9)	219	(9)	247	(9)	301	(10)	370	(11)	452	(14)	540	(19)	600	(23)	320	31.8 (3.6)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.12 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	(SE)
Both													
	1-3	622	252	(11)	106 (16)	131 (14)	178 (13)	238 (12)	312 (15)	392 (23)	447 (31)	120	F
	4-8	919	371	(9)	242 (29)	266 (24)	308 (16)	361 (10)	420 (19)	478 (33)	514 (43)	160	<3
Male													
	9-13	579	485	(20)	336 (38)	367 (34)	422 (27)	489 (22)	564 (30)	641 (47)	693 (62)	250	<3
	14-18	634	585	(20)	330 (34)	378 (32)	469 (28)	584 (27)	711 (33)	838 (47)	922 (59)	330	F
	19-30	578	581	(24)	393 (61)	431 (54)	499 (40)	583 (28)	678 (38)	776 (65)	841 (88)	320	F
	31-50	693	505	(20)	341 (53)	375 (45)	439 (33)	515 (26)	600 (37)	684 (58)	737 (73)	320	F
	51-70	596	438	(17)	227 (28)	268 (26)	343 (21)	431 (21)	537 (26)	659 (42)	749 (62)	320	19.6 (5.2) ^E
	>70	296	406	(22)	232 (23)	262 (23)	318 (22)	392 (24)	484 (31)	590 (47)	668 (64)	320	25.7 (6.9) ^E
	19+	2163	498	(10)	293 (19)	332 (17)	406 (14)	497 (13)	609 (16)	731 (26)	815 (36)	320	8.2 (2.3) ^E
Female													
	9-13	533	424	(14)	342 (38)	362 (33)	397 (25)	438 (19)	482 (23)	523 (34)	549 (42)	250	<3
	14-18	638	412	(18)	222 (23)	254 (23)	316 (21)	395 (22)	487 (26)	582 (33)	644 (39)	330	29.3 (6.6) ^E
	19-30	499	412	(19)	255 (36)	284 (32)	336 (25)	402 (22)	475 (31)	547 (46)	594 (57)	320	F
	31-50	716	382	(13)	285 (36)	305 (31)	340 (24)	382 (18)	429 (21)	477 (33)	507 (43)	320	F
	51-70	745	383	(12)	248 (22)	272 (20)	317 (16)	374 (14)	440 (18)	509 (27)	553 (35)	320	26.4 (6.8) ^E
	>70	510	300	(10)	187 (15)	209 (14)	250 (12)	298 (12)	355 (16)	418 (22)	462 (28)	320	60.8 (5.9)
	19+	2470	379	(7)	226 (13)	254 (12)	306 (10)	371 (9)	447 (11)	526 (16)	579 (21)	320	30.2 (3.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 14.13 Folate (DFE/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	283 (6)	137 (8)	164 (8)	213 (7)	274 (7)	345 (8)	420 (12)	472 (15)	120	2.9	(0.8) ^E
	4-8	3235	396 (6)	250 (10)	277 (9)	327 (8)	390 (7)	462 (10)	537 (15)	587 (19)	160	<3	
Male													
	9-13	2080	490 (10)	344 (18)	372 (16)	424 (13)	488 (11)	562 (14)	637 (21)	687 (27)	250	<3	
	14-18	2288	571 (11)	328 (17)	373 (16)	459 (15)	570 (14)	697 (18)	827 (24)	915 (30)	330	5.2	(1.4) ^E
	19-30	1804	587 (14)	398 (29)	434 (25)	500 (20)	580 (17)	669 (22)	759 (34)	816 (44)	320	<3	
	31-50	2596	528 (11)	325 (18)	363 (17)	433 (15)	521 (13)	622 (17)	726 (25)	793 (31)	320	^F	
	51-70	2550	485 (8)	274 (12)	311 (11)	380 (10)	471 (10)	577 (12)	690 (19)	768 (25)	320	11.5	(1.9)
	>70	1520	425 (11)	235 (11)	267 (11)	326 (11)	403 (14)	494 (19)	593 (27)	664 (35)	320	23.1	(3.2)
	19+	8470	520 (6)	304 (8)	342 (8)	414 (7)	508 (7)	615 (10)	730 (14)	807 (17)	320	6.8	(1.0)
Female													
	9-13	1980	425 (8)	276 (13)	304 (12)	356 (11)	420 (10)	493 (12)	565 (17)	613 (21)	250	^F	
	14-18	2256	445 (8)	246 (12)	282 (11)	349 (10)	434 (10)	532 (13)	635 (19)	705 (24)	330	20.1	(2.6)
	19-30	1854	415 (10)	260 (13)	287 (12)	338 (11)	401 (12)	472 (15)	542 (21)	588 (25)	320	18.8	(3.7) ^E
	31-50	2686	423 (9)	250 (12)	281 (12)	338 (11)	411 (11)	496 (13)	586 (18)	646 (23)	320	19.6	(3.3) ^E
	51-70	3200	400 (7)	236 (9)	265 (9)	320 (8)	390 (8)	472 (10)	555 (13)	610 (17)	320	25.0	(2.6)
	>70	2610	340 (6)	192 (7)	218 (6)	265 (7)	328 (8)	404 (10)	486 (13)	543 (16)	320	47.0	(2.9)
	19+	10350	405 (5)	235 (5)	265 (5)	321 (5)	393 (5)	477 (7)	566 (10)	625 (12)	320	24.6	(1.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

15. Iron (mg/d): Usual intakes from food

Table 15.1 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad-equacy (SE)	UL ³	% >UL (SE)										
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)									
Both																								
	1-3	79	9.6	(0.6)	6.4	(1.0)	6.9	(0.9)	7.9	(0.8)	9.1	(0.8)	10.4	(0.9)	11.6	(1.1)	12.4	(1.3)	3.0	<3	40	0.0	(0.0)	
	4-8	127	14.0	(0.8)	11.7	(2.2) ^E	12.3	(2.0)	13.3	(1.6)	14.5	(1.3)	15.8	(1.2)	17.0	(1.5)	17.7	(1.9)	4.1	<3	40	<3		
Male																								
	9-13	111	16.3	(0.9)	12.6	(1.5)	13.2	(1.4)	14.5	(1.3)	16.0	(1.2)	17.7	(1.4)	19.3	(1.8)	20.2	(2.1)	5.9	<3	40	<3		
	14-18	107	17.9	(1.6)	11.4	(2.2) ^E	12.6	(2.0)	15.0	(1.8)	18.5	(1.7)	23.1	(2.4)	28.5	(3.9)	32.5	(5.3)	7.7	<3	45	<3		
	19-30	77	16.1	(0.9)	11.3	(1.2)	12.3	(1.2)	14.0	(1.3)	16.1	(1.4)	18.3	(1.4)	20.2	(1.4)	21.2	(1.5)	6.0	<3	45	0.0	(0.0)	
	31-50	145	14.1	(0.8)	8.8	(1.2)	9.7	(1.1)	11.3	(1.0)	13.4	(0.9)	15.9	(1.1)	18.9	(1.7)	21.1	(2.4)	6.0	<3	45	<3		
	51-70	182	13.8	(0.7)	9.7	(0.7)	10.4	(0.7)	11.6	(0.8)	13.2	(0.8)	15.0	(1.0)	16.8	(1.1)	18.0	(1.2)	6.0	<3	45	0.0	(0.0)	
	>70	63	15.1	(1.5)	11.5	(1.6)	12.2	(1.7)	13.5	(1.8)	15.3	(2.1)	17.4	(2.7)	19.9	(3.7) ^E	21.6	(4.6) ^E	6.0	<3	45	<3		
	19+	467	14.5	(0.5)	9.9	(0.7)	10.7	(0.7)	12.2	(0.6)	14.1	(0.6)	16.4	(0.6)	18.8	(0.9)	20.5	(1.2)	6.0	<3	45	0.0	(0.0)	
Female																								
	9-13	96	13.5	(0.8)	8.3	(1.6) ^E	9.3	(1.5)	11.2	(1.3)	13.7	(1.2)	16.7	(1.4)	20.0	(1.8)	22.2	(2.2)	5.7	<3	40	<3		
	14-18	105	12.1	(0.8)	7.7	(1.0)	8.4	(1.0)	9.7	(1.0)	11.6	(1.0)	14.0	(1.2)	16.5	(1.7)	18.2	(2.2)	7.7	14.6	(4.8) ^E	45	<3	
	19-30	91	11.0	(0.7)	8.0	(1.0)	8.6	(0.9)	9.6	(0.9)	10.8	(0.9)	12.2	(1.2)	13.5	(1.7)	14.5	(2.1)	7.7	20.5	(6.0) ^E	45	0.0	(0.0)
	31-50	167	12.2	(0.8)	8.5	(1.3)	9.4	(1.2)	10.9	(1.1)	12.9	(1.0)	15.0	(1.2)	17.1	(1.6)	18.4	(1.9)	7.7	F		45	0.0	(0.0)
	51-70	198	12.4	(0.7)	7.5	(1.5) ^E	8.4	(1.3)	10.1	(1.0)	12.3	(0.8)	14.7	(1.2)	17.1	(1.9)	18.7	(2.5)	5.0	<3		45	<3	
	>70	74	13.2	(1.1)	9.3	(1.7) ^E	10.1	(1.5)	11.5	(1.4)	13.1	(1.4)	14.9	(1.6)	16.8	(2.0)	18.1	(2.4)	5.0	<3		45	<3	
	19+	530	12.1	(0.4)	8.2	(0.8)	9.0	(0.7)	10.5	(0.6)	12.4	(0.6)	14.6	(0.7)	16.8	(0.9)	18.1	(1.1)				45	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.2 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	58	10.2	(1.1)	6.0 (1.2) ^E	6.8 (1.1) ^E	8.1 (1.0)	9.8 (1.1)	11.8 (1.4)	14.1 (2.1)	15.8 (2.7) ^E	3.0	<3	40	<3
	4-8	110	11.1	(0.4)	8.1 (0.8)	8.6 (0.7)	9.5 (0.7)	10.4 (0.7)	11.4 (0.8)	12.4 (0.8)	13.0 (0.9)	4.1	^F	40	0.0 (0.0)
Male															
	9-13	95	14.9	(0.7)	11.6 (1.3)	12.1 (1.2)	13.2 (1.1)	14.4 (1.1)	15.8 (1.2)	17.2 (1.6)	18.1 (1.9)	5.9	<3	40	0.0 (0.0)
	14-18	87	17.0	(1.1)	10.6 (2.3) ^E	12.0 (2.0) ^E	14.6 (1.7)	17.7 (1.6)	21.1 (1.9)	24.4 (2.6)	26.5 (3.2)	7.7	^F	45	<3
	19-30	70	18.1	(1.3)	10.4 (2.6) ^E	11.9 (2.4) ^E	14.8 (2.0)	18.5 (1.8)	22.5 (2.1)	27.0 (3.1)	30.4 (4.3)	6.0	<3	45	<3
	31-50	109	16.0	(1.2)	9.7 (1.7) ^E	10.6 (1.6)	12.3 (1.4)	14.5 (1.4)	17.3 (1.8)	20.4 (2.5)	22.5 (3.1)	6.0	<3	45	<3
	51-70	128	14.1	(0.5)	10.7 (1.2)	11.4 (1.1)	12.6 (0.9)	14.0 (0.8)	15.7 (0.9)	17.4 (1.4)	18.5 (1.8)	6.0	<3	45	<3
	>70	65	12.3	(0.6)	7.7 (0.9)	8.5 (0.9)	9.9 (0.8)	11.8 (0.8)	13.9 (0.9)	16.2 (1.3)	17.7 (1.6)	6.0	^F	45	0.0 (0.0)
	19+	372	15.5	(0.6)	9.3 (0.8)	10.3 (0.8)	12.2 (0.8)	14.7 (0.8)	17.8 (1.0)	21.3 (1.3)	23.9 (1.7)	6.0	<3	45	<3
Female															
	9-13	75	11.4	(0.7)	7.2 (1.0)	8.0 (0.9)	9.4 (0.9)	11.0 (0.9)	12.9 (1.1)	14.8 (1.5)	16.1 (1.8)	5.7	^F	40	0.0 (0.0)
	14-18	81	12.1	(0.9)	6.7 (1.3) ^E	7.8 (1.1)	9.7 (1.0)	11.8 (1.0)	14.0 (1.3)	16.3 (1.8)	17.9 (2.3)	7.7	^F	45	<3
	19-30	101	13.0	(0.9)	9.1 (1.2)	10.0 (1.1)	11.7 (1.1)	13.7 (1.2)	16.0 (1.6)	18.2 (2.2)	19.6 (2.5)	7.7	10.6 (3.4) ^E	45	0.0 (0.0)
	31-50	116	11.8	(0.7)	6.8 (0.6)	7.5 (0.7)	9.3 (0.7)	12.0 (0.9)	14.2 (1.1)	16.6 (1.5)	18.5 (1.8)	7.7	20.7 (3.9) ^E	45	0.0 (0.0)
	51-70	146	11.9	(0.4)	8.1 (0.8)	8.9 (0.7)	10.2 (0.6)	11.9 (0.6)	13.7 (0.7)	15.6 (0.9)	16.8 (1.1)	5.0	<3	45	0.0 (0.0)
	>70	94	11.1	(0.8)	6.7 (0.9)	7.5 (0.9)	8.9 (0.8)	10.9 (0.9)	13.4 (1.3)	16.1 (1.8)	18.0 (2.3)	5.0	^F	45	<3
	19+	457	12.0	(0.4)	7.8 (0.4)	8.6 (0.4)	10.2 (0.4)	12.1 (0.4)	14.2 (0.6)	16.4 (0.8)	17.8 (1.0)			45	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.3 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	112	9.5	(0.6)	6.3 (0.8)	6.9 (0.8)	8.0 (0.7)	9.3 (0.7)	11.0 (0.9)	12.8 (1.1)	14.0 (1.4)	3.0	<3	40	0.0 (0.0)
	4-8	177	12.4	(0.5)	10.0 (1.3)	10.6 (1.2)	11.6 (0.9)	12.7 (0.7)	13.9 (0.8)	15.0 (1.2)	15.7 (1.5)	4.1	<3	40	<3
Male															
	9-13	111	15.5	(1.0)	11.8 (1.4)	12.4 (1.3)	13.5 (1.1)	14.9 (1.2)	16.5 (1.6)	18.1 (2.2)	19.0 (2.7)	5.9	<3	40	<3
	14-18	113	15.8	(1.3)	11.4 (1.9) ^E	12.3 (1.8)	14.0 (1.7)	16.0 (1.9)	18.3 (2.4)	20.8 (3.4)	22.5 (4.2) ^E	7.7	^F	45	<3
	19-30	91	17.0	(1.1)	10.2 (2.0) ^E	11.5 (1.8)	13.7 (1.5)	16.3 (1.4)	19.0 (1.8)	21.6 (2.5)	23.3 (3.1)	6.0	<3	45	<3
	31-50	101	16.9	(0.9)	11.0 (1.8) ^E	12.1 (1.7)	14.3 (1.4)	16.9 (1.5)	19.8 (1.9)	22.5 (2.6)	24.2 (3.1)	6.0	<3	45	<3
	51-70	134	16.9	(1.1)	11.8 (2.0) ^E	12.9 (1.8)	14.8 (1.5)	17.1 (1.4)	19.7 (1.6)	22.3 (2.3)	23.9 (3.0)	6.0	<3	45	<3
	>70	56	13.3	(1.1)	10.5 (1.5)	11.0 (1.4)	11.9 (1.4)	13.0 (1.5)	14.3 (1.7)	15.5 (2.0)	16.2 (2.3)	6.0	<3	45	0.0 (0.0)
	19+	382	16.6	(0.6)	10.7 (0.8)	11.9 (0.7)	13.9 (0.7)	16.3 (0.7)	19.2 (0.9)	22.1 (1.1)	23.9 (1.4)	6.0	<3	45	<3
Female															
	9-13	105	11.7	(0.6)	8.0 (0.7)	8.7 (0.7)	10.0 (0.7)	11.5 (0.8)	13.1 (0.8)	14.6 (0.9)	15.5 (0.9)	5.7	<3	40	0.0 (0.0)
	14-18	120	11.2	(0.9)	5.3 (1.5) ^E	6.4 (1.4) ^E	8.5 (1.1)	10.7 (0.9)	13.1 (1.1)	15.9 (1.6)	18.1 (2.2)	7.7	25.3 (7.9) ^E	45	<3
	19-30	91	12.7	(0.7)	10.7 (1.6)	11.1 (1.4)	12.0 (1.3)	13.0 (1.1)	14.1 (1.1)	15.2 (1.3)	15.9 (1.5)	7.7	^F	45	0.0 (0.0)
	31-50	159	12.9	(0.5)	6.9 (0.9)	8.0 (0.8)	10.3 (0.7)	12.7 (0.7)	14.9 (0.9)	17.2 (1.2)	18.7 (1.4)	7.7	17.4 (3.5) ^E	45	0.0 (0.0)
	51-70	174	12.0	(0.6)	7.9 (1.0)	8.6 (1.0)	9.9 (0.8)	11.5 (0.7)	13.3 (0.9)	15.2 (1.3)	16.3 (1.6)	5.0	<3	45	0.0 (0.0)
	>70	80	11.8	(0.7)	7.8 (1.2)	8.6 (1.1)	10.0 (1.0)	11.6 (0.9)	13.6 (1.2)	15.8 (1.7)	17.5 (2.2)	5.0	<3	45	<3
	19+	504	12.5	(0.3)	7.7 (0.5)	8.6 (0.5)	10.2 (0.5)	12.2 (0.5)	14.5 (0.6)	16.7 (0.7)	18.1 (0.8)			45	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.4 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	99	11.1	(1.7)	7.0 (1.2) ^E	7.6 (1.2)	8.9 (1.2)	10.6 (1.4)	12.9 (2.0)	15.8 (3.1) ^E	18.0 (4.1) ^E	3.0	<3	40	<3
	4-8	140	13.2	(0.6)	11.1 (1.6)	11.6 (1.4)	12.4 (1.0)	13.3 (0.8)	14.3 (1.0)	15.2 (1.5)	15.8 (1.9)	4.1	<3	40	0.0 (0.0)
Male															
	9-13	92	15.5	(1.1)	10.6 (1.6)	11.5 (1.5)	13.0 (1.4)	14.8 (1.3)	16.8 (1.5)	19.1 (2.2)	20.6 (2.9)	5.9	<3	40	<3
	14-18	107	20.1	(1.4)	14.6 (2.9) ^E	16.0 (2.6)	18.4 (2.1)	21.4 (1.6)	25.1 (2.0)	29.1 (3.3)	32.0 (4.6)	7.7	<3	45	<3
	19-30	73	22.2	(3.2)	11.9 (2.4) ^E	13.6 (2.4) ^E	16.5 (2.6)	20.8 (3.2)	26.8 (4.3)	32.9 (5.6) ^E	36.6 (6.4) ^E	6.0	<3	45	^F
	31-50	134	16.4	(1.3)	11.9 (2.0) ^E	12.5 (1.8)	13.7 (1.6)	15.3 (1.5)	17.0 (1.9)	18.8 (2.9)	20.0 (3.8) ^E	6.0	<3	45	<3
	51-70	131	15.0	(0.9)	11.9 (1.6)	12.5 (1.5)	13.4 (1.2)	14.6 (1.0)	15.9 (1.4)	17.1 (2.3)	17.9 (3.0)	6.0	<3	45	<3
	>70	55	14.2	(1.0)	8.8 (1.4)	9.6 (1.3)	11.3 (1.1)	13.4 (1.1)	15.8 (1.3)	18.4 (1.9)	20.1 (2.4)	6.0	<3	45	<3
	19+	393	17.0	(0.8)	9.7 (1.0)	10.7 (1.0)	12.9 (0.9)	16.0 (0.9)	20.0 (1.3)	24.6 (2.0)	27.9 (2.7)	6.0	<3	45	<3
Female															
	9-13	79	13.9	(1.0)	10.0 (1.5)	10.9 (1.4)	12.3 (1.2)	14.2 (1.2)	16.5 (1.5)	19.1 (2.4)	21.1 (3.3)	5.7	<3	40	<3
	14-18	104	13.1	(0.9)	8.6 (1.3)	9.4 (1.2)	10.7 (1.1)	12.5 (1.1)	14.6 (1.3)	16.7 (1.7)	18.2 (2.1)	7.7	^F	45	0.0 (0.0)
	19-30	101	12.3	(1.0)	8.4 (1.6) ^E	9.3 (1.5)	11.0 (1.5)	13.1 (1.4)	15.4 (1.4)	17.7 (1.6)	19.3 (1.9)	7.7	^F	45	<3
	31-50	143	12.5	(0.8)	7.9 (1.4) ^E	8.7 (1.3)	10.3 (1.0)	12.2 (0.9)	14.3 (1.2)	16.4 (1.8)	17.9 (2.5)	7.7	16.2 (4.9) ^E	45	<3
	51-70	193	12.0	(0.5)	8.4 (1.1)	9.0 (1.0)	10.0 (0.8)	11.3 (0.6)	12.9 (0.8)	14.4 (1.2)	15.5 (1.5)	5.0	<3	45	0.0 (0.0)
	>70	94	11.8	(0.7)	7.0 (1.0)	7.8 (1.0)	9.4 (0.9)	11.4 (1.0)	13.8 (1.2)	16.1 (1.5)	17.5 (1.6)	5.0	^F	45	0.0 (0.0)
	19+	531	12.2	(0.4)	7.5 (0.5)	8.4 (0.5)	10.1 (0.5)	12.1 (0.5)	14.4 (0.6)	16.8 (0.8)	18.5 (1.1)			45	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.5 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy	(SE)	UL ³	% >UL	(SE)								
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)							95th (<i>SE</i>)							
Both																								
	1-3	311	10.1	(0.5)	6.6	(0.8)	7.3	(0.7)	8.6	(0.6)	10.1	(0.6)	11.8	(0.7)	13.7	(1.0)	15.0	(1.3)	3.0	<3		40	0.0	(0.0)
	4-8	485	13.7	(0.4)	9.2	(1.1)	10.1	(1.0)	11.8	(0.8)	13.9	(0.7)	16.4	(0.9)	19.1	(1.4)	20.9	(1.8)	4.1	<3		40	<3	
Male																								
	9-13	277	17.7	(0.8)	11.9	(1.3)	13.0	(1.2)	15.2	(1.2)	18.0	(1.2)	21.4	(1.4)	25.0	(1.9)	27.4	(2.4)	5.9	<3		40	<3	
	14-18	339	21.2	(1.0)	12.7	(1.2)	14.3	(1.1)	17.4	(1.1)	21.6	(1.2)	26.6	(1.5)	32.4	(2.3)	36.6	(3.2)	7.7	<3		45	F	
	19-30	237	17.7	(0.7)	14.5	(1.8)	15.3	(1.6)	16.6	(1.2)	18.2	(1.1)	19.8	(1.4)	21.4	(2.0)	22.4	(2.5)	6.0	<3		45	<3	
	31-50	423	17.3	(0.7)	11.6	(1.3)	12.7	(1.2)	14.7	(1.0)	17.2	(0.8)	19.8	(1.0)	22.5	(1.5)	24.1	(1.9)	6.0	<3		45	<3	
	51-70	387	15.0	(0.5)	9.6	(0.7)	10.5	(0.7)	12.2	(0.5)	14.4	(0.5)	17.0	(0.7)	19.7	(1.2)	21.7	(1.6)	6.0	<3		45	<3	
	>70	132	12.8	(0.8)	7.2	(1.2)	8.1	(1.1)	10.0	(1.0)	12.4	(1.0)	15.5	(1.2)	18.7	(1.4)	20.9	(1.6)	6.0	F		45	<3	
	19+	1179	16.4	(0.4)	10.4	(0.6)	11.4	(0.5)	13.4	(0.5)	16.1	(0.5)	19.1	(0.6)	22.2	(0.8)	24.2	(1.0)	6.0	<3		45	<3	
Female																								
	9-13	281	14.5	(0.8)	10.6	(1.4)	11.3	(1.3)	12.7	(1.1)	14.3	(1.0)	16.2	(1.1)	18.1	(1.4)	19.3	(1.7)	5.7	<3		40	0.0	(0.0)
	14-18	321	12.8	(0.4)	8.6	(0.6)	9.5	(0.5)	11.1	(0.5)	12.9	(0.5)	15.0	(0.6)	17.4	(0.9)	19.1	(1.2)	7.7	9.1	(1.9) ^E	45	<3	
	19-30	249	14.1	(0.7)	12.2	(1.6)	12.7	(1.4)	13.5	(1.1)	14.4	(0.9)	15.4	(1.1)	16.3	(1.5)	16.9	(1.9)	7.7	F		45	0.0	(0.0)
	31-50	364	13.0	(0.4)	8.4	(0.9)	9.2	(0.8)	11.0	(0.7)	13.0	(0.6)	15.2	(0.7)	17.5	(1.1)	19.0	(1.4)	7.7	13.1	(2.3) ^E	45	<3	
	51-70	467	13.5	(0.4)	9.3	(1.0)	10.1	(0.9)	11.6	(0.6)	13.5	(0.5)	15.6	(0.7)	17.7	(1.2)	19.1	(1.5)	5.0	<3		45	0.0	(0.0)
	>70	215	10.8	(0.4)	7.6	(0.8)	8.2	(0.7)	9.5	(0.7)	10.9	(0.7)	12.4	(0.7)	13.9	(0.9)	14.8	(1.1)	5.0	<3		45	0.0	(0.0)
	19+	1295	13.1	(0.3)	8.9	(0.4)	9.8	(0.4)	11.3	(0.4)	13.1	(0.4)	15.2	(0.4)	17.5	(0.6)	18.9	(0.7)				45	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.6 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy	(SE)	UL ³	% >UL	(SE)								
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)							95th (<i>SE</i>)							
Both																								
	1-3	644	9.7	(0.3)	5.0	(0.5)	5.9	(0.5)	7.4	(0.4)	9.4	(0.3)	11.9	(0.4)	14.6	(0.8)	16.7	(1.1)	3.0	F		40	<3	
	4-8	956	12.5	(0.2)	8.4	(0.6)	9.2	(0.5)	10.6	(0.4)	12.4	(0.3)	14.4	(0.4)	16.5	(0.7)	17.8	(0.9)	4.1	<3		40	0.0	(0.0)
Male																								
	9-13	589	16.0	(0.5)	10.3	(1.0)	11.3	(0.9)	13.2	(0.7)	15.6	(0.5)	18.5	(0.8)	21.8	(1.4)	24.1	(2.1)	5.9	<3		40	<3	
	14-18	639	18.1	(0.5)	10.5	(1.1)	11.9	(1.0)	14.5	(0.7)	17.8	(0.6)	21.8	(0.8)	26.2	(1.4)	29.4	(2.0)	7.7	<3		45	<3	
	19-30	481	16.9	(0.6)	12.0	(1.9)	13.0	(1.6)	14.6	(1.1)	16.7	(0.7)	19.0	(1.2)	21.2	(2.2)	22.7	(2.9)	6.0	<3		45	<3	
	31-50	709	15.7	(0.5)	9.1	(0.9)	10.1	(0.8)	12.3	(0.7)	15.3	(0.6)	19.0	(0.7)	23.2	(1.3)	26.0	(1.7)	6.0	<3		45	<3	
	51-70	758	15.0	(0.4)	8.9	(0.7)	10.0	(0.6)	11.9	(0.5)	14.3	(0.4)	17.3	(0.5)	20.7	(0.9)	23.2	(1.3)	6.0	<3		45	<3	
	>70	734	13.5	(0.3)	7.2	(0.6)	8.4	(0.5)	10.5	(0.4)	12.9	(0.4)	15.9	(0.4)	19.2	(0.6)	21.4	(0.8)	6.0	F		45	0.0	(0.0)
	19+	2682	15.6	(0.3)	9.1	(0.5)	10.2	(0.5)	12.3	(0.4)	15.1	(0.4)	18.5	(0.4)	22.3	(0.7)	24.9	(0.9)	6.0	<3		45	<3	
Female																								
	9-13	585	12.9	(0.3)	8.5	(0.7)	9.4	(0.6)	10.9	(0.4)	12.8	(0.3)	14.8	(0.5)	16.8	(0.8)	18.1	(1.0)	5.7	<3		40	0.0	(0.0)
	14-18	645	13.6	(0.5)	8.8	(1.0)	9.7	(0.9)	11.4	(0.7)	13.4	(0.5)	16.0	(0.7)	18.8	(1.4)	20.8	(2.0)	7.7	7.7	(2.4) ^E	45	<3	
	19-30	514	11.6	(0.4)	6.2	(0.7)	7.0	(0.6)	8.6	(0.5)	10.8	(0.4)	13.3	(0.6)	15.9	(1.0)	17.6	(1.4)	7.7	25.7	(3.2)	45	0.0	(0.0)
	31-50	758	11.9	(0.3)	6.9	(0.5)	7.8	(0.4)	9.5	(0.4)	11.6	(0.3)	14.1	(0.5)	16.7	(0.7)	18.6	(1.0)	7.7	20.2	(2.1)	45	0.0	(0.0)
	51-70	955	12.1	(0.3)	6.8	(0.5)	7.7	(0.5)	9.4	(0.4)	11.5	(0.4)	14.1	(0.5)	17.0	(0.8)	19.0	(1.0)	5.0	F		45	0.0	(0.0)
	>70	1345	11.3	(0.2)	6.5	(0.4)	7.3	(0.4)	8.8	(0.3)	10.6	(0.3)	12.9	(0.3)	15.4	(0.6)	17.2	(0.8)	5.0	2.3	(0.8) ^E	45	0.0	(0.0)
	19+	3572	11.8	(0.2)	6.7	(0.2)	7.6	(0.2)	9.3	(0.2)	11.3	(0.2)	13.8	(0.3)	16.5	(0.4)	18.4	(0.5)				45	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.7 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							% Inad- equacy			% >UL							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR ²	(SE)	UL ³	(SE)							
Both																						
	1-3	324	9.5	(0.4)	5.0	(0.8)	5.8	(0.7)	7.4	(0.6)	9.4	(0.5)	11.7	(0.6)	14.0	(1.0)	15.6	(1.3)	3.0	F	40	<3
	4-8	425	12.2	(0.6)	9.8	(0.5)	10.2	(0.5)	11.0	(0.6)	12.0	(0.6)	13.0	(0.7)	14.0	(0.7)	14.7	(0.8)	4.1	<3	40	0.0 (0.0)
Male																						
	9-13	274	15.5	(0.6)	13.0	(1.6)	13.5	(1.4)	14.4	(1.0)	15.5	(0.7)	16.7	(0.8)	17.8	(1.3)	18.4	(1.8)	5.9	<3	40	0.0 (0.0)
	14-18	297	19.1	(0.8)	10.2	(1.2)	11.7	(1.1)	14.5	(1.1)	18.4	(1.0)	23.1	(1.2)	28.2	(1.6)	31.6	(2.0)	7.7	F	45	<3
	19-30	249	18.6	(1.5)	10.8	(0.8)	12.1	(0.9)	14.4	(1.0)	17.5	(1.3)	21.6	(1.8)	26.2	(2.5)	29.4	(3.0)	6.0	<3	45	<3
	31-50	309	14.6	(0.7)	8.2	(1.1)	9.3	(1.0)	11.4	(0.9)	14.2	(0.8)	17.3	(1.1)	20.6	(1.5)	22.7	(1.9)	6.0	<3	45	<3
	51-70	277	15.2	(0.6)	9.5	(1.1)	10.5	(1.0)	12.4	(0.8)	14.9	(0.7)	17.7	(0.9)	20.6	(1.5)	22.6	(2.0)	6.0	<3	45	<3
	>70	136	13.5	(0.9)	6.1	(1.2) ^E	7.3	(1.1)	9.5	(1.0)	12.4	(0.9)	16.4	(1.3)	20.7	(2.1)	23.4	(2.6)	6.0	F	45	<3
	19+	971	15.5	(0.5)	9.1	(0.9)	10.2	(0.8)	12.3	(0.6)	15.0	(0.5)	18.2	(0.7)	21.7	(1.2)	24.2	(1.6)	6.0	<3	45	<3
Female																						
	9-13	265	12.9	(0.6)	7.9	(1.3)	8.8	(1.1)	10.5	(0.8)	12.5	(0.6)	14.6	(0.9)	17.0	(1.5)	18.7	(2.1)	5.7	F	40	<3
	14-18	290	13.1	(1.0)	8.5	(1.5) ^E	9.3	(1.3)	10.8	(1.1)	12.7	(0.9)	15.0	(1.3)	17.5	(2.1)	19.3	(2.8)	7.7	F	45	<3
	19-30	197	13.4	(1.3)	9.1	(1.5)	9.9	(1.4)	11.2	(1.2)	12.9	(1.2)	14.9	(1.6)	16.9	(2.3)	18.3	(2.8)	7.7	F	45	0.0 (0.0)
	31-50	312	12.6	(0.6)	8.7	(1.2)	9.4	(1.1)	10.8	(0.8)	12.5	(0.7)	14.2	(0.8)	16.0	(1.4)	17.2	(1.9)	7.7	13.8 (3.6) ^E	45	0.0 (0.0)
	51-70	312	11.8	(0.5)	7.7	(0.7)	8.4	(0.6)	9.6	(0.5)	11.2	(0.5)	13.1	(0.7)	15.3	(1.0)	16.7	(1.3)	5.0	<3	45	0.0 (0.0)
	>70	239	10.2	(0.5)	5.8	(0.9)	6.6	(0.8)	8.1	(0.6)	10.0	(0.5)	12.2	(0.7)	14.5	(1.2)	16.0	(1.5)	5.0	F	45	0.0 (0.0)
19+	1060	12.2	(0.3)	6.9	(0.5)	7.8	(0.5)	9.6	(0.4)	11.7	(0.3)	14.2	(0.5)	16.9	(0.9)	19.0	(1.2)			45	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.8 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	129	8.6	(0.4)	5.6 (0.6)	6.2 (0.6)	7.4 (0.5)	8.8 (0.5)	10.4 (0.6)	12.0 (0.9)	13.2 (1.2)	3.0	<3	40	0.0 (0.0)
	4-8	213	12.8	(0.4)	9.9 (1.0)	10.6 (0.9)	11.7 (0.6)	12.9 (0.5)	14.2 (0.6)	15.3 (0.9)	16.1 (1.2)	4.1	<3	40	0.0 (0.0)
Male															
	9-13	122	17.6	(1.8)	15.7 (2.2)	16.3 (2.1)	17.3 (1.9)	18.5 (1.9)	19.9 (2.3)	21.1 (3.1)	21.9 (3.7) ^E	5.9	<3	40	<3
	14-18	150	19.2	(1.0)	11.6 (1.3)	13.0 (1.2)	15.6 (1.1)	18.8 (1.1)	22.6 (1.5)	27.0 (2.1)	30.2 (2.8)	7.7	<3	45	<3
	19-30	106	17.6	(1.4)	9.9 (2.0) ^E	11.2 (1.8)	13.7 (1.6)	16.8 (1.6)	20.6 (2.0)	24.7 (3.1)	27.6 (4.0)	6.0	<3	45	<3
	31-50	155	16.7	(1.0)	9.4 (1.8) ^E	10.7 (1.6)	13.3 (1.4)	16.8 (1.2)	20.5 (1.5)	23.9 (2.0)	25.9 (2.3)	6.0	<3	45	<3
	51-70	122	14.9	(0.8)	9.2 (1.4)	10.3 (1.2)	12.4 (0.9)	15.0 (0.9)	18.2 (1.5)	21.7 (2.6)	24.4 (3.7)	6.0	<3	45	<3
	>70	88	15.2	(0.7)	9.2 (0.8)	10.3 (0.8)	12.2 (0.8)	14.6 (0.9)	17.5 (1.0)	20.8 (1.4)	23.1 (2.0)	6.0	<3	45	<3
	19+	471	16.3	(0.5)	8.7 (0.6)	10.1 (0.6)	12.6 (0.6)	15.9 (0.6)	19.9 (0.8)	24.2 (1.2)	27.2 (1.5)	6.0	<3	45	<3
Female															
	9-13	103	14.2	(0.8)	10.6 (1.2)	11.4 (1.1)	12.7 (1.0)	14.3 (1.0)	16.2 (1.2)	18.1 (1.6)	19.4 (1.8)	5.7	<3	40	0.0 (0.0)
	14-18	142	13.0	(0.7)	7.6 (0.6)	8.5 (0.6)	10.3 (0.7)	12.6 (0.9)	15.2 (1.2)	18.2 (1.5)	20.4 (1.9)	7.7	12.6 (3.5) ^E	45	<3
	19-30	111	11.0	(0.5)	7.2 (0.8)	7.9 (0.8)	9.0 (0.6)	10.4 (0.6)	12.0 (0.9)	13.7 (1.2)	14.7 (1.5)	7.7	24.2 (4.3) ^E	45	0.0 (0.0)
	31-50	146	12.5	(0.6)	6.7 (1.4) ^E	7.6 (1.3) ^E	9.3 (1.1)	11.6 (0.9)	14.3 (0.9)	16.8 (1.3)	18.4 (1.6)	7.7	21.0 (5.3) ^E	45	0.0 (0.0)
	51-70	184	12.5	(0.6)	9.6 (1.2)	10.2 (1.0)	11.2 (0.8)	12.4 (0.7)	13.8 (1.0)	15.2 (1.6)	16.1 (2.2)	5.0	<3	45	<3
	>70	143	11.4	(0.5)	7.5 (0.8)	8.3 (0.7)	9.9 (0.7)	11.8 (0.7)	14.0 (0.8)	16.2 (1.1)	17.6 (1.3)	5.0	<3	45	0.0 (0.0)
	19+	584	12.0	(0.3)	7.3 (0.5)	8.1 (0.5)	9.7 (0.5)	11.7 (0.4)	13.9 (0.5)	16.2 (0.7)	17.7 (0.9)			45	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.9 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad-equacy (<i>SE</i>)	UL ³	% >UL (<i>SE</i>)	
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)					95th (<i>SE</i>)
Both															
	1-3	169	9.5	(0.6)	5.2 (0.7)	5.9 (0.6)	7.2 (0.5)	8.9 (0.5)	11.0 (0.8)	13.4 (1.3)	15.2 (1.7)	3.0	F	40	<3
	4-8	281	12.1	(0.4)	9.7 (0.4)	10.1 (0.4)	10.9 (0.4)	11.9 (0.4)	12.9 (0.5)	13.9 (0.5)	14.5 (0.5)	4.1	<3	40	0.0 (0.0)
Male															
	9-13	183	17.1	(1.1)	12.7 (1.6)	13.5 (1.4)	15.1 (1.2)	17.0 (1.1)	19.4 (1.6)	21.8 (2.6)	23.4 (3.4)	5.9	<3	40	<3
	14-18	187	18.6	(0.8)	12.0 (1.7)	13.3 (1.5)	15.7 (1.1)	18.7 (1.0)	21.9 (1.3)	25.1 (2.1)	27.2 (2.6)	7.7	<3	45	<3
	19-30	223	16.8	(0.7)	11.1 (1.7)	12.3 (1.5)	14.4 (1.0)	16.8 (0.8)	19.2 (1.0)	21.4 (1.5)	22.9 (1.9)	6.0	<3	45	0.0 (0.0)
	31-50	229	16.6	(1.0)	11.4 (1.9)	12.4 (1.7)	14.3 (1.3)	16.8 (1.2)	19.7 (1.7)	22.8 (2.7)	24.9 (3.4)	6.0	<3	45	<3
	51-70	197	13.9	(0.6)	8.1 (1.3)	9.2 (1.2)	11.2 (0.9)	13.6 (0.7)	16.1 (0.9)	18.5 (1.4)	20.0 (1.8)	6.0	F	45	0.0 (0.0)
	>70	72	14.9	(0.7)	9.1 (1.6) ^E	10.3 (1.4)	12.5 (1.1)	15.0 (0.9)	17.4 (1.1)	20.1 (1.7)	22.1 (2.3)	6.0	<3	45	<3
	19+	721	15.9	(0.5)	9.9 (0.9)	11.1 (0.8)	13.1 (0.7)	15.8 (0.6)	18.9 (0.8)	22.2 (1.1)	24.5 (1.5)	6.0	<3	45	<3
Female															
	9-13	165	14.4	(0.9)	9.7 (1.5)	10.6 (1.4)	12.4 (1.1)	14.8 (1.0)	17.6 (1.4)	20.6 (2.2)	22.7 (2.8)	5.7	<3	40	<3
	14-18	206	12.7	(0.6)	10.3 (1.4)	10.8 (1.2)	11.7 (0.9)	12.8 (0.7)	13.9 (1.0)	14.9 (1.5)	15.6 (1.9)	7.7	F	45	0.0 (0.0)
	19-30	191	12.1	(0.7)	7.0 (1.0)	7.8 (1.0)	9.5 (0.9)	11.6 (0.9)	14.2 (1.2)	16.9 (1.6)	18.7 (2.0)	7.7	20.1 (5.2) ^E	45	<3
	31-50	258	10.6	(0.4)	8.3 (1.0)	8.8 (0.9)	9.7 (0.7)	10.7 (0.6)	11.9 (0.8)	13.0 (1.1)	13.8 (1.5)	7.7	20.3 (4.3) ^E	45	0.0 (0.0)
	51-70	249	11.8	(0.6)	7.7 (0.9)	8.5 (0.8)	9.8 (0.6)	11.5 (0.6)	13.5 (0.9)	15.6 (1.5)	17.1 (1.9)	5.0	<3	45	0.0 (0.0)
	>70	128	10.7	(0.6)	6.2 (0.9)	7.2 (0.8)	8.8 (0.8)	10.7 (0.8)	13.3 (1.0)	15.7 (1.2)	17.4 (1.5)	5.0	F	45	<3
	19+	826	11.2	(0.3)	7.1 (0.4)	7.8 (0.4)	9.3 (0.3)	11.1 (0.3)	13.2 (0.4)	15.4 (0.6)	16.9 (0.8)			45	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.10 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad-equacy (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	192	9.7	(0.5)	5.7 (0.8)	6.4 (0.8)	7.7 (0.7)	9.3 (0.8)	11.4 (0.9)	13.7 (1.3)	15.0 (1.5)	3.0	F	40	0.0 (0.0)
	4-8	321	12.9	(0.5)	8.1 (0.7)	9.0 (0.6)	10.6 (0.5)	12.7 (0.5)	15.1 (0.7)	17.7 (1.1)	19.4 (1.4)	4.1	<3	40	0.0 (0.0)
Male															
	9-13	226	16.0	(0.6)	10.4 (1.1)	11.6 (1.0)	13.7 (0.9)	16.2 (0.8)	19.1 (1.0)	22.5 (1.5)	25.0 (1.9)	5.9	<3	40	<3
	14-18	262	20.0	(1.2)	12.9 (2.1)	14.2 (1.9)	16.7 (1.6)	19.8 (1.4)	23.3 (1.8)	27.0 (2.6)	29.4 (3.2)	7.7	<3	45	<3
	19-30	197	19.2	(1.1)	10.9 (1.3)	12.2 (1.2)	14.7 (1.2)	18.2 (1.2)	22.8 (1.7)	27.9 (2.6)	31.6 (3.4)	6.0	<3	45	<3
	31-50	282	19.3	(1.0)	10.7 (1.2)	12.4 (1.1)	15.3 (0.9)	18.8 (0.9)	23.0 (1.2)	27.2 (1.9)	30.1 (2.6)	6.0	<3	45	<3
	51-70	234	16.5	(0.7)	9.3 (1.1)	10.5 (1.0)	12.9 (0.9)	15.9 (0.8)	19.3 (1.1)	22.9 (1.5)	25.3 (1.9)	6.0	<3	45	<3
	>70	119	14.2	(0.9)	7.5 (1.1)	8.7 (1.1)	10.9 (1.1)	13.7 (1.1)	16.9 (1.4)	20.0 (1.7)	22.0 (2.0)	6.0	F	45	0.0 (0.0)
	19+	832	18.0	(0.5)	9.9 (0.5)	11.3 (0.5)	13.9 (0.5)	17.4 (0.5)	21.4 (0.7)	25.7 (1.0)	28.9 (1.4)	6.0	<3	45	<3
Female															
	9-13	226	13.1	(0.7)	8.4 (0.6)	9.2 (0.6)	10.7 (0.7)	12.8 (0.8)	15.4 (1.0)	18.1 (1.4)	20.0 (1.7)	5.7	<3	40	<3
	14-18	242	13.0	(0.6)	7.1 (0.7)	8.1 (0.7)	9.9 (0.7)	12.4 (0.7)	15.4 (1.0)	18.6 (1.3)	20.7 (1.6)	7.7	14.5 (3.2) ^E	45	0.0 (0.0)
	19-30	208	12.3	(0.6)	8.6 (1.1)	9.3 (1.0)	10.6 (0.9)	12.1 (0.8)	13.8 (1.0)	15.5 (1.4)	16.6 (1.6)	7.7	15.0 (4.1) ^E	45	0.0 (0.0)
	31-50	263	14.2	(0.8)	7.7 (0.8)	8.8 (0.7)	10.7 (0.7)	13.2 (0.8)	16.4 (1.2)	20.5 (2.2)	23.9 (3.1)	7.7	14.2 (2.6) ^E	45	<3
	51-70	322	11.9	(0.6)	8.0 (1.1)	8.7 (0.9)	10.0 (0.7)	11.7 (0.6)	13.6 (0.9)	15.7 (1.4)	17.1 (1.9)	5.0	<3	45	<3
	>70	198	11.0	(0.4)	6.9 (0.7)	7.6 (0.6)	9.1 (0.6)	10.8 (0.5)	12.7 (0.6)	14.5 (0.8)	15.7 (0.9)	5.0	F	45	0.0 (0.0)
	19+	991	12.8	(0.4)	7.7 (0.4)	8.6 (0.4)	10.2 (0.4)	12.4 (0.4)	15.0 (0.6)	17.8 (1.0)	19.8 (1.3)			45	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.11 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						% Inad- equacy			% >UL		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	EAR ²	(SE)	UL ³	(SE)	
Both																
	1-3	348	10.1	(0.6)	6.1 (0.5)	6.8 (0.5)	8.0 (0.5)	9.6 (0.6)	11.7 (0.8)	14.3 (1.3)	16.3 (1.8)	3.0	<3	40	<3	
	4-8	554	12.9	(0.3)	9.9 (1.0)	10.6 (0.9)	11.7 (0.7)	13.2 (0.5)	14.7 (0.5)	16.1 (0.8)	17.0 (1.0)	4.1	<3	40	<3	
Male																
	9-13	409	15.6	(0.6)	11.3 (1.0)	12.1 (0.9)	13.4 (0.7)	15.1 (0.7)	17.1 (0.9)	19.0 (1.2)	20.3 (1.5)	5.9	<3	40	<3	
	14-18	414	17.7	(0.8)	11.6 (1.1)	12.9 (1.1)	15.2 (1.0)	18.3 (1.0)	22.0 (1.3)	26.0 (1.8)	28.6 (2.4)	7.7	<3	45	<3	
	19-30	311	18.6	(1.1)	11.5 (1.0)	12.8 (1.0)	15.2 (1.0)	18.2 (1.1)	21.8 (1.5)	25.6 (2.1)	28.1 (2.6)	6.0	<3	45	<3	
	31-50	489	16.0	(0.6)	9.7 (0.7)	10.7 (0.7)	12.7 (0.7)	15.2 (0.7)	18.3 (0.9)	21.7 (1.2)	24.0 (1.5)	6.0	<3	45	<3	
	51-70	575	15.4	(0.6)	11.3 (1.2)	12.1 (1.0)	13.5 (0.8)	15.3 (0.7)	17.5 (0.9)	19.6 (1.4)	20.9 (1.7)	6.0	<3	45	0.0 (0.0)	
	>70	239	13.9	(0.7)	9.2 (0.6)	10.0 (0.6)	11.5 (0.7)	13.5 (0.8)	15.8 (1.0)	18.3 (1.2)	20.0 (1.5)	6.0	<3	45	<3	
	19+	1614	16.2	(0.4)	10.0 (0.4)	11.1 (0.4)	13.0 (0.4)	15.6 (0.4)	18.8 (0.6)	22.2 (0.8)	24.6 (1.0)	6.0	<3	45	<3	
Female																
	9-13	355	12.8	(0.4)	8.6 (0.7)	9.5 (0.6)	10.9 (0.6)	12.8 (0.6)	15.0 (0.6)	17.4 (0.9)	19.1 (1.2)	5.7	<3	40	<3	
	14-18	410	12.0	(0.5)	6.4 (0.7)	7.4 (0.6)	9.2 (0.5)	11.4 (0.5)	14.1 (0.6)	17.0 (0.9)	19.0 (1.1)	7.7	19.4 (3.4) ^E	45	0.0 (0.0)	
	19-30	384	12.2	(0.5)	8.8 (0.7)	9.6 (0.7)	10.9 (0.6)	12.7 (0.7)	14.6 (0.8)	16.5 (1.0)	17.8 (1.2)	7.7	13.0 (2.7) ^E	45	0.0 (0.0)	
	31-50	585	12.5	(0.4)	6.9 (0.5)	8.0 (0.5)	10.1 (0.5)	12.6 (0.5)	15.2 (0.6)	17.9 (0.8)	19.7 (1.1)	7.7	17.6 (2.5)	45	0.0 (0.0)	
	51-70	711	12.1	(0.3)	7.6 (0.4)	8.3 (0.4)	9.7 (0.4)	11.6 (0.4)	14.0 (0.6)	16.2 (0.8)	17.6 (0.9)	5.0	<3	45	0.0 (0.0)	
	>70	342	12.0	(0.4)	7.2 (0.6)	8.0 (0.6)	9.6 (0.5)	11.7 (0.6)	14.2 (0.7)	16.8 (0.9)	18.5 (1.1)	5.0	<3	45	0.0 (0.0)	
	19+	2022	12.3	(0.2)	7.7 (0.3)	8.5 (0.3)	10.2 (0.3)	12.2 (0.3)	14.6 (0.4)	17.1 (0.5)	18.6 (0.6)			45	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.12 Iron (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% Inad- equacy (<i>SE</i>)	% UL ³ >UL (<i>SE</i>)										
		n	Mean	(<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)	95th (<i>SE</i>)													
Both																								
	1-3	622	9.4	(0.4)	5.0	(0.5)	5.7	(0.4)	7.2	(0.3)	9.0	(0.4)	11.1	(0.5)	13.7	(0.9)	15.7	(1.2)	3.0	2.2	(0.7) ^E	40	<3	
	4-8	919	12.2	(0.3)	9.8	(0.9)	10.3	(0.8)	11.1	(0.5)	12.1	(0.3)	13.1	(0.5)	14.1	(0.9)	14.7	(1.2)	4.1	<3		40	0.0	(0.0)
Male																								
	9-13	579	16.9	(0.7)	11.9	(0.5)	12.9	(0.5)	14.6	(0.6)	16.9	(0.8)	19.7	(1.1)	22.7	(1.5)	24.7	(1.9)	5.9	<3		40	<3	
	14-18	634	18.8	(0.5)	10.8	(0.7)	12.3	(0.6)	15.1	(0.6)	18.5	(0.7)	22.6	(0.8)	26.8	(1.1)	29.7	(1.4)	7.7	<3		45	<3	
	19-30	578	17.3	(0.6)	10.9	(1.4)	12.2	(1.2)	14.4	(0.9)	17.1	(0.7)	20.0	(0.9)	23.0	(1.5)	25.0	(2.0)	6.0	<3		45	<3	
	31-50	693	16.2	(0.7)	9.7	(1.1)	10.9	(1.0)	13.0	(0.9)	15.9	(0.8)	19.5	(1.0)	23.5	(1.5)	26.2	(1.9)	6.0	<3		45	<3	
	51-70	596	14.4	(0.4)	8.2	(0.6)	9.4	(0.5)	11.5	(0.5)	14.1	(0.5)	17.1	(0.6)	20.3	(0.8)	22.6	(1.1)	6.0	<3		45	<3	
	>70	296	14.6	(0.5)	8.3	(0.7)	9.5	(0.6)	11.7	(0.6)	14.3	(0.6)	17.1	(0.7)	20.1	(0.9)	22.4	(1.2)	6.0	<3		45	<3	
	19+	2163	15.9	(0.3)	9.1	(0.5)	10.4	(0.4)	12.6	(0.4)	15.5	(0.4)	19.0	(0.5)	22.8	(0.7)	25.6	(0.9)	6.0	<3		45	<3	
Female																								
	9-13	533	14.0	(0.6)	9.3	(0.8)	10.2	(0.7)	12.0	(0.6)	14.2	(0.7)	16.7	(0.8)	19.3	(1.2)	21.0	(1.5)	5.7	<3		40	<3	
	14-18	638	12.8	(0.4)	8.1	(0.6)	8.9	(0.5)	10.5	(0.5)	12.6	(0.5)	15.0	(0.6)	17.4	(0.9)	19.1	(1.1)	7.7	11.1	(2.3) ^E	45	0.0	(0.0)
	19-30	499	12.2	(0.5)	7.3	(0.7)	8.1	(0.7)	9.6	(0.6)	11.6	(0.6)	13.9	(0.8)	16.3	(1.1)	17.9	(1.4)	7.7	19.3	(3.3) ^E	45	0.0	(0.0)
	31-50	716	11.3	(0.3)	7.3	(0.8)	8.0	(0.7)	9.4	(0.6)	11.2	(0.5)	13.2	(0.5)	15.3	(0.8)	16.7	(1.1)	7.7	20.8	(3.1)	45	0.0	(0.0)
	51-70	745	11.9	(0.4)	7.9	(0.5)	8.7	(0.5)	10.0	(0.4)	11.6	(0.4)	13.6	(0.6)	15.7	(0.9)	17.1	(1.2)	5.0	<3		45	0.0	(0.0)
	>70	510	10.8	(0.4)	6.0	(0.5)	6.9	(0.5)	8.6	(0.5)	10.8	(0.5)	13.3	(0.6)	15.9	(0.7)	17.6	(0.8)	5.0	^F		45	0.0	(0.0)
	19+	2470	11.6	(0.2)	6.9	(0.3)	7.7	(0.2)	9.3	(0.2)	11.3	(0.2)	13.7	(0.3)	16.2	(0.4)	17.9	(0.6)				45	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 15.13 Iron (mg/d): Usual intakes from food, by DRI age-sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% Inad- equacy	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	2117	9.7	(0.2)	5.5 (0.2)	6.3 (0.2)	7.7 (0.2)	9.5 (0.2)	11.7 (0.3)	14.2 (0.4)	15.9 (0.6)	3.0	1.4	(0.3) ^E	40	0.0 (0.0)
	4-8	3235	12.8	(0.2)	8.7 (0.3)	9.5 (0.3)	10.9 (0.2)	12.7 (0.2)	14.8 (0.3)	16.9 (0.4)	18.3 (0.6)	4.1	0.6	(0.2) ^E	40	0.0 (0.0)
Male																
	9-13	2080	16.5	(0.3)	11.7 (0.5)	12.6 (0.5)	14.2 (0.4)	16.4 (0.4)	19.0 (0.5)	21.8 (0.7)	23.8 (1.0)	5.9	<3		40	<3
	14-18	2288	19.1	(0.4)	11.3 (0.5)	12.7 (0.4)	15.4 (0.4)	18.9 (0.5)	23.3 (0.6)	28.1 (0.8)	31.5 (1.1)	7.7	<3		45	<3
	19-30	1804	17.6	(0.4)	11.5 (0.7)	12.6 (0.6)	14.7 (0.5)	17.3 (0.5)	20.3 (0.7)	23.5 (1.0)	25.7 (1.3)	6.0	<3		45	<3
	31-50	2596	16.7	(0.3)	9.8 (0.4)	10.9 (0.4)	13.2 (0.3)	16.2 (0.4)	19.9 (0.4)	23.9 (0.6)	26.5 (0.9)	6.0	<3		45	<3
	51-70	2550	15.1	(0.2)	9.3 (0.3)	10.3 (0.3)	12.2 (0.3)	14.6 (0.3)	17.5 (0.3)	20.6 (0.5)	22.7 (0.7)	6.0	<3		45	0.0 (0.0)
	>70	1520	13.7	(0.3)	7.7 (0.3)	8.8 (0.3)	10.7 (0.3)	13.2 (0.3)	16.2 (0.4)	19.3 (0.5)	21.4 (0.6)	6.0	1.9	(0.5) ^E	45	0.0 (0.0)
	19+	8470	16.2	(0.2)	9.6 (0.2)	10.7 (0.2)	12.9 (0.2)	15.7 (0.2)	19.1 (0.3)	22.9 (0.4)	25.4 (0.5)	6.0	0.4	(0.1) ^E	45	<3
Female																
	9-13	1980	13.5	(0.3)	8.7 (0.3)	9.6 (0.3)	11.2 (0.3)	13.3 (0.3)	15.7 (0.4)	18.2 (0.5)	19.9 (0.6)	5.7	<3		40	0.0 (0.0)
	14-18	2256	13.1	(0.2)	7.6 (0.2)	8.7 (0.2)	10.5 (0.2)	12.8 (0.2)	15.6 (0.3)	18.9 (0.6)	21.3 (0.8)	7.7	11.9	(1.0)	45	<3
	19-30	1854	12.4	(0.3)	7.8 (0.4)	8.6 (0.3)	10.2 (0.3)	12.0 (0.3)	14.2 (0.4)	16.4 (0.5)	17.9 (0.6)	7.7	16.8	(1.5)	45	0.0 (0.0)
	31-50	2686	12.4	(0.2)	7.2 (0.2)	8.1 (0.2)	9.8 (0.2)	12.1 (0.2)	14.7 (0.3)	17.7 (0.5)	19.7 (0.6)	7.7	18.3	(1.1)	45	0.0 (0.0)
	51-70	3200	12.4	(0.2)	7.9 (0.3)	8.7 (0.3)	10.2 (0.2)	12.1 (0.2)	14.4 (0.3)	16.8 (0.4)	18.5 (0.6)	5.0	<3		45	0.0 (0.0)
	>70	2610	11.1	(0.2)	6.6 (0.2)	7.4 (0.2)	9.0 (0.2)	10.8 (0.2)	13.0 (0.3)	15.3 (0.3)	16.9 (0.4)	5.0	2.0	(0.4) ^E	45	0.0 (0.0)
	19+	10350	12.3	(0.1)	7.4 (0.1)	8.2 (0.1)	9.9 (0.1)	11.9 (0.1)	14.3 (0.2)	17.0 (0.3)	18.8 (0.3)				45	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. Comparisons to the EAR are determined using the probability approach. For additional detail, see Appendix B.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

16. Linoleic acid (g/d): Usual intakes from food

Table 16.1 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	5.6 (0.6)	3.2 (0.7) ^E	3.6 (0.7) ^E	4.3 (0.6)	5.1 (0.7)	6.1 (0.9)	7.1 (1.3) ^E	7.8 (1.5) ^E	7	F	
	4-8	127	10.0 (0.7)	6.5 (1.2) ^E	7.2 (1.1)	8.5 (1.0)	10.2 (1.0)	12.1 (1.2)	13.8 (1.5)	14.9 (1.7)	10	F	
Male	9-13	111	13.7 (0.9)	8.3 (1.5) ^E	9.3 (1.4)	11.3 (1.3)	13.9 (1.3)	16.9 (1.7)	20.1 (2.2)	22.1 (2.7)	12	68.0	(13.1) ^E
	14-18	107	12.8 (0.9)	6.8 (1.8) ^E	7.9 (1.7) ^E	9.9 (1.4)	12.7 (1.2)	16.1 (1.4)	19.8 (2.3)	22.3 (3.1)	16	F	
	19-30	77	12.1 (1.2)	6.6 (1.1)	7.5 (1.1)	9.1 (1.3)	11.2 (1.5)	13.7 (1.9)	16.4 (2.3)	18.1 (2.5)	17	F	
	31-50	145	12.3 (0.9)	6.6 (0.9)	7.6 (0.9)	9.5 (0.9)	11.9 (1.0)	14.8 (1.4)	17.8 (1.8)	19.6 (2.1)	17	F	
	51-70	182	9.9 (0.7)	6.0 (0.6)	6.6 (0.7)	7.7 (0.7)	9.0 (0.8)	10.6 (0.9)	12.1 (1.0)	13.0 (1.0)	14	F	
	>70	63	10.9 (0.7)	7.8 (1.5) ^E	8.6 (1.3)	10.2 (1.1)	12.0 (1.1)	14.0 (1.3)	15.9 (1.8)	17.0 (2.2)	14	F	
	19+	467	11.5 (0.5)	6.3 (0.7)	7.2 (0.7)	8.9 (0.6)	11.1 (0.6)	13.6 (0.8)	16.2 (1.2)	17.8 (1.4)			
	Female	9-13	96	9.7 (0.6)	6.9 (1.0)	7.6 (0.9)	8.7 (0.8)	10.1 (0.8)	11.7 (0.9)	13.3 (1.3)	14.3 (1.6)	10	52.1
14-18	105	11.0 (1.1)	6.2 (1.6) ^E	6.9 (1.5) ^E	8.3 (1.5) ^E	10.4 (1.5)	12.8 (1.7)	15.1 (2.3)	16.6 (2.7)	11	F		
19-30	91	9.3 (1.2)	6.5 (0.9)	7.0 (0.9)	7.9 (1.1)	9.0 (1.2)	10.2 (1.5)	11.5 (1.7)	12.4 (2.0)	12	F		
31-50	167	9.0 (0.7)	4.3 (0.9) ^E	5.1 (0.9) ^E	6.5 (0.9)	8.7 (0.9)	11.7 (1.2)	15.0 (1.8)	17.2 (2.3)	12	F		
51-70	198	7.7 (0.5)	3.1 (0.6) ^E	3.8 (0.6)	5.2 (0.6)	7.4 (0.7)	10.5 (0.9)	13.7 (1.3)	15.8 (1.6)	11	21.7	(6.1) ^E	
>70	74	8.2 (0.8)	4.4 (0.7)	5.1 (0.7)	6.4 (0.8)	8.3 (1.0)	10.7 (1.4)	13.3 (1.9)	15.1 (2.3)	11	F		
19+	530	8.6 (0.4)	4.2 (0.4)	4.9 (0.4)	6.3 (0.4)	8.3 (0.4)	10.8 (0.6)	13.6 (0.8)	15.4 (1.0)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.2 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	58	5.1 (0.3)	2.8 (0.7) ^E	3.2 (0.6) ^E	3.8 (0.6)	4.6 (0.5)	5.5 (0.5)	6.3 (0.6)	6.8 (0.7)	7	F	
	4-8	110	7.8 (0.4)	5.5 (0.7)	6.0 (0.6)	6.8 (0.5)	7.7 (0.5)	8.6 (0.6)	9.5 (0.9)	10.0 (1.0)	10	F	
Male	9-13	95	10.1 (0.8)	7.0 (1.2)	7.5 (1.1)	8.5 (1.0)	9.7 (1.0)	11.1 (1.3)	12.6 (1.7)	13.5 (2.0)	12	F	
	14-18	87	14.0 (1.2)	9.9 (2.0) ^E	10.8 (1.9) ^E	12.4 (1.6)	14.4 (1.5)	16.6 (1.8)	18.9 (2.5)	20.4 (3.1)	16	F	
	19-30	70	17.2 (1.7)	11.9 (2.3) ^E	13.1 (2.1)	15.5 (2.0)	18.7 (2.1)	22.2 (2.6)	25.4 (3.3)	27.3 (3.8)	17	63.7 (17.8) ^E	
	31-50	109	14.2 (1.2)	10.7 (2.0) ^E	11.6 (1.9)	13.1 (1.8)	15.0 (1.7)	17.1 (1.8)	19.0 (2.3)	20.2 (2.7)	17	F	
	51-70	128	11.2 (0.7)	6.6 (1.5) ^E	7.5 (1.4) ^E	9.2 (1.2)	11.4 (1.1)	14.1 (1.3)	17.0 (1.9)	19.0 (2.8)	14	F	
	>70	65	9.3 (0.7)	5.8 (1.2) ^E	6.4 (1.1)	7.4 (0.9)	8.8 (0.9)	10.4 (1.2)	12.1 (1.8)	13.3 (2.2) ^E	14	F	
	19+	372	13.5 (0.7)	8.0 (0.9)	9.1 (0.9)	11.0 (0.9)	13.7 (0.9)	17.1 (1.1)	20.7 (1.5)	23.1 (1.8)			
	Female	9-13	75	10.7 (1.2)	4.4 (1.0) ^E	5.3 (1.1) ^E	7.2 (1.1)	9.9 (1.3)	13.7 (1.9)	18.2 (2.9)	21.5 (3.8) ^E	10	49.4 (12.6) ^E
14-18	81	10.3 (1.0)	4.5 (1.0) ^E	5.4 (1.0) ^E	7.1 (1.1)	9.6 (1.2)	13.0 (1.7)	17.6 (2.8)	21.4 (4.0) ^E	11	37.9 (11.5) ^E		
19-30	101	9.6 (0.7)	7.4 (1.3) ^E	8.0 (1.2)	9.0 (1.0)	10.3 (1.1)	11.7 (1.5)	13.2 (2.2) ^E	14.1 (2.8) ^E	12	F		
31-50	116	9.8 (0.8)	5.7 (1.1) ^E	6.4 (1.0)	7.9 (0.9)	9.7 (0.9)	11.9 (1.2)	14.0 (1.6)	15.4 (2.0)	12	F		
51-70	146	8.4 (0.6)	5.2 (0.9) ^E	5.7 (0.8)	6.7 (0.7)	8.0 (0.6)	9.5 (0.8)	10.9 (1.2)	11.8 (1.5)	11	F		
>70	94	7.5 (0.8)	4.6 (1.0) ^E	5.1 (0.9) ^E	6.0 (0.8)	7.1 (0.7)	8.4 (0.9)	9.8 (1.4)	10.8 (1.8) ^E	11	F		
19+	457	9.1 (0.4)	5.4 (0.6)	6.0 (0.5)	7.3 (0.5)	9.0 (0.5)	11.1 (0.7)	13.2 (1.0)	14.5 (1.3)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.3 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	112	5.6 (0.3)	3.4 (0.6) ^E	3.8 (0.5)	4.6 (0.5)	5.5 (0.4)	6.6 (0.5)	7.7 (0.7)	8.4 (0.9)	7	F	
	4-8	177	8.4 (0.5)	5.5 (0.5)	6.1 (0.5)	7.2 (0.6)	8.6 (0.6)	10.1 (0.7)	11.5 (0.8)	12.4 (0.9)	10	F	
Male	9-13	111	11.4 (0.6)	8.1 (0.7)	8.8 (0.7)	9.9 (0.8)	11.3 (0.8)	12.9 (0.9)	14.3 (1.0)	15.1 (1.0)	12	F	
	14-18	113	13.9 (1.4)	8.9 (2.2) ^E	9.7 (2.1) ^E	11.4 (2.0) ^E	13.8 (1.9)	16.6 (2.1)	19.5 (2.8)	21.3 (3.4)	16	F	
	19-30	91	12.2 (0.9)	6.4 (1.6) ^E	7.3 (1.5) ^E	9.0 (1.2)	11.3 (1.1)	14.1 (1.6)	17.1 (2.6)	19.1 (3.5) ^E	17	F	
	31-50	101	13.6 (1.1)	5.8 (1.5) ^E	6.8 (1.5) ^E	9.0 (1.5) ^E	12.4 (1.6)	16.4 (1.9)	20.5 (2.4)	23.1 (2.8)	17	F	
	51-70	134	12.0 (1.1)	8.5 (1.1)	9.2 (1.2)	10.5 (1.3)	12.0 (1.4)	13.7 (1.6)	15.2 (1.7)	16.2 (1.8)	14	F	
	>70	56	8.5 (1.0)	4.6 (0.7)	5.2 (0.7)	6.3 (0.9)	7.8 (1.1)	9.6 (1.5)	11.6 (1.9)	12.9 (2.2) ^E	14	F	
	19+	382	12.3 (0.6)	6.1 (0.9)	7.0 (0.9)	9.0 (0.8)	11.6 (0.7)	14.7 (1.0)	17.8 (1.6)	20.0 (2.1)			
	Female	9-13	105	9.4 (0.7)	6.4 (0.6)	7.1 (0.6)	8.2 (0.7)	9.5 (0.8)	10.9 (0.9)	12.4 (1.1)	13.3 (1.2)	10	F
14-18	120	9.5 (1.2)	F	5.4 (1.5) ^E	6.9 (1.4) ^E	8.9 (1.2)	11.4 (1.3)	13.9 (2.0)	15.7 (2.6) ^E	11	F		
19-30	91	10.5 (1.2)	F	6.0 (1.6) ^E	7.6 (1.6) ^E	9.5 (1.6) ^E	11.8 (1.8)	14.4 (2.3)	16.3 (2.8) ^E	12	F		
31-50	159	10.0 (1.0)	5.8 (1.3) ^E	6.5 (1.2) ^E	7.9 (1.1)	9.7 (1.1)	11.7 (1.3)	13.7 (1.7)	15.1 (2.1)	12	F		
51-70	174	10.7 (1.3)	6.5 (1.6) ^E	7.2 (1.6) ^E	8.4 (1.4) ^E	10.0 (1.3)	11.9 (1.5)	13.8 (2.1)	15.1 (2.7) ^E	11	F		
>70	80	7.1 (0.5)	4.3 (0.8) ^E	4.9 (0.8)	6.0 (0.7)	7.3 (0.7)	8.6 (0.9)	10.0 (1.2)	11.0 (1.5)	11	F		
19+	504	9.9 (0.6)	5.0 (0.6)	5.8 (0.6)	7.4 (0.6)	9.6 (0.7)	12.4 (1.0)	15.5 (1.4)	17.7 (1.7)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.4 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	99	6.4 (0.6)	4.0 (0.8) ^E	4.4 (0.8) ^E	5.3 (0.7)	6.3 (0.7)	7.4 (0.8)	8.4 (1.1)	9.1 (1.2)	7	F	
	4-8	140	8.5 (0.4)	6.1 (0.3)	6.5 (0.4)	7.3 (0.4)	8.3 (0.5)	9.4 (0.6)	10.5 (0.7)	11.3 (0.8)	10	F	
Male	9-13	92	15.7 (3.1) ^E	F	8.3 (2.7) ^E	10.6 (2.7) ^E	14.4 (2.9) ^E	19.4 (3.9) ^E	25.1 (5.7) ^E	29.0 (7.2) ^E	12	64.9	(20.6) ^E
	14-18	107	14.5 (1.2)	8.9 (1.9) ^E	9.9 (1.8) ^E	11.8 (1.7)	14.3 (1.7)	17.2 (1.9)	20.5 (2.4)	22.8 (2.9)	16	F	
	19-30	73	14.6 (1.1)	7.8 (1.7) ^E	9.1 (1.5) ^E	11.8 (1.3)	15.3 (1.4)	19.5 (2.1)	23.9 (3.0)	26.7 (3.7)	17	38.9	(12.3) ^E
	31-50	134	13.0 (0.9)	6.0 (1.4) ^E	7.0 (1.3) ^E	9.0 (1.2)	11.7 (1.1)	15.2 (1.4)	19.2 (2.2)	22.0 (2.8)	17	F	
	51-70	131	13.0 (1.4)	8.9 (0.9)	9.6 (1.0)	10.8 (1.1)	12.3 (1.3)	14.2 (1.7)	16.2 (2.2)	17.6 (2.6)	14	F	
	>70	55	10.6 (1.2)	F	5.9 (1.8) ^E	7.7 (1.6) ^E	10.1 (1.5)	13.1 (1.9)	16.3 (2.7) ^E	18.5 (3.5) ^E	14	F	
	19+	393	13.1 (0.6)	6.7 (0.7)	7.7 (0.7)	9.7 (0.7)	12.4 (0.7)	16.1 (1.0)	20.3 (1.5)	23.5 (2.0)			
	Female	9-13	79	10.6 (1.3)	5.6 (1.6) ^E	6.6 (1.6) ^E	8.6 (1.6) ^E	10.9 (1.6)	13.6 (1.7)	16.4 (1.9)	18.3 (2.1)	10	60.5
14-18	104	9.5 (1.1)	5.7 (0.7)	6.3 (0.8)	7.5 (0.9)	9.1 (1.2)	11.2 (1.5)	13.6 (2.0)	15.2 (2.4)	11	F		
19-30	101	10.3 (1.0)	6.7 (1.7) ^E	7.5 (1.6) ^E	9.0 (1.4)	10.9 (1.3)	13.0 (1.5)	15.1 (2.1)	16.6 (2.6)	12	F		
31-50	143	8.2 (0.4)	5.0 (0.4)	5.7 (0.4)	6.8 (0.5)	8.3 (0.6)	9.8 (0.6)	11.4 (0.6)	12.4 (0.7)	12	F		
51-70	193	8.8 (0.5)	5.8 (1.1) ^E	6.3 (1.0)	7.3 (0.8)	8.4 (0.7)	9.7 (0.8)	10.9 (1.1)	11.7 (1.3)	11	F		
>70	94	6.8 (0.4)	3.6 (0.5)	4.1 (0.5)	4.9 (0.5)	6.1 (0.5)	7.6 (0.7)	9.3 (1.0)	10.5 (1.3)	11	F		
19+	531	8.6 (0.3)	5.1 (0.6)	5.8 (0.6)	7.0 (0.5)	8.5 (0.4)	10.3 (0.4)	12.0 (0.7)	13.1 (0.8)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.5 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI	(SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both																					
	1-3	311	5.8	(0.3)	2.9	(0.4)	3.5	(0.4)	4.5	(0.4)	5.7	(0.4)	7.2	(0.6)	8.7	(0.8)	9.6	(0.9)	7	27.9	(8.0) ^E
	4-8	485	9.1	(0.3)	5.4	(0.6)	6.1	(0.6)	7.3	(0.5)	9.0	(0.4)	11.1	(0.6)	13.3	(1.0)	14.8	(1.4)	10	37.3	(7.0) ^E
Male																					
	9-13	277	13.6	(0.8)	7.8	(0.9)	9.0	(0.9)	11.2	(0.9)	14.1	(1.1)	17.5	(1.5)	21.8	(2.1)	25.1	(2.7)	12	68.7	(9.0)
	14-18	339	14.5	(0.7)	9.0	(1.3)	10.0	(1.2)	12.0	(1.1)	14.7	(1.0)	17.9	(1.2)	21.4	(1.8)	23.8	(2.4)	16	38.5	(9.7) ^E
	19-30	237	14.7	(0.8)	11.6	(1.7)	12.3	(1.6)	13.6	(1.4)	15.2	(1.3)	16.9	(1.5)	18.6	(1.9)	19.7	(2.3)	17		F
	31-50	423	13.2	(1.0)	6.6	(1.4) ^E	7.7	(1.3) ^E	9.8	(1.3)	12.9	(1.3)	16.5	(1.5)	20.5	(2.2)	23.2	(2.7)	17		F
	51-70	387	12.3	(0.5)	5.6	(0.6)	6.6	(0.6)	8.5	(0.6)	11.2	(0.6)	14.7	(0.8)	18.7	(1.1)	21.5	(1.5)	14	29.3	(4.7)
	>70	132	10.1	(0.8)	5.8	(1.3) ^E	6.7	(1.2) ^E	8.3	(1.1)	10.4	(1.1)	12.6	(1.3)	14.8	(1.6)	16.2	(1.8)	14		F
	19+	1179	13.0	(0.5)	6.8	(0.6)	7.8	(0.6)	9.9	(0.6)	12.7	(0.6)	16.0	(0.8)	19.6	(1.1)	22.1	(1.3)			
Female																					
	9-13	281	10.6	(0.6)	6.0	(0.8)	6.8	(0.8)	8.3	(0.7)	10.2	(0.7)	12.5	(0.9)	15.2	(1.3)	17.0	(1.7)	10	52.3	(9.6) ^E
	14-18	321	11.4	(0.7)	8.1	(1.5) ^E	8.9	(1.4)	10.4	(1.2)	12.2	(1.2)	14.4	(1.4)	16.5	(1.9)	18.0	(2.3)	11	67.0	(16.9) ^E
	19-30	249	10.1	(0.7)	6.6	(1.1)	7.2	(1.0)	8.2	(0.8)	9.5	(0.7)	10.9	(0.9)	12.3	(1.4)	13.3	(1.8)	12		F
	31-50	364	10.0	(0.7)	6.0	(0.9)	6.7	(0.8)	8.0	(0.8)	9.9	(0.8)	12.5	(1.3)	15.7	(2.5)	18.4	(3.8) ^E	12		F
	51-70	467	9.8	(0.5)	4.8	(0.6)	5.6	(0.6)	7.3	(0.6)	9.5	(0.5)	12.1	(0.7)	14.7	(0.9)	16.5	(1.2)	11	34.1	(6.1) ^E
	>70	215	8.1	(0.6)	5.3	(1.0) ^E	5.8	(1.0) ^E	6.7	(0.9)	7.9	(0.9)	9.4	(1.1)	11.1	(1.5)	12.4	(2.0)	11		F
	19+	1295	9.7	(0.3)	5.5	(0.4)	6.3	(0.4)	7.6	(0.3)	9.4	(0.3)	11.7	(0.5)	14.3	(0.9)	16.3	(1.4)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.6 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI	(SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both																					
	1-3	644	5.4	(0.2)	3.0	(0.5)	3.4	(0.4)	4.2	(0.3)	5.3	(0.2)	6.5	(0.3)	7.7	(0.6)	8.5	(0.8)	7	F	
	4-8	956	8.1	(0.3)	4.8	(0.5)	5.4	(0.5)	6.5	(0.4)	7.9	(0.3)	9.7	(0.4)	11.6	(0.7)	12.9	(1.0)	10	21.6	(4.4) ^E
Male																					
	9-13	589	10.7	(0.3)	6.3	(0.8)	7.0	(0.7)	8.4	(0.6)	10.3	(0.4)	12.6	(0.6)	15.1	(1.0)	16.8	(1.4)	12	30.6	(5.9) ^E
	14-18	639	12.8	(0.4)	8.8	(1.4)	9.6	(1.2)	11.1	(0.8)	13.0	(0.5)	15.0	(0.8)	17.0	(1.4)	18.3	(1.9)	16	F	
	19-30	481	13.4	(0.7)	6.7	(1.2) ^E	7.9	(1.1)	10.0	(0.9)	12.8	(0.8)	16.3	(1.1)	20.4	(1.9)	23.3	(2.8)	17	21.6	(6.3) ^E
	31-50	709	11.6	(0.5)	6.8	(1.4) ^E	7.6	(1.2)	9.1	(0.9)	11.0	(0.6)	13.4	(0.9)	15.9	(1.8)	17.6	(2.4)	17	F	
	51-70	758	11.5	(0.6)	5.4	(1.2) ^E	6.3	(1.1) ^E	8.1	(0.9)	10.6	(0.7)	14.0	(0.7)	18.0	(1.4)	20.9	(2.1)	14	24.8	(4.9) ^E
	>70	734	8.9	(0.3)	4.1	(0.7) ^E	4.9	(0.7)	6.4	(0.5)	8.4	(0.4)	10.9	(0.5)	13.8	(0.9)	15.9	(1.3)	14	9.5	(3.1) ^E
	19+	2682	11.8	(0.3)	5.7	(0.5)	6.6	(0.5)	8.4	(0.4)	11.0	(0.4)	14.3	(0.5)	18.1	(0.8)	20.8	(1.1)			
Female																					
	9-13	585	9.5	(0.3)	5.1	(0.6)	5.8	(0.6)	7.2	(0.5)	9.1	(0.4)	11.4	(0.5)	13.9	(0.9)	15.7	(1.3)	10	39.2	(5.1)
	14-18	645	9.9	(0.3)	5.3	(0.7)	6.1	(0.7)	7.6	(0.5)	9.6	(0.4)	12.1	(0.6)	14.6	(1.1)	16.4	(1.4)	11	34.6	(5.3)
	19-30	514	8.6	(0.4)	4.7	(0.9) ^E	5.4	(0.8)	6.5	(0.6)	8.1	(0.5)	10.0	(0.7)	12.0	(1.2)	13.4	(1.7)	12	F	
	31-50	758	9.5	(0.4)	4.7	(0.5)	5.4	(0.5)	6.9	(0.4)	9.0	(0.4)	11.5	(0.5)	14.3	(0.9)	16.2	(1.3)	12	21.7	(4.3) ^E
	51-70	955	8.7	(0.4)	4.7	(0.8)	5.3	(0.7)	6.6	(0.6)	8.3	(0.5)	10.5	(0.5)	13.0	(0.9)	14.6	(1.3)	11	21.1	(5.1) ^E
	>70	1345	7.8	(0.2)	4.2	(0.5)	4.8	(0.5)	5.9	(0.4)	7.5	(0.3)	9.3	(0.4)	11.3	(0.7)	12.7	(1.0)	11	11.8	(3.7) ^E
19+	3572	8.9	(0.2)	4.5	(0.3)	5.2	(0.3)	6.6	(0.3)	8.4	(0.2)	10.7	(0.3)	13.3	(0.5)	15.0	(0.7)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.7 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake											AI ²	% >AI	(SE)						
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both																					
	1-3	324	5.2	(0.3)	2.4	(0.5) ^E	2.9	(0.5)	3.8	(0.4)	4.9	(0.4)	6.2	(0.4)	7.7	(0.7)	8.8	(0.9)	7	F	
	4-8	425	8.4	(0.3)	4.8	(0.9) ^E	5.4	(0.8)	6.5	(0.6)	8.1	(0.4)	10.0	(0.5)	12.1	(1.1)	13.7	(1.6)	10	24.7	(7.4) ^E
Male																					
	9-13	274	11.5	(0.7)	6.0	(1.4) ^E	6.9	(1.2) ^E	8.7	(1.0)	11.1	(0.8)	14.1	(1.0)	17.4	(1.7)	19.9	(2.4)	12	41.1	(9.6) ^E
	14-18	297	14.4	(0.9)	6.8	(1.6) ^E	8.2	(1.4) ^E	10.7	(1.2)	13.9	(1.1)	17.8	(1.4)	22.2	(2.3)	25.4	(3.1)	16	34.9	(8.5) ^E
	19-30	249	15.5	(1.2)	8.9	(0.7)	10.0	(0.8)	12.1	(1.1)	15.3	(1.4)	19.4	(1.7)	23.7	(2.2)	26.6	(2.6)	17	38.3	(10.8) ^E
	31-50	309	11.4	(0.8)	6.7	(0.7)	7.5	(0.7)	9.1	(0.8)	11.2	(1.0)	13.7	(1.3)	16.5	(1.8)	18.5	(2.2)	17	F	
	51-70	277	10.8	(0.6)	6.6	(1.3) ^E	7.4	(1.2)	8.8	(1.0)	10.6	(0.8)	12.5	(1.0)	14.4	(1.6)	15.7	(2.1)	14	F	
	>70	136	9.2	(0.8)	4.6	(0.8) ^E	5.2	(0.7)	6.5	(0.7)	8.2	(0.7)	10.5	(1.0)	13.0	(1.6)	14.8	(2.1)	14	F	
	19+	971	11.9	(0.5)	6.8	(1.4) ^E	7.7	(1.3) ^E	9.4	(1.0)	11.7	(0.6)	14.6	(0.8)	17.8	(1.7)	20.1	(2.5)			
Female																					
	9-13	265	9.2	(0.4)	5.4	(0.9)	6.0	(0.8)	7.2	(0.6)	8.5	(0.5)	10.1	(0.7)	11.8	(1.1)	12.9	(1.4)	10	F	
	14-18	290	9.8	(0.8)	5.4	(1.3) ^E	6.1	(1.1) ^E	7.6	(0.9)	9.6	(0.7)	12.0	(1.0)	14.6	(1.8)	16.4	(2.4)	11	F	
	19-30	197	9.0	(0.8)	5.4	(0.6)	6.1	(0.6)	7.2	(0.7)	8.7	(0.8)	10.4	(0.9)	12.1	(1.1)	13.2	(1.1)	12	F	
	31-50	312	9.0	(0.5)	5.4	(0.4)	6.0	(0.4)	7.2	(0.5)	8.9	(0.6)	10.8	(0.8)	12.9	(0.9)	14.3	(1.0)	12	F	
	51-70	312	8.1	(0.4)	4.2	(0.6)	4.8	(0.6)	6.0	(0.5)	7.7	(0.5)	9.7	(0.7)	12.0	(1.1)	13.6	(1.4)	11	F	
	>70	239	7.0	(0.4)	4.1	(0.8) ^E	4.6	(0.7)	5.4	(0.6)	6.6	(0.5)	7.9	(0.6)	9.2	(1.0)	10.2	(1.3)	11	F	
19+	1060	8.4	(0.3)	5.1	(0.7)	5.7	(0.6)	6.8	(0.4)	8.2	(0.3)	9.9	(0.5)	11.6	(0.9)	12.8	(1.2)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.8 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	6.0 (0.4)	3.8 (0.9) ^E	4.2 (0.8) ^E	5.1 (0.6)	6.2 (0.5)	7.4 (0.7)	8.5 (0.9)	9.2 (1.2)	7	F	
	4-8	213	10.0 (0.9)	6.7 (1.0)	7.2 (1.0)	8.3 (0.9)	9.6 (0.8)	11.2 (1.0)	12.8 (1.6)	13.9 (2.0)	10	42.8	(13.4) ^E
Male	9-13	122	15.5 (4.5) ^E	7.7 (2.2) ^E	8.9 (2.2) ^E	11.3 (2.4) ^E	14.4 (3.1) ^E	18.1 (4.3) ^E	22.5 (5.9) ^E	25.6 (7.1) ^E	12	F	
	14-18	150	16.3 (1.2)	10.8 (1.0)	11.9 (1.1)	13.9 (1.3)	16.5 (1.7)	19.5 (2.2)	22.5 (2.8)	24.5 (3.3)	16	54.8	(15.8) ^E
	19-30	106	15.3 (1.4)	6.9 (2.3) ^E	8.3 (2.1) ^E	10.9 (1.8) ^E	14.6 (1.7)	19.1 (2.3)	24.5 (3.9)	28.4 (5.4) ^E	17	F	
	31-50	155	14.6 (1.5)	8.8 (2.1) ^E	9.9 (2.0) ^E	12.0 (1.8)	14.5 (1.8)	17.5 (2.3)	20.8 (3.3)	22.9 (4.1) ^E	17	F	
	51-70	122	12.5 (1.2)	6.7 (1.4) ^E	7.8 (1.4) ^E	9.7 (1.2)	12.4 (1.3)	16.0 (1.9)	20.3 (3.2)	23.2 (4.4) ^E	14	F	
	>70	88	11.1 (0.8)	5.7 (0.9)	6.7 (0.9)	8.5 (0.8)	11.0 (0.9)	14.0 (1.4)	17.3 (2.1)	19.7 (2.8)	14	F	
	19+	471	13.8 (0.7)	6.3 (0.8)	7.6 (0.8)	10.0 (0.8)	13.2 (0.8)	17.3 (1.1)	22.4 (1.7)	26.3 (2.3)			
	Female	9-13	103	9.7 (0.7)	7.1 (1.3) ^E	7.6 (1.2)	8.7 (1.1)	10.0 (1.1)	11.5 (1.2)	13.0 (1.5)	14.0 (1.7)	10	F
14-18	142	10.0 (0.5)	5.6 (0.9)	6.4 (0.8)	8.0 (0.7)	9.9 (0.8)	12.0 (1.1)	14.3 (1.6)	16.0 (2.0)	11	35.3	(11.4) ^E	
19-30	111	9.8 (0.7)	5.9 (0.6)	6.5 (0.7)	7.9 (0.8)	9.6 (0.9)	11.8 (1.1)	14.1 (1.4)	15.7 (1.7)	12	F		
31-50	146	8.5 (0.7)	4.6 (0.6)	5.1 (0.6)	6.2 (0.7)	7.6 (0.8)	9.4 (1.0)	11.2 (1.2)	12.4 (1.4)	12	F		
51-70	184	9.8 (0.7)	4.7 (1.2) ^E	5.5 (1.1) ^E	7.1 (1.0)	9.4 (0.8)	12.2 (1.1)	15.4 (1.9)	17.7 (2.6)	11	34.1	(9.3) ^E	
>70	143	8.1 (0.7)	4.6 (1.0) ^E	5.2 (1.0) ^E	6.5 (0.9)	8.2 (1.0)	10.5 (1.3)	13.0 (2.0)	14.7 (2.5) ^E	11	F		
19+	584	9.1 (0.3)	5.0 (0.7)	5.6 (0.7)	6.9 (0.6)	8.6 (0.5)	10.7 (0.6)	13.1 (0.9)	14.8 (1.2)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.9 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	169	5.2 (0.4)	2.7 (0.6) ^E	3.1 (0.6) ^E	3.9 (0.5)	4.9 (0.4)	6.1 (0.5)	7.4 (0.9)	8.3 (1.1)	7	F	
	4-8	281	8.1 (0.3)	5.2 (0.9) ^E	5.7 (0.8)	6.7 (0.6)	7.9 (0.4)	9.1 (0.6)	10.4 (1.1)	11.2 (1.5)	10	F	
Male													
	9-13	183	11.8 (0.8)	7.5 (1.4) ^E	8.2 (1.3)	9.7 (1.1)	11.7 (1.0)	14.2 (1.3)	17.0 (2.1)	18.9 (2.9)	12	47.0 (13.0) ^E	
	14-18	187	13.8 (1.1)	8.2 (1.7) ^E	9.2 (1.5)	11.0 (1.3)	13.4 (1.1)	16.3 (1.5)	19.7 (2.4)	22.1 (3.4)	16	F	
	19-30	223	12.1 (0.7)	5.5 (1.0) ^E	6.6 (1.0)	8.7 (0.9)	11.6 (0.9)	15.2 (1.2)	19.2 (1.8)	21.9 (2.4)	17	F	
	31-50	229	13.8 (1.3)	8.4 (0.9)	9.5 (1.0)	11.6 (1.1)	14.2 (1.4)	17.3 (1.9)	20.7 (2.5)	22.9 (3.0)	17	F	
	51-70	197	10.9 (0.8)	5.3 (1.4) ^E	6.2 (1.3) ^E	8.0 (1.1)	10.6 (1.0)	13.8 (1.3)	17.5 (2.2)	20.1 (3.0)	14	F	
	>70	72	10.6 (1.1)	5.8 (1.3) ^E	6.6 (1.2) ^E	8.2 (1.1)	10.3 (1.1)	12.9 (1.6)	15.6 (2.5)	17.5 (3.3) ^E	14	F	
	19+	721	12.5 (0.6)	6.9 (1.3) ^E	7.9 (1.2)	9.9 (1.0)	12.5 (0.8)	15.7 (0.9)	19.1 (1.4)	21.5 (1.9)			
Female													
	9-13	165	9.4 (0.8)	5.5 (1.4) ^E	6.2 (1.3) ^E	7.6 (1.2)	9.4 (1.1)	11.7 (1.3)	14.2 (2.1)	16.0 (2.8) ^E	10	F	
	14-18	206	9.8 (0.6)	6.1 (1.2) ^E	6.8 (1.1)	8.1 (0.9)	9.6 (0.7)	11.4 (1.0)	13.2 (1.6)	14.4 (2.2)	11	F	
	19-30	191	8.9 (0.6)	3.8 (0.9) ^E	4.6 (0.9) ^E	6.2 (0.8)	8.4 (0.9)	11.2 (1.2)	14.4 (1.8)	16.8 (2.6)	12	F	
	31-50	258	9.8 (0.8)	5.3 (0.5)	6.0 (0.5)	7.4 (0.7)	9.3 (0.9)	11.7 (1.2)	14.3 (1.6)	16.1 (2.0)	12	F	
	51-70	249	8.5 (0.5)	4.0 (0.7) ^E	4.7 (0.7)	6.2 (0.6)	8.1 (0.6)	10.5 (0.7)	13.0 (1.1)	14.8 (1.5)	11	21.1 (6.0) ^E	
	>70	128	6.7 (0.8)	2.8 (0.7) ^E	3.3 (0.7) ^E	4.4 (0.8) ^E	6.1 (1.0)	8.5 (1.3)	11.3 (1.7)	13.4 (2.2)	11	F	
	19+	826	8.9 (0.4)	4.8 (0.7)	5.5 (0.6)	6.9 (0.6)	8.7 (0.5)	10.9 (0.6)	13.3 (0.9)	15.0 (1.2)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.10 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	6.2 (0.4)	4.0 (0.7) ^E	4.3 (0.6)	5.0 (0.6)	5.8 (0.6)	6.8 (0.6)	7.7 (0.9)	8.3 (1.0)	7	F	
	4-8	321	8.9 (0.5)	4.8 (0.3)	5.5 (0.4)	6.8 (0.5)	8.6 (0.6)	10.8 (0.9)	13.2 (1.2)	15.0 (1.5)	10	32.8	(8.8) ^E
Male													
	9-13	226	12.1 (0.7)	6.1 (0.9)	7.1 (0.9)	8.9 (0.9)	11.4 (1.0)	15.1 (1.3)	19.5 (2.0)	22.8 (2.7)	12	45.1	(8.7) ^E
	14-18	262	17.1 (1.9)	10.7 (2.3) ^E	11.9 (2.2) ^E	13.9 (1.9)	16.5 (1.8)	19.5 (2.2)	22.6 (3.2)	24.6 (4.0)	16	F	
	19-30	197	15.4 (1.0)	9.2 (1.6) ^E	10.3 (1.4)	12.4 (1.2)	15.3 (1.3)	18.9 (1.9)	22.9 (3.0)	25.7 (4.0)	17	F	
	31-50	282	15.1 (1.0)	7.2 (1.4) ^E	8.5 (1.3)	11.0 (1.2)	14.5 (1.2)	19.0 (1.6)	24.2 (2.5)	28.0 (3.4)	17	34.6	(8.1) ^E
	51-70	234	11.9 (0.7)	6.0 (1.4) ^E	6.8 (1.3) ^E	8.6 (1.1)	11.0 (0.9)	14.2 (1.2)	17.9 (2.1)	20.7 (3.0)	14	26.3	(8.7) ^E
	>70	119	10.4 (0.8)	4.4 (0.8) ^E	5.3 (0.9)	7.2 (0.9)	9.7 (1.1)	12.9 (1.4)	16.5 (1.8)	18.9 (2.1)	14	F	
	19+	832	13.8 (0.5)	6.9 (0.5)	8.0 (0.5)	10.2 (0.5)	13.4 (0.6)	17.3 (0.8)	21.8 (1.2)	25.1 (1.6)			
	Female												
9-13		226	9.3 (0.6)	6.5 (1.0)	7.0 (0.9)	8.1 (0.8)	9.4 (0.7)	10.9 (0.9)	12.4 (1.2)	13.3 (1.5)	10	F	
14-18		242	10.2 (0.7)	5.1 (1.1) ^E	5.9 (1.0) ^E	7.6 (0.9)	9.9 (0.9)	12.5 (1.0)	15.3 (1.4)	17.2 (1.7)	11	38.3	(09.9) ^E
19-30		208	10.3 (0.7)	4.3 (0.8) ^E	5.2 (0.8)	6.9 (0.8)	9.4 (0.8)	12.5 (1.2)	16.2 (2.0)	19.0 (2.6)	12	28.4	(7.5) ^E
31-50		263	10.8 (0.7)	6.1 (1.1) ^E	6.9 (1.0)	8.5 (0.9)	10.5 (0.8)	13.0 (1.1)	15.7 (1.7)	17.6 (2.3)	12	33.7	(10.4) ^E
51-70		322	8.7 (0.6)	3.8 (0.8) ^E	4.5 (0.8) ^E	5.9 (0.7)	8.0 (0.7)	10.8 (0.9)	13.9 (1.5)	16.2 (1.9)	11	23.6	(7.2) ^E
>70		198	7.4 (0.4)	4.3 (0.7)	4.8 (0.6)	5.7 (0.6)	6.9 (0.5)	8.4 (0.7)	10.0 (0.9)	11.1 (1.2)	11	F	
19+		991	9.6 (0.3)	4.5 (0.3)	5.3 (0.3)	6.9 (0.3)	9.1 (0.4)	11.9 (0.5)	15.1 (0.8)	17.3 (1.0)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.11 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake											AI ²	% >AI	(SE)						
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both	1-3	348	5.8	(0.3)	3.3	(0.4)	3.7	(0.3)	4.6	(0.3)	5.6	(0.3)	6.9	(0.4)	8.1	(0.5)	8.9	(0.6)	7	23.0	(6.7) ^E
	4-8	554	8.7	(0.3)	6.0	(0.6)	6.6	(0.6)	7.6	(0.5)	8.9	(0.4)	10.3	(0.5)	11.7	(0.8)	12.6	(1.0)	10	29.5	(8.7) ^E
Male	9-13	409	13.2	(1.1)	8.5	(1.1)	9.5	(1.0)	11.3	(1.0)	13.4	(1.1)	15.7	(1.5)	18.2	(2.3)	19.8	(2.9)	12	67.2	(13.0) ^E
	14-18	414	13.8	(0.7)	8.5	(0.9)	9.5	(0.9)	11.3	(0.9)	13.8	(0.9)	16.7	(1.1)	19.8	(1.4)	21.9	(1.7)	16	30.1	(8.6) ^E
	19-30	311	13.3	(0.6)	7.5	(0.9)	8.5	(0.9)	10.4	(0.8)	13.0	(0.7)	16.1	(1.0)	19.2	(1.5)	21.4	(1.9)	17	19.6	(6.4) ^E
	31-50	489	13.2	(0.6)	6.3	(0.6)	7.3	(0.6)	9.3	(0.6)	12.2	(0.7)	15.8	(0.9)	19.5	(1.2)	22.0	(1.4)	17	18.9	(4.5) ^E
	51-70	575	11.8	(0.7)	7.9	(1.3) ^E	8.7	(1.2)	10.1	(1.0)	11.7	(0.8)	13.5	(1.0)	15.5	(1.6)	16.9	(2.2)	14		^F
	>70	239	9.7	(0.6)	5.4	(0.7)	6.2	(0.7)	7.7	(0.7)	9.6	(0.8)	12.0	(1.0)	14.4	(1.3)	16.1	(1.6)	14		^F
	19+	1614	12.5	(0.3)	6.3	(0.4)	7.3	(0.4)	9.2	(0.4)	11.8	(0.4)	15.1	(0.5)	18.7	(0.8)	21.1	(1.0)			
	Female	9-13	355	9.9	(0.5)	5.9	(0.7)	6.8	(0.6)	8.2	(0.6)	10.0	(0.7)	12.2	(0.8)	14.5	(1.1)	16.1	(1.4)	10	49.7
14-18	410	9.9	(0.6)	5.1	(0.6)	5.9	(0.6)	7.3	(0.7)	9.3	(0.8)	11.9	(0.9)	14.8	(1.3)	16.7	(1.6)	11	32.6	(8.2) ^E	
19-30	384	10.1	(0.6)	6.7	(1.0)	7.4	(0.9)	8.6	(0.8)	10.2	(0.8)	11.9	(1.0)	13.7	(1.3)	14.9	(1.6)	12		^F	
31-50	585	9.2	(0.5)	5.1	(0.6)	5.9	(0.6)	7.2	(0.5)	9.0	(0.6)	11.2	(0.7)	13.3	(0.9)	14.8	(1.1)	12	18.0	(5.6) ^E	
51-70	711	9.3	(0.6)	4.9	(0.6)	5.6	(0.6)	7.0	(0.6)	8.8	(0.6)	11.0	(0.7)	13.3	(0.9)	14.8	(1.1)	11	25.4	(6.5) ^E	
>70	342	7.2	(0.3)	4.1	(0.4)	4.6	(0.4)	5.7	(0.4)	7.1	(0.4)	8.7	(0.5)	10.5	(0.7)	11.8	(0.9)	11		^F	
19+	2022	9.2	(0.3)	4.9	(0.3)	5.6	(0.3)	7.1	(0.3)	9.0	(0.3)	11.4	(0.4)	13.9	(0.6)	15.5	(0.7)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.12 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI (SE)								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both																					
	1-3	622	5.3	(0.2)	2.7	(0.3)	3.1	(0.3)	4.0	(0.3)	5.0	(0.3)	6.4	(0.4)	7.8	(0.5)	8.8	(0.7)	7	17.1	(4.7) ^E
	4-8	919	8.5	(0.3)	4.5	(0.4)	5.1	(0.4)	6.3	(0.3)	8.0	(0.3)	10.0	(0.4)	12.2	(0.7)	13.8	(1.0)	10	24.7	(4.3) ^E
Male																					
	9-13	579	12.5	(0.9)	7.0	(0.9)	7.8	(0.9)	9.5	(0.9)	11.9	(0.9)	15.1	(1.2)	18.9	(1.8)	21.8	(2.4)	12	48.9	(9.6) ^E
	14-18	634	14.4	(0.7)	9.6	(1.5)	10.5	(1.3)	12.3	(1.1)	14.5	(0.8)	17.1	(1.0)	19.8	(1.7)	21.7	(2.3)	16	34.0	(10.3) ^E
	19-30	578	13.3	(0.6)	7.3	(1.6) ^E	8.4	(1.4) ^E	10.6	(1.1)	13.3	(0.8)	16.5	(1.1)	19.8	(1.9)	22.1	(2.6)	17		^F
	31-50	693	13.5	(0.9)	8.1	(1.7) ^E	9.1	(1.5)	11.0	(1.2)	13.5	(1.0)	16.5	(1.2)	19.7	(2.0)	21.8	(2.6)	17		^F
	51-70	596	11.2	(0.5)	5.4	(0.8)	6.4	(0.8)	8.3	(0.7)	10.9	(0.7)	14.1	(0.8)	17.8	(1.3)	20.4	(1.8)	14	25.8	(5.2) ^E
	>70	296	10.4	(0.6)	5.3	(0.6)	6.1	(0.6)	7.7	(0.6)	9.9	(0.6)	12.6	(0.9)	15.6	(1.4)	17.7	(1.9)	14		^F
	19+	2163	12.6	(0.4)	6.5	(0.7)	7.5	(0.6)	9.6	(0.6)	12.4	(0.5)	15.9	(0.6)	19.8	(0.9)	22.5	(1.2)			
Female																					
	9-13	533	9.4	(0.5)	5.9	(1.0)	6.6	(0.9)	7.8	(0.8)	9.4	(0.7)	11.3	(0.8)	13.2	(1.1)	14.6	(1.4)	10	41.1	(11.3) ^E
	14-18	638	9.8	(0.4)	5.6	(0.6)	6.3	(0.6)	7.8	(0.5)	9.5	(0.5)	11.6	(0.6)	13.9	(1.0)	15.5	(1.3)	11	31.2	(6.8) ^E
	19-30	499	9.0	(0.4)	5.3	(0.9) ^E	6.0	(0.8)	7.2	(0.7)	8.9	(0.6)	10.8	(0.8)	12.8	(1.3)	14.1	(1.7)	12		^F
	31-50	716	9.4	(0.5)	5.2	(0.3)	5.8	(0.3)	7.1	(0.4)	9.1	(0.6)	11.2	(0.7)	13.4	(1.0)	14.9	(1.2)	12	18.2	(5.5) ^E
	51-70	745	8.7	(0.3)	4.3	(0.4)	5.0	(0.4)	6.4	(0.4)	8.3	(0.4)	10.6	(0.5)	13.1	(0.8)	14.9	(1.0)	11	22.0	(4.0) ^E
	>70	510	7.1	(0.4)	3.4	(0.5)	3.9	(0.5)	5.1	(0.5)	6.7	(0.6)	8.8	(0.7)	11.1	(0.9)	12.8	(1.2)	11		^F
	19+	2470	8.9	(0.2)	4.7	(0.4)	5.4	(0.3)	6.7	(0.3)	8.5	(0.3)	10.7	(0.4)	13.2	(0.6)	14.9	(0.8)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 16.13 Linoleic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	5.6 (0.1)	3.0 (0.2)	3.4 (0.2)	4.3 (0.2)	5.4 (0.2)	6.7 (0.2)	8.0 (0.3)	8.9 (0.3)	7	20.5	(2.8)
	4-8	3235	8.5 (0.2)	5.1 (0.2)	5.7 (0.2)	6.8 (0.2)	8.4 (0.2)	10.2 (0.2)	12.2 (0.4)	13.5 (0.5)	10	27.2	(2.7)
Male													
	9-13	2080	12.0 (0.3)	6.8 (0.3)	7.6 (0.3)	9.3 (0.3)	11.7 (0.3)	14.7 (0.5)	18.1 (0.7)	20.5 (1.0)	12	46.8	(3.5)
	14-18	2288	14.1 (0.3)	8.8 (0.6)	9.8 (0.5)	11.7 (0.5)	14.3 (0.4)	17.2 (0.6)	20.4 (0.9)	22.6 (1.3)	16	34.1	(4.2)
	19-30	1804	13.9 (0.4)	8.1 (0.6)	9.2 (0.5)	11.2 (0.5)	13.7 (0.5)	16.9 (0.7)	20.3 (1.0)	22.7 (1.3)	17	24.3	(4.2) ^E
	31-50	2596	12.9 (0.4)	6.2 (0.5)	7.3 (0.5)	9.4 (0.4)	12.3 (0.4)	15.9 (0.6)	19.8 (0.9)	22.5 (1.1)	17	19.6	(2.9)
	51-70	2550	11.7 (0.3)	5.7 (0.4)	6.6 (0.4)	8.4 (0.4)	11.0 (0.4)	14.3 (0.4)	18.0 (0.7)	20.8 (0.9)	14	26.7	(2.5)
	>70	1520	9.7 (0.3)	4.8 (0.3)	5.6 (0.3)	7.2 (0.3)	9.3 (0.3)	11.9 (0.4)	14.7 (0.6)	16.7 (0.7)	14	12.9	(2.2) ^E
	19+	8470	12.5 (0.2)	6.1 (0.2)	7.1 (0.2)	9.2 (0.2)	11.9 (0.2)	15.4 (0.3)	19.4 (0.4)	22.1 (0.6)			
Female													
	9-13	1980	9.7 (0.2)	5.6 (0.3)	6.4 (0.3)	7.7 (0.3)	9.5 (0.3)	11.7 (0.3)	14.0 (0.5)	15.6 (0.6)	10	43.6	(3.7)
	14-18	2256	10.2 (0.2)	5.9 (0.3)	6.7 (0.3)	8.2 (0.3)	10.2 (0.3)	12.6 (0.4)	15.0 (0.5)	16.7 (0.7)	11	40.6	(3.6)
	19-30	1854	9.4 (0.3)	5.1 (0.4)	5.8 (0.3)	7.2 (0.3)	8.8 (0.3)	10.8 (0.4)	13.0 (0.6)	14.4 (0.8)	12	15.4	(3.5) ^E
	31-50	2686	9.8 (0.3)	5.4 (0.4)	6.1 (0.3)	7.6 (0.3)	9.5 (0.3)	11.9 (0.4)	14.4 (0.7)	16.2 (1.0)	12	23.8	(3.3)
	51-70	3200	9.0 (0.2)	4.3 (0.3)	5.0 (0.3)	6.5 (0.2)	8.6 (0.2)	11.1 (0.3)	14.0 (0.5)	15.9 (0.6)	11	26.1	(2.4)
	>70	2610	7.7 (0.2)	4.1 (0.2)	4.6 (0.2)	5.8 (0.2)	7.3 (0.2)	9.3 (0.3)	11.6 (0.5)	13.2 (0.7)	11	12.6	(2.3) ^E
	19+	10350	9.2 (0.1)	4.7 (0.1)	5.5 (0.1)	6.9 (0.1)	8.8 (0.2)	11.2 (0.2)	13.9 (0.3)	15.8 (0.5)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

17. Percentage of total energy intake from linoleic acid

Table 17.1 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	79	3.0 (0.2)	1.9 (0.4) ^E	2.1 (0.3)	2.4 (0.3)	2.8 (0.3)	3.3 (0.3)	3.7 (0.4)	4.0 (0.5)
	4-8	127	4.1 (0.2)	3.2 (0.3)	3.4 (0.3)	3.7 (0.2)	4.1 (0.2)	4.6 (0.3)	5.0 (0.3)	5.2 (0.4)
Male										
	9-13	111	4.7 (0.2)	4.1 (0.4)	4.3 (0.4)	4.5 (0.3)	4.8 (0.3)	5.2 (0.4)	5.5 (0.5)	5.7 (0.6)
	14-18	107	4.2 (0.2)	2.4 (0.4) ^E	2.8 (0.4)	3.4 (0.3)	4.1 (0.2)	4.8 (0.3)	5.5 (0.4)	6.0 (0.5)
	19-30	77	4.2 (0.4)	2.8 (0.5) ^E	3.0 (0.5)	3.4 (0.5)	4.0 (0.4)	4.6 (0.5)	5.2 (0.6)	5.5 (0.7)
	31-50	145	4.3 (0.2)	3.3 (0.4)	3.5 (0.3)	3.9 (0.3)	4.3 (0.3)	4.7 (0.3)	5.1 (0.5)	5.4 (0.5)
	51-70	182	4.0 (0.2)	3.4 (0.2)	3.5 (0.2)	3.7 (0.2)	3.9 (0.2)	4.1 (0.2)	4.3 (0.2)	4.4 (0.2)
	>70	63	4.6 (0.2)	4.1 (0.6)	4.2 (0.5)	4.5 (0.4)	4.8 (0.3)	5.1 (0.4)	5.4 (0.5)	5.5 (0.5)
	19+	467	4.2 (0.1)	3.1 (0.2)	3.3 (0.2)	3.7 (0.2)	4.1 (0.2)	4.6 (0.2)	5.1 (0.3)	5.4 (0.3)
Female										
	9-13	96	4.2 (0.2)	3.6 (0.4)	3.7 (0.3)	4.0 (0.3)	4.3 (0.3)	4.6 (0.3)	4.9 (0.4)	5.1 (0.5)
	14-18	105	4.3 (0.2)	3.3 (0.4)	3.5 (0.4)	3.8 (0.3)	4.2 (0.3)	4.6 (0.4)	5.0 (0.4)	5.2 (0.5)
	19-30	91	4.3 (0.3)	3.6 (0.3)	3.7 (0.3)	4.0 (0.3)	4.3 (0.3)	4.7 (0.4)	5.0 (0.4)	5.2 (0.4)
	31-50	167	4.6 (0.3)	3.8 (0.5)	4.0 (0.4)	4.3 (0.4)	4.6 (0.3)	5.0 (0.4)	5.3 (0.5)	5.5 (0.6)
	51-70	198	4.3 (0.2)	2.7 (0.5) ^E	3.0 (0.4)	3.5 (0.3)	4.3 (0.3)	5.3 (0.4)	6.3 (0.6)	7.1 (0.8)
	>70	74	4.2 (0.2)	3.1 (0.4)	3.4 (0.3)	3.8 (0.2)	4.4 (0.2)	5.0 (0.3)	5.7 (0.5)	6.2 (0.6)
	19+	530	4.4 (0.1)	3.2 (0.3)	3.5 (0.3)	3.9 (0.2)	4.5 (0.2)	5.1 (0.2)	5.7 (0.3)	6.0 (0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.2 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	58	3.1 (0.2)	1.8 (0.4) ^E	2.0 (0.3)	2.3 (0.3)	2.8 (0.2)	3.2 (0.2)	3.6 (0.3)	3.8 (0.3)
	4-8	110	3.8 (0.1)	2.5 (0.2)	2.8 (0.2)	3.2 (0.2)	3.7 (0.2)	4.3 (0.2)	4.8 (0.3)	5.2 (0.3)
Male										
	9-13	95	4.0 (0.2)	3.1 (0.4)	3.2 (0.4)	3.5 (0.3)	3.8 (0.3)	4.2 (0.3)	4.5 (0.4)	4.7 (0.4)
	14-18	87	4.5 (0.4)	3.2 (0.4)	3.4 (0.4)	3.9 (0.3)	4.4 (0.4)	5.0 (0.5)	5.6 (0.7)	6.0 (0.8)
	19-30	70	5.0 (0.3)	4.3 (0.6)	4.5 (0.5)	4.8 (0.4)	5.2 (0.4)	5.6 (0.4)	6.0 (0.6)	6.2 (0.7)
	31-50	109	5.3 (0.3)	4.4 (0.6)	4.6 (0.6)	5.0 (0.5)	5.5 (0.5)	6.0 (0.5)	6.5 (0.6)	6.8 (0.7)
	51-70	128	4.7 (0.3)	3.8 (0.6)	4.0 (0.5)	4.3 (0.4)	4.7 (0.3)	5.1 (0.3)	5.5 (0.5)	5.8 (0.7)
	>70	65	4.7 (0.3)	3.7 (0.5)	3.9 (0.4)	4.2 (0.4)	4.6 (0.5)	4.9 (0.6)	5.3 (0.8)	5.5 (1.0) ^E
	19+	372	5.0 (0.2)	4.0 (0.4)	4.2 (0.4)	4.6 (0.3)	5.1 (0.3)	5.7 (0.3)	6.2 (0.3)	6.5 (0.4)
Female										
	9-13	75	4.5 (0.3)	3.0 (0.3)	3.3 (0.3)	3.8 (0.2)	4.4 (0.3)	5.1 (0.4)	5.9 (0.5)	6.6 (0.7)
	14-18	81	4.4 (0.3)	2.7 (0.5)	3.0 (0.4)	3.6 (0.4)	4.4 (0.4)	5.3 (0.6)	6.5 (0.9)	7.3 (1.2) ^E
	19-30	101	4.2 (0.2)	3.3 (0.3)	3.5 (0.3)	3.8 (0.3)	4.1 (0.3)	4.4 (0.3)	4.7 (0.3)	4.9 (0.3)
	31-50	116	4.8 (0.3)	3.7 (0.4)	3.9 (0.4)	4.3 (0.3)	4.7 (0.3)	5.2 (0.4)	5.7 (0.6)	6.0 (0.7)
	51-70	146	4.4 (0.2)	3.0 (0.2)	3.2 (0.2)	3.7 (0.2)	4.2 (0.3)	4.7 (0.3)	5.2 (0.3)	5.5 (0.4)
	>70	94	4.5 (0.3)	3.0 (0.3)	3.3 (0.3)	3.8 (0.3)	4.4 (0.4)	5.1 (0.4)	5.9 (0.5)	6.4 (0.5)
	19+	457	4.5 (0.2)	3.7 (0.3)	3.9 (0.3)	4.1 (0.2)	4.5 (0.2)	4.8 (0.2)	5.2 (0.3)	5.4 (0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.3 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	112	3.4 (0.1)	2.5 (0.2)	2.7 (0.2)	3.0 (0.2)	3.4 (0.2)	3.8 (0.2)	4.3 (0.3)	4.6 (0.4)
	4-8	177	3.9 (0.2)	2.7 (0.4)	3.0 (0.4)	3.5 (0.3)	4.1 (0.3)	4.7 (0.3)	5.2 (0.4)	5.6 (0.5)
Male										
	9-13	111	4.2 (0.2)	3.4 (0.4)	3.5 (0.3)	3.8 (0.3)	4.2 (0.2)	4.6 (0.3)	4.9 (0.4)	5.1 (0.5)
	14-18	113	4.5 (0.2)	3.7 (0.4)	3.9 (0.4)	4.3 (0.3)	4.6 (0.3)	5.1 (0.3)	5.5 (0.4)	5.7 (0.5)
	19-30	91	3.9 (0.2)	2.7 (0.4)	2.9 (0.4)	3.3 (0.3)	3.8 (0.3)	4.4 (0.4)	5.0 (0.5)	5.4 (0.7)
	31-50	101	4.4 (0.2)	2.8 (0.5)	3.1 (0.4)	3.5 (0.4)	4.1 (0.3)	4.8 (0.4)	5.4 (0.5)	5.8 (0.6)
	51-70	134	4.6 (0.3)	4.1 (0.4)	4.2 (0.4)	4.4 (0.4)	4.7 (0.4)	4.9 (0.5)	5.2 (0.5)	5.3 (0.5)
	>70	56	4.0 (0.3)	3.4 (0.4)	3.5 (0.4)	3.6 (0.3)	3.8 (0.4)	4.0 (0.4)	4.2 (0.6)	4.3 (0.7)
	19+	382	4.3 (0.1)	3.0 (0.3)	3.2 (0.3)	3.7 (0.2)	4.2 (0.2)	4.8 (0.3)	5.3 (0.4)	5.7 (0.5)
Female										
	9-13	105	4.1 (0.2)	3.5 (0.4)	3.7 (0.4)	3.9 (0.3)	4.2 (0.2)	4.4 (0.2)	4.6 (0.4)	4.8 (0.4)
	14-18	120	4.0 (0.3)	2.9 (0.4)	3.1 (0.4)	3.4 (0.4)	3.9 (0.4)	4.4 (0.4)	4.9 (0.4)	5.2 (0.5)
	19-30	91	4.6 (0.5)	3.4 (0.6) ^E	3.6 (0.6) ^E	3.9 (0.6)	4.3 (0.6)	4.7 (0.7)	5.1 (0.7)	5.3 (0.7)
	31-50	159	4.4 (0.2)	3.2 (0.5)	3.4 (0.5)	3.8 (0.4)	4.3 (0.3)	4.8 (0.4)	5.3 (0.5)	5.6 (0.6)
	51-70	174	5.3 (0.4)	3.0 (0.6) ^E	3.4 (0.5)	4.2 (0.4)	5.1 (0.4)	6.2 (0.6)	7.3 (0.9)	8.0 (1.1)
	>70	80	4.1 (0.2)	2.6 (0.3)	2.9 (0.3)	3.3 (0.3)	3.9 (0.2)	4.6 (0.3)	5.3 (0.5)	5.7 (0.7)
	19+	504	4.7 (0.2)	3.0 (0.3)	3.3 (0.3)	3.9 (0.2)	4.6 (0.2)	5.4 (0.3)	6.1 (0.4)	6.6 (0.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.4 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	99	3.6 (0.2)	2.3 (0.4)	2.5 (0.3)	3.0 (0.3)	3.5 (0.3)	4.1 (0.4)	4.6 (0.5)	4.9 (0.5)
	4-8	140	3.8 (0.2)	3.0 (0.3)	3.1 (0.3)	3.4 (0.2)	3.7 (0.2)	4.0 (0.2)	4.3 (0.4)	4.5 (0.4)
Male										
	9-13	92	5.2 (0.7)	3.6 (0.6) ^E	3.9 (0.6)	4.4 (0.6)	5.1 (0.7)	5.9 (0.9)	6.7 (1.2) ^E	7.3 (1.4) ^E
	14-18	107	4.4 (0.4)	3.1 (0.3)	3.3 (0.3)	3.7 (0.4)	4.2 (0.4)	4.7 (0.5)	5.2 (0.6)	5.5 (0.6)
	19-30	73	4.5 (0.4)	3.3 (0.6) ^E	3.5 (0.5)	4.0 (0.5)	4.6 (0.5)	5.1 (0.5)	5.7 (0.6)	6.0 (0.7)
	31-50	134	4.5 (0.3)	2.8 (0.4)	3.1 (0.4)	3.6 (0.3)	4.3 (0.3)	5.1 (0.4)	6.0 (0.5)	6.6 (0.7)
	51-70	131	4.9 (0.3)	3.4 (0.5)	3.6 (0.4)	4.1 (0.4)	4.6 (0.3)	5.2 (0.4)	5.8 (0.6)	6.2 (0.7)
	>70	55	4.4 (0.2)	2.9 (0.6) ^E	3.2 (0.5)	3.8 (0.4)	4.4 (0.3)	5.0 (0.4)	5.5 (0.5)	5.9 (0.5)
	19+	393	4.6 (0.2)	3.0 (0.3)	3.3 (0.2)	3.8 (0.2)	4.5 (0.2)	5.2 (0.2)	6.0 (0.3)	6.5 (0.4)
Female										
	9-13	79	4.4 (0.5)	2.9 (0.5) ^E	3.2 (0.5) ^E	3.7 (0.6)	4.2 (0.6)	4.9 (0.6)	5.5 (0.6)	5.9 (0.6)
	14-18	104	4.2 (0.3)	3.3 (0.3)	3.4 (0.3)	3.8 (0.3)	4.2 (0.4)	4.7 (0.4)	5.2 (0.5)	5.5 (0.6)
	19-30	101	5.2 (0.5)	3.1 (0.7) ^E	3.5 (0.7) ^E	4.2 (0.6)	5.1 (0.6)	6.0 (0.7)	6.9 (0.9)	7.6 (1.1)
	31-50	143	4.5 (0.2)	3.7 (0.3)	3.9 (0.3)	4.2 (0.3)	4.6 (0.3)	5.0 (0.3)	5.3 (0.3)	5.6 (0.3)
	51-70	193	4.8 (0.2)	3.6 (0.2)	3.8 (0.2)	4.2 (0.2)	4.7 (0.3)	5.3 (0.3)	5.8 (0.3)	6.1 (0.3)
	>70	94	4.3 (0.3)	3.0 (0.4)	3.3 (0.4)	3.7 (0.4)	4.3 (0.3)	4.9 (0.4)	5.6 (0.5)	6.0 (0.5)
	19+	531	4.7 (0.2)	3.3 (0.3)	3.6 (0.3)	4.1 (0.2)	4.7 (0.2)	5.3 (0.2)	6.0 (0.3)	6.4 (0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.5 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	311	3.2	(0.1)	1.9	(0.2)	2.2	(0.2)	2.6	(0.2)	3.1	(0.2)	3.7	(0.2)	4.3	(0.3)	4.7	(0.3)
	4-8	485	3.9	(0.1)	2.8	(0.2)	3.0	(0.2)	3.4	(0.1)	3.9	(0.1)	4.5	(0.2)	5.0	(0.3)	5.4	(0.4)
Male																		
	9-13	277	4.3	(0.2)	3.3	(0.3)	3.5	(0.3)	4.0	(0.2)	4.5	(0.2)	5.1	(0.3)	5.7	(0.4)	6.1	(0.5)
	14-18	339	4.0	(0.2)	2.9	(0.3)	3.1	(0.3)	3.5	(0.2)	3.9	(0.2)	4.4	(0.3)	4.9	(0.4)	5.1	(0.4)
	19-30	237	4.6	(0.3)	4.1	(0.5)	4.3	(0.5)	4.5	(0.4)	4.7	(0.3)	5.0	(0.4)	5.3	(0.6)	5.5	(0.8)
	31-50	423	4.2	(0.2)	3.5	(0.4)	3.6	(0.3)	3.9	(0.3)	4.3	(0.2)	4.6	(0.3)	5.0	(0.4)	5.2	(0.5)
	51-70	387	4.6	(0.1)	2.6	(0.2)	3.0	(0.2)	3.6	(0.2)	4.4	(0.2)	5.3	(0.3)	6.3	(0.6)	7.0	(0.8)
	>70	132	4.6	(0.3)	3.1	(0.3)	3.4	(0.3)	3.9	(0.3)	4.6	(0.4)	5.3	(0.4)	6.0	(0.4)	6.5	(0.4)
	19+	1179	4.4	(0.1)	3.2	(0.2)	3.4	(0.2)	3.9	(0.2)	4.4	(0.1)	5.0	(0.2)	5.6	(0.2)	6.0	(0.3)
Female																		
	9-13	281	4.2	(0.1)	3.1	(0.3)	3.3	(0.3)	3.7	(0.2)	4.2	(0.2)	4.8	(0.2)	5.5	(0.4)	5.9	(0.5)
	14-18	321	4.5	(0.2)	3.8	(0.3)	4.0	(0.3)	4.3	(0.3)	4.8	(0.3)	5.2	(0.4)	5.6	(0.4)	5.9	(0.5)
	19-30	249	4.1	(0.2)	3.0	(0.3)	3.2	(0.3)	3.5	(0.2)	3.9	(0.2)	4.3	(0.3)	4.8	(0.4)	5.0	(0.5)
	31-50	364	4.2	(0.2)	3.1	(0.3)	3.3	(0.3)	3.7	(0.2)	4.2	(0.2)	4.8	(0.2)	5.4	(0.4)	5.7	(0.5)
	51-70	467	4.7	(0.2)	3.0	(0.3)	3.4	(0.3)	3.9	(0.2)	4.6	(0.2)	5.3	(0.2)	6.1	(0.4)	6.6	(0.4)
	>70	215	4.6	(0.2)	3.5	(0.5)	3.7	(0.4)	4.1	(0.4)	4.7	(0.3)	5.2	(0.3)	5.7	(0.5)	6.1	(0.6)
	19+	1295	4.4	(0.1)	3.0	(0.2)	3.3	(0.1)	3.8	(0.1)	4.3	(0.1)	4.9	(0.1)	5.6	(0.2)	6.0	(0.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.6 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	644	3.2 (0.1)	2.3 (0.3)	2.5 (0.2)	2.8 (0.1)	3.2 (0.1)	3.6 (0.2)	4.0 (0.3)	4.2 (0.4)
	4-8	956	3.8 (0.1)	2.5 (0.2)	2.7 (0.2)	3.2 (0.1)	3.7 (0.1)	4.4 (0.1)	5.1 (0.2)	5.5 (0.3)
Male										
	9-13	589	4.0 (0.1)	3.3 (0.3)	3.4 (0.3)	3.7 (0.2)	4.0 (0.1)	4.3 (0.2)	4.6 (0.3)	4.8 (0.4)
	14-18	639	4.2 (0.1)	3.5 (0.4)	3.6 (0.3)	3.9 (0.2)	4.2 (0.1)	4.4 (0.2)	4.7 (0.3)	4.9 (0.4)
	19-30	481	4.4 (0.1)	3.0 (0.4)	3.3 (0.3)	3.8 (0.2)	4.3 (0.2)	5.0 (0.2)	5.7 (0.4)	6.1 (0.6)
	31-50	709	4.3 (0.1)	2.8 (0.4)	3.0 (0.3)	3.5 (0.2)	4.0 (0.2)	4.7 (0.2)	5.4 (0.4)	5.9 (0.6)
	51-70	758	4.5 (0.1)	2.9 (0.4)	3.2 (0.3)	3.7 (0.3)	4.4 (0.2)	5.2 (0.2)	6.1 (0.4)	6.6 (0.6)
	>70	734	4.3 (0.1)	2.5 (0.3)	2.9 (0.3)	3.5 (0.2)	4.2 (0.1)	5.1 (0.2)	6.0 (0.3)	6.7 (0.4)
	19+	2682	4.4 (0.1)	2.6 (0.1)	2.9 (0.1)	3.5 (0.1)	4.2 (0.1)	5.0 (0.1)	5.9 (0.2)	6.6 (0.3)
Female										
	9-13	585	4.2 (0.1)	4.0 (0.7)	4.0 (0.5)	4.1 (0.3)	4.3 (0.1)	4.4 (0.3)	4.5 (0.6)	4.6 (0.8) ^E
	14-18	645	4.3 (0.1)	2.9 (0.3)	3.2 (0.3)	3.6 (0.2)	4.2 (0.1)	4.8 (0.2)	5.5 (0.4)	5.9 (0.5)
	19-30	514	4.2 (0.2)	3.0 (0.1)	3.2 (0.1)	3.6 (0.2)	4.1 (0.2)	4.6 (0.2)	5.1 (0.2)	5.4 (0.3)
	31-50	758	4.6 (0.1)	3.0 (0.4)	3.2 (0.3)	3.8 (0.2)	4.5 (0.2)	5.4 (0.3)	6.3 (0.5)	6.9 (0.6)
	51-70	955	4.4 (0.1)	3.3 (0.4)	3.5 (0.4)	3.9 (0.3)	4.4 (0.2)	4.9 (0.2)	5.5 (0.4)	5.8 (0.5)
	>70	1345	4.4 (0.1)	2.8 (0.2)	3.1 (0.2)	3.6 (0.2)	4.3 (0.1)	5.2 (0.2)	6.0 (0.3)	6.5 (0.3)
	19+	3572	4.5 (0.1)	3.1 (0.2)	3.3 (0.2)	3.8 (0.1)	4.4 (0.1)	5.1 (0.1)	5.8 (0.2)	6.2 (0.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.7 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	324	3.3 (0.1)	2.0 (0.3)	2.2 (0.3)	2.6 (0.2)	3.2 (0.1)	3.7 (0.2)	4.3 (0.3)	4.7 (0.4)
	4-8	425	4.0 (0.2)	3.1 (0.4)	3.3 (0.4)	3.6 (0.3)	4.1 (0.2)	4.6 (0.3)	5.1 (0.4)	5.4 (0.6)
Male										
	9-13	274	4.3 (0.2)	2.9 (0.5)	3.1 (0.4)	3.6 (0.3)	4.2 (0.2)	4.9 (0.3)	5.6 (0.4)	6.0 (0.6)
	14-18	297	4.1 (0.1)	3.1 (0.3)	3.3 (0.3)	3.6 (0.2)	4.0 (0.2)	4.4 (0.2)	4.8 (0.3)	5.1 (0.4)
	19-30	249	4.8 (0.2)	4.2 (0.2)	4.3 (0.2)	4.6 (0.2)	4.9 (0.2)	5.3 (0.3)	5.6 (0.3)	5.8 (0.3)
	31-50	309	4.3 (0.2)	3.6 (0.7) ^E	3.8 (0.6)	4.0 (0.4)	4.3 (0.3)	4.7 (0.5)	5.0 (1.0) ^E	5.3 (1.4) ^E
	51-70	277	4.5 (0.2)	3.4 (0.2)	3.7 (0.2)	4.0 (0.2)	4.5 (0.2)	5.0 (0.2)	5.5 (0.3)	5.8 (0.3)
	>70	136	4.6 (0.2)	2.6 (0.4)	2.9 (0.4)	3.6 (0.3)	4.4 (0.3)	5.4 (0.3)	6.4 (0.5)	7.1 (0.6)
	19+	971	4.5 (0.1)	3.4 (0.4)	3.6 (0.3)	4.0 (0.2)	4.5 (0.1)	5.1 (0.2)	5.6 (0.4)	6.0 (0.5)
Female										
	9-13	265	4.1 (0.2)	2.9 (0.2)	3.1 (0.2)	3.5 (0.2)	3.9 (0.2)	4.4 (0.2)	4.9 (0.3)	5.2 (0.3)
	14-18	290	4.3 (0.1)	3.2 (0.4)	3.4 (0.3)	3.8 (0.2)	4.2 (0.2)	4.6 (0.3)	5.1 (0.4)	5.4 (0.5)
	19-30	197	3.8 (0.2)	3.0 (0.2)	3.2 (0.2)	3.5 (0.2)	3.9 (0.2)	4.3 (0.2)	4.7 (0.3)	5.0 (0.3)
	31-50	312	4.3 (0.2)	3.3 (0.2)	3.5 (0.2)	3.9 (0.2)	4.3 (0.2)	4.8 (0.2)	5.3 (0.2)	5.6 (0.3)
	51-70	312	4.4 (0.2)	2.8 (0.3)	3.0 (0.3)	3.5 (0.3)	4.3 (0.3)	5.1 (0.4)	6.0 (0.6)	6.6 (0.7)
	>70	239	4.3 (0.2)	3.1 (0.4)	3.3 (0.3)	3.7 (0.2)	4.1 (0.2)	4.5 (0.3)	4.9 (0.4)	5.1 (0.6)
	19+	1060	4.2 (0.1)	2.9 (0.3)	3.2 (0.2)	3.6 (0.2)	4.2 (0.1)	4.8 (0.2)	5.4 (0.3)	5.8 (0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.8 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	129	3.6 (0.2)	2.3 (0.3)	2.5 (0.3)	3.0 (0.3)	3.6 (0.2)	4.3 (0.3)	5.0 (0.4)	5.5 (0.5)
	4-8	213	4.4 (0.2)	3.2 (0.4)	3.4 (0.3)	3.8 (0.3)	4.3 (0.2)	4.8 (0.3)	5.3 (0.4)	5.7 (0.6)
Male										
	9-13	122	5.0 (0.8)	2.8 (0.4)	3.1 (0.4)	3.7 (0.5)	4.5 (0.6)	5.7 (0.9)	6.9 (1.2) ^E	7.9 (1.4) ^E
	14-18	150	4.7 (0.2)	4.4 (0.3)	4.4 (0.3)	4.6 (0.3)	4.8 (0.3)	5.0 (0.3)	5.1 (0.3)	5.2 (0.3)
	19-30	106	4.6 (0.2)	3.4 (0.6) ^E	3.6 (0.5)	4.0 (0.4)	4.6 (0.3)	5.2 (0.4)	5.7 (0.6)	6.1 (0.7)
	31-50	155	4.9 (0.3)	4.0 (0.3)	4.2 (0.3)	4.6 (0.4)	5.0 (0.4)	5.4 (0.4)	5.8 (0.5)	6.1 (0.6)
	51-70	122	5.1 (0.3)	4.2 (0.3)	4.4 (0.3)	4.8 (0.4)	5.3 (0.4)	5.8 (0.5)	6.3 (0.6)	6.7 (0.6)
	>70	88	4.8 (0.2)	3.3 (0.4)	3.6 (0.4)	4.1 (0.3)	4.8 (0.3)	5.7 (0.4)	6.5 (0.6)	7.1 (0.7)
	19+	471	4.9 (0.2)	3.4 (0.4)	3.7 (0.4)	4.3 (0.3)	4.9 (0.2)	5.6 (0.3)	6.3 (0.4)	6.8 (0.5)
Female										
	9-13	103	4.1 (0.2)	3.4 (0.4)	3.6 (0.4)	3.9 (0.3)	4.2 (0.3)	4.6 (0.3)	5.0 (0.4)	5.3 (0.5)
	14-18	142	4.3 (0.3)	3.3 (0.3)	3.4 (0.3)	3.8 (0.3)	4.1 (0.4)	4.6 (0.4)	5.0 (0.5)	5.3 (0.5)
	19-30	111	4.5 (0.2)	3.5 (0.2)	3.7 (0.3)	4.1 (0.3)	4.5 (0.3)	5.0 (0.3)	5.5 (0.4)	5.8 (0.4)
	31-50	146	4.2 (0.3)	2.8 (0.3)	3.1 (0.3)	3.4 (0.3)	3.9 (0.3)	4.4 (0.4)	4.9 (0.4)	5.2 (0.4)
	51-70	184	4.8 (0.2)	3.2 (0.5)	3.5 (0.4)	4.0 (0.4)	4.7 (0.3)	5.4 (0.4)	6.2 (0.6)	6.7 (0.8)
	>70	143	4.4 (0.2)	3.4 (0.2)	3.6 (0.3)	4.0 (0.3)	4.4 (0.3)	4.9 (0.4)	5.4 (0.4)	5.7 (0.4)
	19+	584	4.4 (0.1)	3.4 (0.4)	3.6 (0.3)	3.9 (0.2)	4.3 (0.2)	4.8 (0.2)	5.2 (0.3)	5.5 (0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.9 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	169	3.3 (0.2)	2.0 (0.3) ^E	2.2 (0.3)	2.7 (0.2)	3.2 (0.2)	3.8 (0.2)	4.4 (0.4)	4.7 (0.5)
	4-8	281	3.9 (0.1)	2.6 (0.3)	2.8 (0.3)	3.2 (0.2)	3.8 (0.1)	4.4 (0.2)	5.1 (0.4)	5.5 (0.5)
Male										
	9-13	183	4.2 (0.2)	2.9 (0.4)	3.2 (0.3)	3.6 (0.3)	4.0 (0.2)	4.6 (0.3)	5.2 (0.5)	5.7 (0.6)
	14-18	187	4.3 (0.2)	3.0 (0.4)	3.2 (0.4)	3.6 (0.3)	4.2 (0.3)	4.8 (0.3)	5.6 (0.6)	6.1 (0.9)
	19-30	223	4.1 (0.2)	3.4 (0.2)	3.5 (0.2)	3.8 (0.2)	4.1 (0.2)	4.5 (0.2)	4.8 (0.3)	5.0 (0.3)
	31-50	229	4.7 (0.3)	3.8 (0.3)	4.0 (0.3)	4.4 (0.3)	4.8 (0.4)	5.3 (0.4)	5.8 (0.5)	6.1 (0.5)
	51-70	197	4.4 (0.2)	2.8 (0.5) ^E	3.1 (0.4)	3.7 (0.4)	4.4 (0.3)	5.2 (0.4)	6.0 (0.6)	6.6 (0.7)
	>70	72	5.1 (0.5)	3.8 (0.6)	4.1 (0.5)	4.5 (0.4)	5.1 (0.5)	5.7 (0.6)	6.4 (1.0)	6.8 (1.3) ^E
	19+	721	4.5 (0.2)	3.1 (0.4)	3.4 (0.4)	3.9 (0.3)	4.6 (0.2)	5.3 (0.3)	6.0 (0.4)	6.5 (0.6)
Female										
	9-13	165	4.1 (0.2)	3.0 (0.2)	3.2 (0.2)	3.6 (0.2)	4.0 (0.3)	4.5 (0.3)	5.1 (0.4)	5.5 (0.4)
	14-18	206	4.5 (0.3)	3.4 (0.2)	3.6 (0.2)	4.0 (0.3)	4.4 (0.3)	4.9 (0.3)	5.4 (0.3)	5.7 (0.4)
	19-30	191	4.1 (0.2)	3.0 (0.5)	3.2 (0.4)	3.6 (0.4)	4.1 (0.3)	4.6 (0.3)	5.1 (0.5)	5.4 (0.7)
	31-50	258	4.7 (0.3)	3.4 (0.2)	3.6 (0.2)	4.0 (0.2)	4.5 (0.3)	5.1 (0.3)	5.7 (0.4)	6.1 (0.4)
	51-70	249	4.5 (0.2)	3.1 (0.4)	3.4 (0.4)	3.8 (0.3)	4.4 (0.2)	5.0 (0.3)	5.5 (0.5)	5.9 (0.6)
	>70	128	4.1 (0.4)	2.7 (0.6) ^E	3.0 (0.6) ^E	3.4 (0.5)	3.9 (0.5)	4.5 (0.5)	5.2 (0.6)	5.6 (0.7)
	19+	826	4.5 (0.1)	3.1 (0.3)	3.3 (0.3)	3.8 (0.2)	4.3 (0.2)	4.9 (0.2)	5.6 (0.4)	6.0 (0.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.10 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	192	3.7 (0.2)	2.7 (0.3)	2.9 (0.3)	3.2 (0.2)	3.5 (0.2)	3.9 (0.3)	4.2 (0.4)	4.5 (0.5)
	4-8	321	4.2 (0.2)	2.8 (0.1)	3.0 (0.2)	3.5 (0.2)	4.0 (0.2)	4.7 (0.3)	5.4 (0.4)	5.9 (0.5)
Male										
	9-13	226	4.4 (0.2)	3.1 (0.4)	3.3 (0.3)	3.7 (0.3)	4.2 (0.3)	4.8 (0.3)	5.4 (0.5)	5.8 (0.5)
	14-18	262	4.8 (0.3)	3.5 (0.3)	3.8 (0.3)	4.2 (0.3)	4.7 (0.3)	5.2 (0.4)	5.7 (0.4)	6.0 (0.4)
	19-30	197	4.6 (0.2)	3.3 (0.4)	3.6 (0.3)	4.0 (0.3)	4.5 (0.3)	5.2 (0.4)	5.8 (0.6)	6.3 (0.7)
	31-50	282	4.5 (0.2)	3.1 (0.4)	3.4 (0.4)	3.9 (0.3)	4.5 (0.3)	5.2 (0.3)	5.9 (0.5)	6.4 (0.6)
	51-70	234	4.2 (0.1)	3.2 (0.4)	3.4 (0.4)	3.8 (0.3)	4.2 (0.2)	4.7 (0.2)	5.1 (0.4)	5.4 (0.6)
	>70	119	4.5 (0.2)	3.0 (0.4)	3.3 (0.4)	3.8 (0.3)	4.4 (0.3)	5.1 (0.4)	5.8 (0.5)	6.3 (0.6)
	19+	832	4.4 (0.1)	3.1 (0.2)	3.4 (0.2)	3.8 (0.2)	4.4 (0.1)	5.1 (0.2)	5.7 (0.2)	6.1 (0.3)
Female										
	9-13	226	3.9 (0.2)	3.4 (0.4)	3.5 (0.3)	3.7 (0.3)	3.9 (0.2)	4.2 (0.3)	4.4 (0.4)	4.5 (0.5)
	14-18	242	4.5 (0.2)	3.2 (0.5)	3.5 (0.5)	3.9 (0.4)	4.5 (0.3)	5.1 (0.4)	5.7 (0.6)	6.1 (0.6)
	19-30	208	4.6 (0.3)	2.7 (0.4)	3.0 (0.4)	3.5 (0.3)	4.3 (0.3)	5.1 (0.4)	6.1 (0.6)	6.7 (0.8)
	31-50	263	4.6 (0.2)	3.6 (0.4)	3.8 (0.4)	4.2 (0.3)	4.7 (0.2)	5.2 (0.3)	5.6 (0.4)	5.9 (0.6)
	51-70	322	4.4 (0.2)	2.9 (0.4)	3.2 (0.4)	3.7 (0.3)	4.2 (0.2)	4.9 (0.3)	5.5 (0.4)	6.0 (0.5)
	>70	198	4.1 (0.2)	2.9 (0.3)	3.1 (0.3)	3.5 (0.2)	3.9 (0.2)	4.5 (0.3)	5.0 (0.4)	5.4 (0.5)
	19+	991	4.5 (0.1)	3.0 (0.2)	3.3 (0.2)	3.8 (0.1)	4.4 (0.1)	5.0 (0.2)	5.7 (0.3)	6.1 (0.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.11 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	348	3.3	(0.1)	2.3	(0.2)	2.5	(0.2)	2.8	(0.1)	3.2	(0.1)	3.7	(0.2)	4.1	(0.2)	4.4	(0.3)
	4-8	554	3.9	(0.1)	2.8	(0.2)	3.0	(0.2)	3.4	(0.1)	3.9	(0.1)	4.4	(0.2)	4.9	(0.2)	5.2	(0.3)
Male																		
	9-13	409	4.6	(0.2)	3.8	(0.3)	4.0	(0.3)	4.3	(0.3)	4.7	(0.3)	5.2	(0.4)	5.6	(0.5)	5.9	(0.6)
	14-18	414	4.4	(0.2)	3.3	(0.2)	3.5	(0.2)	3.9	(0.2)	4.4	(0.2)	4.9	(0.2)	5.4	(0.3)	5.7	(0.3)
	19-30	311	4.2	(0.2)	2.9	(0.3)	3.2	(0.3)	3.6	(0.3)	4.2	(0.2)	4.8	(0.2)	5.3	(0.3)	5.7	(0.4)
	31-50	489	4.5	(0.1)	3.0	(0.2)	3.2	(0.2)	3.7	(0.2)	4.3	(0.2)	4.9	(0.2)	5.6	(0.3)	6.0	(0.3)
	51-70	575	4.5	(0.2)	4.0	(0.5)	4.1	(0.4)	4.3	(0.3)	4.6	(0.2)	4.9	(0.2)	5.1	(0.4)	5.3	(0.5)
	>70	239	4.3	(0.2)	3.0	(0.2)	3.2	(0.2)	3.7	(0.2)	4.3	(0.2)	4.8	(0.3)	5.4	(0.3)	5.7	(0.3)
	19+	1614	4.4	(0.1)	3.1	(0.1)	3.3	(0.1)	3.8	(0.1)	4.3	(0.1)	4.9	(0.1)	5.6	(0.2)	5.9	(0.2)
Female																		
	9-13	355	4.2	(0.2)	3.2	(0.2)	3.4	(0.2)	3.8	(0.2)	4.2	(0.2)	4.7	(0.2)	5.2	(0.2)	5.5	(0.2)
	14-18	410	4.2	(0.2)	3.1	(0.3)	3.3	(0.3)	3.6	(0.2)	4.1	(0.2)	4.6	(0.3)	5.1	(0.4)	5.5	(0.5)
	19-30	384	4.7	(0.3)	3.5	(0.5)	3.8	(0.4)	4.2	(0.4)	4.6	(0.3)	5.1	(0.4)	5.6	(0.5)	5.9	(0.6)
	31-50	585	4.5	(0.1)	3.6	(0.4)	3.8	(0.3)	4.1	(0.2)	4.5	(0.2)	4.8	(0.2)	5.2	(0.3)	5.4	(0.4)
	51-70	711	4.8	(0.2)	3.4	(0.3)	3.7	(0.3)	4.2	(0.2)	4.8	(0.2)	5.5	(0.3)	6.2	(0.3)	6.6	(0.4)
	>70	342	4.2	(0.1)	3.0	(0.2)	3.2	(0.2)	3.7	(0.2)	4.2	(0.2)	4.8	(0.2)	5.4	(0.3)	5.8	(0.3)
	19+	2022	4.6	(0.1)	3.3	(0.2)	3.6	(0.2)	4.0	(0.1)	4.6	(0.1)	5.2	(0.1)	5.8	(0.2)	6.2	(0.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.12 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	622	3.4	(0.1)	2.1	(0.1)	2.3	(0.1)	2.8	(0.1)	3.3	(0.1)	3.8	(0.2)	4.4	(0.2)	4.7	(0.2)
	4-8	919	4.0	(0.1)	2.6	(0.2)	2.8	(0.2)	3.3	(0.1)	3.9	(0.1)	4.6	(0.2)	5.3	(0.2)	5.8	(0.3)
Male																		
	9-13	579	4.4	(0.2)	2.8	(0.2)	3.1	(0.2)	3.6	(0.2)	4.2	(0.2)	5.0	(0.3)	5.8	(0.4)	6.3	(0.5)
	14-18	634	4.3	(0.1)	3.8	(0.5)	3.9	(0.4)	4.2	(0.3)	4.4	(0.2)	4.6	(0.2)	4.9	(0.4)	5.0	(0.6)
	19-30	578	4.3	(0.1)	3.6	(0.5)	3.8	(0.4)	4.0	(0.3)	4.4	(0.2)	4.7	(0.2)	5.1	(0.4)	5.3	(0.6)
	31-50	693	4.7	(0.2)	3.7	(0.5)	3.9	(0.5)	4.3	(0.4)	4.8	(0.3)	5.3	(0.3)	5.7	(0.5)	6.0	(0.6)
	51-70	596	4.6	(0.1)	3.4	(0.4)	3.7	(0.3)	4.1	(0.2)	4.6	(0.2)	5.2	(0.3)	5.7	(0.4)	6.0	(0.5)
	>70	296	4.9	(0.3)	3.0	(0.2)	3.4	(0.2)	4.0	(0.2)	4.8	(0.2)	5.7	(0.4)	6.8	(0.6)	7.6	(0.8)
	19+	2163	4.6	(0.1)	3.3	(0.3)	3.5	(0.2)	4.0	(0.2)	4.6	(0.1)	5.3	(0.2)	6.0	(0.3)	6.5	(0.4)
Female																		
	9-13	533	4.1	(0.1)	3.0	(0.3)	3.2	(0.3)	3.6	(0.2)	4.1	(0.2)	4.6	(0.2)	5.1	(0.3)	5.5	(0.4)
	14-18	638	4.4	(0.2)	3.3	(0.1)	3.5	(0.2)	3.9	(0.2)	4.3	(0.2)	4.8	(0.2)	5.2	(0.2)	5.5	(0.3)
	19-30	499	4.1	(0.1)	3.1	(0.1)	3.3	(0.2)	3.7	(0.2)	4.1	(0.2)	4.6	(0.2)	5.0	(0.2)	5.4	(0.2)
	31-50	716	4.6	(0.2)	3.2	(0.1)	3.5	(0.1)	3.9	(0.2)	4.4	(0.2)	4.9	(0.2)	5.5	(0.2)	5.8	(0.3)
	51-70	745	4.5	(0.1)	2.9	(0.2)	3.2	(0.2)	3.7	(0.2)	4.4	(0.2)	5.1	(0.2)	5.9	(0.3)	6.4	(0.4)
	>70	510	4.2	(0.2)	3.1	(0.4)	3.3	(0.3)	3.7	(0.3)	4.1	(0.2)	4.6	(0.3)	5.1	(0.4)	5.4	(0.5)
	19+	2470	4.4	(0.1)	3.1	(0.2)	3.4	(0.2)	3.8	(0.1)	4.3	(0.1)	4.9	(0.1)	5.5	(0.2)	5.9	(0.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 17.13 Percentage of total energy intake from linoleic acid, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both											
	1-3	2117	3.3	(0.1)	2.1 (0.1)	2.3 (0.1)	2.7 (0.1)	3.2 (0.1)	3.8 (0.1)	4.3 (0.1)	4.7 (0.1)
	4-8	3235	3.9	(0.1)	2.8 (0.1)	3.0 (0.1)	3.4 (0.1)	3.9 (0.1)	4.4 (0.1)	5.0 (0.1)	5.4 (0.2)
Male											
	9-13	2080	4.2	(0.1)	3.1 (0.1)	3.3 (0.1)	3.7 (0.1)	4.2 (0.1)	4.8 (0.1)	5.4 (0.2)	5.8 (0.2)
	14-18	2288	4.3	(0.1)	3.4 (0.2)	3.6 (0.2)	3.9 (0.1)	4.3 (0.1)	4.6 (0.1)	5.0 (0.2)	5.3 (0.3)
	19-30	1804	4.5	(0.1)	3.3 (0.2)	3.6 (0.2)	4.0 (0.1)	4.5 (0.1)	5.1 (0.2)	5.6 (0.2)	6.0 (0.3)
	31-50	2596	4.4	(0.1)	2.9 (0.2)	3.2 (0.1)	3.7 (0.1)	4.3 (0.1)	5.0 (0.1)	5.7 (0.2)	6.2 (0.2)
	51-70	2550	4.5	(0.1)	3.1 (0.2)	3.3 (0.2)	3.8 (0.1)	4.4 (0.1)	5.1 (0.1)	5.8 (0.2)	6.2 (0.3)
	>70	1520	4.5	(0.1)	2.9 (0.1)	3.2 (0.1)	3.8 (0.1)	4.5 (0.1)	5.2 (0.1)	5.9 (0.2)	6.4 (0.2)
	19+	8470	4.4	(0.0)	3.0 (0.1)	3.2 (0.1)	3.7 (0.1)	4.4 (0.1)	5.1 (0.1)	5.8 (0.1)	6.3 (0.1)
Female											
	9-13	1980	4.2	(0.1)	3.1 (0.1)	3.3 (0.1)	3.7 (0.1)	4.2 (0.1)	4.7 (0.1)	5.2 (0.2)	5.6 (0.2)
	14-18	2256	4.4	(0.1)	3.4 (0.2)	3.6 (0.2)	4.0 (0.1)	4.4 (0.1)	4.9 (0.1)	5.3 (0.2)	5.6 (0.3)
	19-30	1854	4.2	(0.1)	3.1 (0.2)	3.3 (0.2)	3.7 (0.1)	4.1 (0.1)	4.6 (0.2)	5.1 (0.2)	5.5 (0.3)
	31-50	2686	4.5	(0.1)	3.3 (0.2)	3.5 (0.2)	3.9 (0.1)	4.5 (0.1)	5.1 (0.1)	5.7 (0.2)	6.1 (0.3)
	51-70	3200	4.5	(0.1)	3.0 (0.1)	3.3 (0.1)	3.8 (0.1)	4.4 (0.1)	5.2 (0.1)	5.9 (0.2)	6.4 (0.2)
	>70	2610	4.4	(0.1)	2.9 (0.1)	3.2 (0.1)	3.7 (0.1)	4.3 (0.1)	5.0 (0.1)	5.7 (0.2)	6.2 (0.2)
	19+	10350	4.4	(0.0)	3.1 (0.1)	3.3 (0.1)	3.8 (0.1)	4.4 (0.1)	5.0 (0.1)	5.7 (0.1)	6.1 (0.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

18. Magnesium (mg/d): Usual intakes from food

Table 18.1 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	197 (12)	155 (19)	163 (17)	176 (14)	191 (14)	207 (19)	223 (25)	233 (31)	65	<3	
	4-8	127	239 (10)	169 (19)	182 (18)	207 (15)	237 (13)	272 (15)	308 (21)	332 (26)	110	<3	
Male	9-13	111	320 (21)	230 (23)	247 (22)	277 (22)	314 (24)	356 (31)	398 (45)	427 (56)	200	F	
	14-18	107	320 (21)	188 (21)	209 (22)	253 (23)	322 (26)	417 (38)	520 (59)	592 (77)	340	55.6	(10.0) ^E
	19-30	77	306 (16)	203 (15)	226 (16)	265 (17)	309 (20)	354 (23)	399 (25)	429 (28)	330	62.6	(11.9) ^E
	31-50	145	311 (24)	211 (32)	230 (28)	263 (24)	303 (22)	349 (25)	397 (35)	429 (44)	350	75.6	(12.3)
	51-70	182	312 (17)	172 (28)	193 (26)	235 (23)	290 (20)	356 (23)	431 (37)	486 (51)	350	73.3	(7.9)
	>70	63	349 (38)	227 (37)	249 (37)	291 (41)	348 (50)	420 (68)	501 (95) ^E	558 (117) ^E	350	F	
	19+	467	314 (12)	181 (9)	205 (9)	250 (10)	304 (12)	365 (15)	433 (20)	482 (26)			
	Female												
Female	9-13	96	245 (10)	167 (18)	184 (17)	215 (15)	253 (15)	295 (17)	336 (23)	362 (27)	200	F	
	14-18	105	246 (25)	128 (28) ^E	147 (27) ^E	179 (28)	232 (30)	309 (39)	397 (60)	464 (81) ^E	300	72.5	(12.3) ^E
	19-30	91	228 (14)	164 (19)	175 (18)	194 (17)	218 (17)	247 (21)	277 (30)	297 (38)	255	80.2	(12.7)
	31-50	167	248 (17)	164 (25)	180 (23)	212 (22)	255 (23)	306 (28)	360 (38)	395 (46)	265	55.6	(13.0) ^E
	51-70	198	256 (12)	148 (19)	168 (17)	205 (15)	252 (14)	304 (19)	357 (27)	391 (33)	265	57.1	(9.2)
	>70	74	263 (24)	159 (31) ^E	179 (29)	212 (27)	250 (26)	291 (26)	334 (31)	363 (38)	265	60.4	(17.1) ^E
	19+	530	248 (9)	147 (11)	166 (10)	201 (10)	247 (11)	303 (14)	360 (19)	399 (23)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.2 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	58	231 (17)	170 (21)	183 (19)	207 (17)	234 (18)	264 (22)	293 (28)	312 (33)	65	<3	
	4-8	110	250 (14)	180 (22)	196 (19)	223 (17)	254 (17)	286 (21)	315 (26)	331 (29)	110	<3	
Male													
	9-13	95	298 (17)	208 (32)	227 (29)	260 (25)	295 (24)	333 (25)	370 (29)	394 (33)	200	F	
	14-18	87	340 (18)	188 (34) ^E	220 (30)	276 (23)	344 (21)	425 (30)	516 (53)	583 (74)	340	48.5 (8.6) ^E	
	19-30	70	416 (31)	232 (50) ^E	268 (48) ^E	339 (45)	426 (42)	517 (43)	604 (53)	660 (65)	330	F	
	31-50	109	348 (23)	206 (37) ^E	230 (35)	277 (30)	340 (29)	416 (37)	498 (54)	554 (66)	350	53.8 (14.3) ^E	
	51-70	128	354 (14)	231 (22)	256 (19)	299 (16)	347 (14)	403 (19)	470 (37)	525 (58)	350	51.4 (8.0)	
	>70	65	282 (15)	179 (22)	198 (21)	231 (19)	272 (19)	316 (23)	359 (31)	387 (36)	350	87.6 (8.0)	
	19+	372	357 (12)	210 (15)	238 (14)	288 (14)	353 (14)	431 (18)	518 (27)	580 (35)			
Female													
	9-13	75	261 (20)	181 (33) ^E	198 (31)	228 (28)	264 (29)	302 (34)	339 (42)	362 (48)	200	F	
	14-18	81	252 (15)	124 (27) ^E	149 (24)	193 (19)	242 (18)	291 (23)	346 (33)	388 (44)	300	78.2 (8.7)	
	19-30	101	287 (21)	170 (25)	193 (24)	236 (24)	290 (27)	352 (36)	415 (47)	456 (56)	255	F	
	31-50	116	275 (17)	159 (20)	181 (20)	224 (19)	272 (21)	325 (27)	384 (33)	419 (37)	265	46.3 (11.4) ^E	
	51-70	146	283 (9)	164 (15)	184 (14)	223 (14)	273 (14)	330 (17)	388 (21)	425 (24)	265	46.0 (7.4)	
	>70	94	234 (14)	143 (18)	157 (17)	184 (15)	224 (17)	272 (26)	321 (40)	356 (53)	265	72.0 (11.8)	
	19+	457	274 (8)	159 (8)	180 (8)	219 (9)	269 (11)	326 (14)	382 (17)	419 (19)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.3 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	112	211 (10)	150 (20)	162 (19)	182 (15)	208 (13)	236 (15)	264 (21)	282 (25)	65	0.0 (0.0)	
	4-8	177	257 (11)	180 (21)	196 (19)	224 (16)	258 (14)	296 (17)	333 (25)	356 (32)	110	<3	
Male	9-13	111	309 (14)	195 (23)	217 (20)	257 (16)	306 (17)	360 (24)	413 (33)	446 (40)	200	F	
	14-18	113	315 (22)	224 (38) ^E	245 (36)	280 (32)	317 (31)	354 (37)	394 (49)	421 (57)	340	66.7 (20.0) ^E	
	19-30	91	380 (31)	192 (36) ^E	222 (34)	279 (31)	354 (31)	446 (42)	542 (61)	608 (76)	330	41.9 (11.7) ^E	
	31-50	101	371 (19)	250 (36)	273 (31)	315 (25)	365 (23)	418 (29)	469 (40)	500 (49)	350	42.0 (13.9) ^E	
	51-70	134	387 (16)	263 (34)	288 (30)	333 (25)	393 (21)	453 (25)	504 (34)	542 (47)	350	F	
	>70	56	299 (23)	252 (35)	263 (31)	282 (27)	305 (26)	330 (32)	353 (41)	368 (50)	350	88.5 (20.8) ^E	
	19+	382	371 (11)	221 (16)	249 (15)	300 (14)	363 (14)	433 (16)	500 (20)	545 (25)			
	Female	9-13	105	250 (18)	170 (26)	184 (24)	212 (22)	245 (21)	283 (24)	326 (32)	354 (40)	200	F
14-18	120	237 (18)	114 (31) ^E	149 (26) ^E	192 (21)	231 (19)	282 (22)	336 (31)	377 (37)	300	81.7 (7.7)		
19-30	91	280 (16)	199 (30)	218 (26)	246 (19)	279 (18)	317 (25)	352 (37)	377 (46)	255	F		
31-50	159	309 (13)	177 (20)	202 (20)	251 (19)	308 (17)	370 (21)	434 (29)	480 (37)	265	30.5 (7.8) ^E		
51-70	174	293 (12)	187 (25)	208 (22)	244 (17)	286 (14)	332 (16)	380 (25)	413 (35)	265	37.0 (10.2) ^E		
>70	80	269 (13)	165 (18)	187 (16)	223 (13)	263 (14)	313 (22)	389 (36)	449 (48)	265	51.4 (8.9) ^E		
19+	504	294 (7)	173 (11)	196 (10)	239 (9)	292 (9)	349 (11)	408 (15)	449 (19)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.4 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	99	244 (11)	190 (20)	203 (17)	226 (14)	253 (13)	282 (18)	310 (24)	328 (29)	65	0.0	(0.0)
	4-8	140	261 (14)	180 (21)	196 (19)	226 (16)	261 (16)	299 (22)	339 (33)	366 (41)	110	<3	
Male	9-13	92	281 (19)	207 (26)	220 (24)	246 (23)	277 (24)	310 (29)	341 (35)	360 (40)	200	F	
	14-18	107	366 (24)	232 (18)	257 (19)	304 (22)	366 (28)	443 (36)	530 (45)	593 (53)	340	39.4	(10.8) ^E
	19-30	73	422 (38)	223 (32)	260 (33)	332 (35)	426 (44)	548 (65)	695 (97)	806 (122)	330	F	
	31-50	134	358 (22)	258 (33)	273 (31)	300 (28)	335 (26)	375 (30)	416 (42)	444 (55)	350	60.6	(18.1) ^E
	51-70	131	359 (27)	213 (37) ^E	238 (35)	286 (31)	348 (31)	422 (40)	502 (57)	556 (73)	350	50.7	(13.8) ^E
	>70	55	315 (24)	153 (34) ^E	178 (33) ^E	227 (30)	295 (26)	380 (33)	475 (51)	541 (67)	350	67.5	(9.4)
	19+	393	367 (13)	196 (17)	226 (17)	282 (17)	354 (17)	446 (20)	551 (27)	624 (33)			
	Female	9-13	79	262 (18)	166 (24)	185 (26)	224 (28)	274 (27)	319 (24)	357 (23)	381 (24)	200	F
14-18	104	261 (14)	214 (34)	227 (28)	249 (21)	271 (17)	292 (19)	311 (27)	322 (34)	300	82.4	(13.9) ^E	
19-30	101	262 (16)	182 (27)	200 (26)	231 (24)	271 (23)	317 (24)	363 (29)	393 (34)	255	F		
31-50	143	290 (16)	169 (30) ^E	193 (25)	232 (19)	276 (16)	331 (26)	396 (47)	441 (64)	265	43.6	(10.9) ^E	
51-70	193	260 (10)	162 (21)	179 (18)	210 (14)	247 (13)	290 (17)	333 (25)	360 (31)	265	61.3	(10.0)	
>70	94	260 (16)	141 (18)	160 (19)	197 (19)	246 (20)	304 (24)	366 (28)	408 (31)	265	59.3	(10.0) ^E	
19+	531	272 (7)	159 (10)	180 (9)	219 (8)	268 (8)	321 (10)	376 (14)	415 (19)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.5 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	
Both													
	1-3	311	231	(9)	148 (17)	166 (16)	196 (14)	232 (13)	270 (14)	308 (17)	333 (20)	65	<3
	4-8	485	265	(8)	179 (15)	196 (13)	226 (10)	263 (9)	306 (14)	348 (22)	376 (27)	110	<3
Male													
	9-13	277	330	(14)	208 (20)	232 (19)	276 (18)	334 (19)	400 (25)	464 (33)	506 (39)	200	F
	14-18	339	376	(15)	243 (19)	270 (18)	320 (18)	387 (20)	467 (25)	555 (36)	617 (47)	340	32.5 (6.8) ^E
	19-30	237	393	(18)	284 (32)	309 (29)	352 (24)	400 (22)	453 (30)	508 (45)	545 (57)	330	F
	31-50	423	379	(15)	244 (24)	269 (22)	317 (19)	377 (18)	446 (21)	520 (31)	573 (40)	350	38.6 (8.2) ^E
	51-70	387	338	(10)	214 (16)	237 (14)	280 (13)	333 (12)	394 (16)	460 (24)	504 (32)	350	57.8 (5.8)
	>70	132	298	(20)	150 (30) ^E	180 (26)	231 (23)	291 (26)	370 (30)	453 (36)	505 (42)	350	69.9 (9.3)
	19+	1179	364	(8)	220 (12)	247 (12)	298 (10)	362 (10)	434 (12)	510 (18)	565 (23)		
	Female												
9-13		281	268	(12)	158 (17)	176 (16)	211 (14)	255 (13)	306 (17)	360 (24)	395 (30)	200	F
14-18		321	275	(8)	195 (16)	215 (15)	247 (12)	283 (11)	324 (12)	367 (16)	395 (21)	300	61.3 (7.3)
19-30		249	316	(16)	233 (28)	251 (25)	281 (20)	318 (19)	357 (24)	396 (33)	420 (40)	255	F
31-50		364	321	(12)	199 (13)	221 (12)	261 (13)	315 (14)	380 (20)	454 (30)	510 (41)	265	26.6 (5.6) ^E
51-70		467	312	(11)	197 (19)	220 (16)	262 (13)	312 (12)	368 (16)	427 (25)	468 (33)	265	26.5 (6.2) ^E
>70		215	248	(11)	161 (17)	178 (17)	210 (16)	250 (16)	295 (19)	341 (25)	373 (31)	265	59.1 (10.4) ^E
19+		1295	308	(7)	196 (8)	218 (8)	258 (7)	308 (8)	368 (11)	431 (15)	475 (19)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.6 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	644	221	(5)	141 (13)	159 (11)	188 (8)	223 (6)	261 (7)	298 (12)	322 (16)	65	<3
	4-8	956	256	(4)	175 (10)	190 (9)	218 (7)	253 (5)	294 (8)	334 (13)	360 (17)	110	<3
Male													
	9-13	589	309	(8)	196 (16)	218 (15)	256 (12)	303 (10)	357 (12)	416 (20)	457 (26)	200	F
	14-18	639	349	(8)	220 (22)	245 (20)	291 (15)	348 (11)	413 (13)	481 (22)	528 (30)	340	46.6 (5.9)
	19-30	481	369	(13)	222 (32)	250 (27)	300 (19)	362 (15)	432 (23)	506 (39)	556 (53)	330	36.7 (7.9) ^E
	31-50	709	360	(10)	225 (26)	249 (23)	292 (17)	346 (12)	412 (15)	485 (29)	533 (40)	350	51.7 (6.3)
	51-70	758	356	(8)	207 (11)	232 (10)	277 (9)	338 (9)	413 (12)	498 (19)	557 (25)	350	54.8 (3.6)
	>70	734	305	(7)	174 (11)	196 (10)	237 (9)	292 (8)	357 (9)	425 (14)	470 (19)	350	72.7 (3.0)
	19+	2682	356	(6)	208 (8)	233 (8)	281 (7)	341 (6)	415 (9)	498 (13)	554 (17)		
Female													
	9-13	585	264	(7)	144 (11)	166 (10)	207 (8)	257 (8)	312 (11)	367 (15)	401 (19)	200	22.0 (3.5)
	14-18	645	274	(7)	158 (12)	179 (11)	218 (9)	269 (9)	330 (11)	396 (18)	441 (24)	300	64.0 (4.0)
	19-30	514	267	(9)	142 (16)	163 (15)	204 (12)	255 (10)	311 (12)	371 (19)	414 (26)	255	50.1 (5.4)
	31-50	758	304	(8)	164 (10)	188 (9)	233 (8)	292 (8)	364 (12)	442 (19)	503 (27)	265	38.8 (3.6)
	51-70	955	301	(8)	160 (14)	185 (13)	229 (10)	285 (9)	355 (11)	433 (18)	489 (25)	265	41.1 (4.6)
	>70	1345	275	(6)	149 (6)	169 (6)	208 (6)	260 (6)	322 (8)	389 (11)	436 (14)	265	52.4 (2.9)
	19+	3572	292	(4)	154 (5)	178 (5)	222 (4)	279 (5)	347 (6)	424 (9)	480 (13)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.7 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	(SE)
Both													
	1-3	324	205	(6)	128 (14)	143 (12)	170 (9)	206 (7)	246 (9)	283 (14)	309 (20)	65	<3
	4-8	425	233	(8)	155 (14)	167 (12)	192 (10)	224 (9)	263 (12)	305 (20)	333 (26)	110	<3
Male													
	9-13	274	295	(9)	198 (23)	217 (19)	252 (14)	292 (11)	336 (13)	378 (20)	405 (25)	200	F
	14-18	297	364	(16)	183 (18)	214 (17)	274 (16)	352 (18)	444 (24)	543 (34)	609 (42)	340	46.1 (5.9)
	19-30	249	376	(26)	248 (43) ^E	270 (38)	311 (29)	363 (24)	422 (33)	480 (52)	518 (66)	330	F
	31-50	309	334	(15)	204 (24)	229 (22)	273 (19)	326 (16)	385 (26)	447 (33)	488 (39)	350	61.2 (8.1)
	51-70	277	327	(12)	196 (22)	219 (20)	262 (16)	316 (14)	379 (18)	444 (28)	486 (35)	350	64.6 (6.7)
	>70	136	309	(18)	205 (29)	224 (26)	259 (20)	301 (19)	348 (26)	394 (39)	424 (49)	350	75.9 (11.2)
	19+	971	339	(9)	205 (16)	229 (15)	274 (12)	329 (10)	389 (13)	450 (20)	494 (26)		
Female													
	9-13	265	246	(10)	143 (18)	163 (16)	197 (13)	238 (12)	285 (17)	334 (25)	368 (31)	200	26.8 (8.3) ^E
	14-18	290	255	(12)	163 (20)	180 (18)	210 (15)	247 (14)	291 (17)	339 (27)	373 (35)	300	78.7 (8.0)
	19-30	197	289	(18)	181 (29)	203 (26)	242 (22)	291 (21)	345 (27)	397 (37)	430 (46)	255	F
	31-50	312	295	(11)	182 (17)	204 (16)	243 (14)	287 (13)	339 (18)	400 (27)	444 (33)	265	36.7 (7.7) ^E
	51-70	312	298	(16)	190 (28)	208 (25)	241 (21)	284 (17)	334 (18)	387 (28)	422 (37)	265	38.8 (12.3) ^E
	>70	239	260	(12)	143 (15)	162 (14)	199 (12)	250 (12)	314 (15)	385 (24)	435 (32)	265	56.9 (5.8)
	19+	1060	290	(7)	169 (9)	190 (8)	230 (7)	279 (8)	340 (10)	407 (14)	452 (18)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.8 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	215 (10)	134 (16)	152 (15)	183 (13)	219 (13)	257 (15)	292 (17)	314 (19)	65	<3	
	4-8	213	242 (9)	158 (13)	173 (12)	201 (10)	236 (9)	275 (12)	314 (17)	340 (20)	110	<3	
Male	9-13	122	304 (16)	237 (23)	251 (21)	277 (19)	310 (19)	346 (24)	382 (31)	405 (36)	200	F	
	14-18	150	379 (23)	218 (23)	247 (22)	302 (22)	373 (27)	455 (36)	539 (48)	594 (57)	340	38.3	(9.2) ^E
	19-30	106	365 (23)	177 (34) ^E	211 (31)	273 (26)	349 (26)	439 (37)	538 (56)	607 (74)	330	43.6	(9.1) ^E
	31-50	155	358 (20)	186 (38) ^E	219 (34)	280 (28)	353 (24)	432 (29)	513 (42)	564 (52)	350	49.0	(9.9) ^E
	51-70	122	339 (16)	204 (29)	229 (25)	274 (20)	333 (20)	402 (32)	474 (51)	523 (67)	350	57.0	(12.0) ^E
	>70	88	365 (21)	202 (16)	223 (17)	268 (18)	338 (21)	427 (29)	518 (40)	577 (49)	350	53.9	(7.9)
	19+	471	356 (11)	182 (14)	214 (13)	273 (12)	344 (12)	427 (16)	517 (22)	579 (28)			
	Female	9-13	103	275 (15)	191 (20)	207 (19)	235 (18)	272 (18)	316 (22)	365 (30)	398 (37)	200	F
14-18	142	250 (13)	146 (15)	166 (14)	201 (14)	246 (18)	303 (21)	363 (27)	401 (31)	300	73.9	(8.0)	
19-30	111	269 (16)	173 (27)	192 (24)	224 (22)	262 (23)	306 (29)	353 (40)	385 (50)	255	F		
31-50	146	299 (15)	159 (28) ^E	184 (26)	226 (22)	279 (20)	343 (23)	406 (29)	447 (35)	265	43.2	(10.7) ^E	
51-70	184	304 (17)	207 (29)	226 (25)	260 (20)	301 (18)	347 (26)	396 (42)	430 (55)	265	F		
>70	143	294 (12)	190 (16)	214 (15)	254 (13)	301 (14)	353 (18)	406 (25)	442 (30)	265	30.6	(7.2) ^E	
19+	584	293 (8)	171 (11)	193 (10)	233 (10)	283 (10)	342 (13)	404 (17)	445 (21)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.9 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	169	202 (9)	114 (14)	130 (13)	159 (12)	197 (11)	243 (14)	290 (20)	322 (26)	65	<3	
	4-8	281	248 (8)	185 (19)	198 (16)	220 (12)	247 (9)	276 (13)	305 (21)	323 (28)	110	<3	
Male													
	9-13	183	331 (22)	205 (35) ^E	226 (32)	266 (27)	320 (24)	390 (32)	475 (56)	539 (80)	200	F	
	14-18	187	367 (18)	240 (46) ^E	268 (39)	317 (27)	372 (20)	434 (29)	500 (48)	544 (63)	340	F	
	19-30	223	349 (16)	235 (34)	257 (30)	295 (23)	342 (18)	395 (22)	447 (34)	482 (44)	330	43.3 (12.0) ^E	
	31-50	229	359 (16)	243 (33)	266 (29)	308 (24)	362 (21)	426 (30)	497 (47)	546 (62)	350	44.4 (12.2) ^E	
	51-70	197	346 (16)	188 (20)	214 (20)	266 (19)	334 (20)	416 (23)	504 (32)	563 (41)	350	55.7 (7.4)	
	>70	72	332 (20)	217 (33)	241 (29)	281 (25)	331 (22)	390 (27)	456 (41)	504 (55)	350	59.3 (12.0) ^E	
	19+	721	352 (9)	220 (14)	246 (13)	291 (12)	349 (11)	417 (13)	493 (18)	544 (23)			
	Female												
9-13		165	264 (14)	182 (29)	199 (26)	231 (21)	271 (19)	317 (25)	363 (37)	394 (47)	200	F	
14-18		206	253 (9)	163 (20)	180 (17)	212 (14)	250 (11)	292 (13)	333 (21)	361 (27)	300	78.9 (6.9)	
19-30		191	269 (15)	157 (20)	174 (20)	212 (18)	262 (18)	315 (30)	373 (39)	418 (49)	255	46.5 (10.0) ^E	
31-50		258	275 (12)	195 (28)	211 (25)	239 (20)	274 (16)	315 (20)	359 (31)	389 (40)	265	F	
51-70		249	280 (11)	157 (15)	179 (14)	219 (12)	269 (12)	329 (16)	395 (24)	438 (29)	265	48.1 (6.2)	
>70		128	263 (12)	172 (17)	192 (16)	225 (15)	265 (15)	314 (21)	376 (33)	424 (45)	265	49.9 (9.5) ^E	
19+		826	274 (7)	166 (9)	186 (9)	222 (8)	268 (9)	323 (10)	386 (14)	429 (18)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.10 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	223 (8)	151 (14)	164 (12)	188 (10)	216 (10)	248 (12)	278 (17)	298 (20)	65	0.0	(0.0)
	4-8	321	269 (7)	176 (14)	194 (13)	227 (10)	268 (9)	312 (11)	356 (17)	384 (21)	110	<3	
Male													
	9-13	226	318 (12)	187 (14)	211 (13)	255 (13)	311 (15)	383 (19)	469 (27)	536 (36)	200	F	
	14-18	262	402 (29)	227 (37)	258 (35)	315 (31)	391 (30)	484 (37)	586 (53)	658 (68)	340	32.9	(10.2) ^E
	19-30	197	422 (20)	221 (33)	251 (32)	318 (28)	405 (26)	524 (36)	674 (63)	771 (84)	330	28.3	(8.2) ^E
	31-50	282	432 (19)	255 (19)	283 (18)	338 (17)	414 (19)	505 (28)	604 (43)	675 (59)	350	28.9	(5.7) ^E
	51-70	234	384 (17)	221 (22)	249 (21)	302 (20)	367 (21)	444 (26)	528 (36)	585 (44)	350	43.3	(8.2) ^E
	>70	119	367 (29)	167 (31) ^E	201 (30)	266 (31)	354 (36)	458 (45)	566 (60)	639 (71)	350	49.0	(10.8) ^E
	19+	832	410 (11)	225 (10)	256 (9)	316 (9)	394 (11)	491 (17)	598 (23)	674 (30)			
Female													
	9-13	226	277 (13)	168 (16)	189 (16)	228 (16)	277 (16)	334 (18)	392 (21)	429 (24)	200	F	
	14-18	242	275 (15)	138 (15)	160 (15)	203 (14)	258 (16)	327 (23)	412 (34)	475 (45)	300	66.7	(7.2)
	19-30	208	303 (15)	217 (28)	235 (26)	267 (21)	306 (19)	352 (24)	400 (35)	433 (45)	255	F	
	31-50	263	323 (13)	200 (21)	221 (19)	258 (15)	304 (15)	357 (22)	413 (35)	452 (43)	265	28.6	(8.2) ^E
	51-70	322	309 (14)	174 (22)	199 (20)	245 (18)	304 (18)	370 (21)	437 (28)	480 (35)	265	33.1	(8.0) ^E
	>70	198	286 (11)	178 (17)	198 (16)	235 (15)	282 (15)	335 (18)	389 (24)	424 (30)	265	40.9	(8.7) ^E
	19+	991	311 (7)	187 (10)	210 (9)	251 (8)	302 (8)	360 (11)	422 (16)	465 (21)			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.11 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	(SE)
Both													
	1-3	348	220	(6)	158 (11)	171 (10)	193 (8)	220 (8)	249 (9)	279 (12)	297 (14)	65	0.0 (0.0)
	4-8	554	254	(7)	169 (9)	186 (8)	217 (8)	256 (9)	299 (10)	345 (14)	375 (17)	110	<3
Male													
	9-13	409	302	(9)	203 (12)	221 (12)	255 (11)	297 (11)	343 (14)	390 (18)	420 (21)	200	F
	14-18	414	334	(12)	199 (13)	224 (14)	272 (15)	334 (16)	407 (20)	489 (25)	544 (29)	340	52.4 (6.4)
	19-30	311	379	(18)	213 (16)	242 (15)	297 (15)	370 (18)	459 (25)	559 (36)	630 (46)	330	36.2 (5.9)
	31-50	489	352	(11)	213 (15)	237 (14)	281 (13)	336 (14)	400 (17)	466 (22)	511 (27)	350	56.4 (6.4)
	51-70	575	359	(12)	206 (16)	235 (16)	287 (14)	352 (14)	427 (17)	507 (23)	565 (31)	350	49.2 (5.5)
	>70	239	313	(15)	189 (18)	211 (18)	255 (17)	312 (18)	375 (19)	442 (26)	490 (35)	350	66.3 (7.5)
	19+	1614	356	(7)	203 (8)	230 (8)	280 (7)	345 (8)	421 (10)	505 (13)	564 (16)		
Female													
	9-13	355	253	(9)	166 (13)	183 (13)	214 (13)	255 (13)	302 (14)	349 (18)	380 (21)	200	F
	14-18	410	247	(10)	135 (17)	162 (14)	200 (11)	241 (11)	297 (15)	357 (21)	395 (25)	300	76.0 (5.3)
	19-30	384	263	(9)	180 (12)	196 (11)	225 (11)	262 (11)	307 (14)	355 (19)	387 (23)	255	45.0 (7.7) ^E
	31-50	585	287	(8)	154 (10)	180 (10)	229 (10)	286 (10)	350 (13)	421 (18)	469 (23)	265	40.2 (4.6)
	51-70	711	273	(6)	156 (9)	177 (8)	216 (8)	265 (8)	320 (10)	374 (12)	408 (14)	265	50.0 (4.5)
	>70	342	263	(9)	153 (9)	174 (9)	211 (9)	254 (10)	304 (12)	365 (17)	409 (19)	265	56.6 (5.7)
	19+	2022	276	(4)	160 (5)	182 (5)	223 (5)	272 (5)	330 (6)	391 (9)	431 (11)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.12 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	622	205	(6)	122 (8)	138 (8)	166 (7)	203 (7)	245 (9)	288 (11)	316 (15)	65	<3
	4-8	919	244	(5)	160 (8)	176 (7)	204 (6)	238 (6)	278 (8)	320 (11)	348 (14)	110	<3
Male													
	9-13	579	319	(13)	218 (19)	236 (18)	270 (16)	314 (15)	368 (19)	427 (30)	469 (40)	200	F
	14-18	634	369	(12)	208 (16)	242 (15)	300 (13)	363 (13)	438 (19)	531 (28)	597 (34)	340	40.3 (5.1)
	19-30	578	357	(12)	198 (22)	227 (20)	280 (16)	344 (14)	418 (17)	500 (27)	556 (36)	330	44.2 (6.1)
	31-50	693	354	(11)	221 (19)	248 (17)	295 (15)	353 (15)	420 (17)	495 (22)	545 (28)	350	48.6 (6.8)
	51-70	596	340	(10)	191 (11)	217 (11)	266 (11)	329 (12)	407 (15)	490 (21)	547 (28)	350	57.6 (4.8)
	>70	296	335	(12)	190 (18)	214 (18)	262 (17)	324 (13)	391 (29)	468 (74)	527 (115) ^E	350	60.7 (5.6)
	19+	2163	350	(6)	201 (8)	230 (7)	279 (7)	342 (7)	417 (9)	497 (12)	554 (15)		
	Female												
9-13		533	262	(9)	169 (13)	187 (12)	220 (11)	262 (12)	311 (15)	362 (19)	398 (23)	200	15.2 (5.1) ^E
14-18		638	253	(6)	152 (9)	172 (8)	206 (8)	248 (8)	296 (9)	349 (13)	385 (17)	300	76.5 (3.7)
19-30		499	274	(11)	165 (12)	184 (12)	220 (12)	267 (12)	319 (15)	376 (21)	416 (26)	255	43.5 (7.0)
31-50		716	283	(8)	177 (16)	196 (15)	233 (13)	276 (11)	328 (13)	385 (17)	423 (22)	265	43.7 (7.1)
51-70		745	289	(8)	173 (10)	194 (9)	230 (8)	277 (8)	335 (12)	397 (17)	439 (21)	265	43.7 (4.4)
>70		510	270	(7)	158 (8)	181 (8)	221 (8)	269 (9)	327 (12)	392 (16)	439 (21)	265	47.8 (4.7)
19+		2470	281	(4)	164 (6)	185 (6)	224 (5)	273 (6)	331 (7)	396 (9)	442 (11)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 18.13 Magnesium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)	
Both												
	1-3	2117	220 (3)	137 (5)	154 (5)	185 (4)	221 (4)	260 (4)	299 (5)	324 (6)	65	<3
	4-8	3235	257 (3)	167 (4)	184 (4)	215 (3)	254 (4)	299 (5)	344 (7)	374 (8)	110	<3
Male												
	9-13	2080	316 (5)	201 (7)	223 (6)	262 (6)	311 (6)	371 (8)	436 (11)	480 (14)	200	4.7 (1.1) ^E
	14-18	2288	364 (6)	217 (8)	245 (8)	296 (7)	362 (8)	441 (10)	529 (13)	591 (17)	340	41.5 (2.9)
	19-30	1804	380 (8)	218 (9)	248 (9)	302 (9)	372 (9)	453 (12)	541 (17)	603 (21)	330	34.8 (3.2)
	31-50	2596	372 (6)	221 (8)	248 (7)	298 (6)	361 (7)	437 (9)	522 (13)	581 (17)	350	45.7 (2.7)
	51-70	2550	353 (5)	205 (6)	232 (5)	280 (5)	341 (6)	414 (8)	494 (11)	550 (14)	350	53.6 (2.3)
	>70	1520	318 (8)	173 (8)	198 (7)	245 (8)	307 (9)	383 (11)	463 (14)	518 (17)	350	65.3 (3.3)
	19+	8470	364 (3)	208 (4)	236 (4)	288 (4)	353 (4)	431 (5)	518 (8)	579 (10)		
Female												
	9-13	1980	265 (5)	157 (5)	177 (5)	215 (5)	260 (5)	311 (7)	366 (9)	400 (11)	200	18.3 (2.1)
	14-18	2256	269 (4)	157 (5)	178 (5)	217 (5)	266 (5)	322 (6)	384 (8)	427 (10)	300	66.3 (2.4)
	19-30	1854	284 (6)	176 (8)	196 (7)	233 (7)	278 (6)	328 (8)	381 (11)	417 (13)	255	36.6 (3.7)
	31-50	2686	306 (5)	171 (5)	195 (4)	238 (4)	296 (5)	365 (7)	441 (11)	497 (15)	265	36.4 (2.1)
	51-70	3200	301 (5)	172 (6)	196 (5)	238 (5)	291 (5)	356 (7)	425 (10)	473 (12)	265	37.5 (2.3)
	>70	2610	268 (4)	157 (5)	177 (5)	214 (5)	263 (5)	320 (6)	381 (8)	423 (10)	265	51.1 (2.7)
	19+	10350	296 (2)	169 (3)	192 (2)	234 (2)	288 (3)	351 (4)	420 (6)	468 (7)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

19. Niacin (NE/d): Usual intakes from food

Table 19.1 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both												
	1-3	79	22.9 (1.4)	17.2 (2.4)	18.3 (2.1)	20.2 (1.7)	22.5 (1.6)	25.0 (2.1)	27.5 (2.9)	29.0 (3.5)	5	<3
	4-8	127	31.9 (2.3)	22.2 (3.1)	23.8 (3.0)	26.8 (2.8)	30.9 (2.8)	35.8 (3.4)	41.1 (5.0)	44.8 (6.8)	6	0.0 (0.0)
Male												
	9-13	111	43.6 (2.9)	32.8 (4.0)	34.7 (3.8)	38.3 (3.6)	42.7 (3.7)	47.5 (4.5)	52.2 (5.8)	55.1 (7.0)	9	0.0 (0.0)
	14-18	107	44.4 (2.8)	33.6 (5.1)	35.8 (4.7)	40.0 (4.0)	45.4 (3.5)	51.7 (4.1)	58.2 (6.1)	62.7 (7.8)	12	0.0 (0.0)
	19-30	77	48.5 (2.2)	40.3 (6.2)	42.3 (5.3)	45.8 (3.9)	49.7 (3.0)	53.7 (3.6)	57.5 (5.2)	59.8 (6.7)	12	<3
	31-50	145	50.8 (5.3)	29.5 (5.1) ^E	33.1 (4.7)	40.0 (4.3)	48.6 (5.1)	58.3 (7.3)	69.5 (10.9)	77.9 (14.0) ^E	12	<3
	51-70	182	40.4 (2.2)	27.2 (2.1)	29.5 (2.2)	33.7 (2.4)	38.9 (2.6)	44.5 (3.0)	49.9 (3.4)	53.4 (3.8)	12	0.0 (0.0)
	>70	63	44.8 (3.1)	35.9 (5.1)	38.4 (5.0)	42.8 (4.8)	48.4 (5.1)	54.6 (6.1)	60.8 (7.8)	64.8 (9.2)	12	<3
	19+	467	46.7 (2.3)	32.1 (4.1)	35.0 (3.7)	40.4 (3.1)	46.6 (2.7)	53.5 (3.0)	60.3 (4.2)	64.9 (5.3)	12	0.0 (0.0)
Female												
	9-13	96	30.9 (1.8)	20.4 (3.2)	22.2 (3.0)	25.7 (2.7)	30.5 (2.5)	36.2 (2.8)	42.0 (3.9)	46.0 (5.0)	9	<3
	14-18	105	36.0 (3.4)	25.3 (4.7) ^E	27.4 (4.5)	31.4 (4.1)	36.5 (4.1)	42.5 (4.8)	48.7 (6.4)	52.7 (7.7)	11	<3
	19-30	91	29.7 (1.6)	21.5 (2.4)	22.9 (2.3)	25.4 (2.1)	28.4 (2.1)	31.6 (2.5)	34.8 (3.3)	36.8 (4.0)	11	<3
	31-50	167	32.4 (2.2)	20.4 (3.2)	22.8 (3.0)	27.2 (2.8)	32.8 (2.8)	39.3 (3.5)	46.1 (4.8)	50.9 (5.8)	11	<3
	51-70	198	33.4 (2.3)	18.9 (4.1) ^E	21.5 (3.5)	25.9 (2.8)	31.6 (2.4)	40.0 (3.6)	49.7 (6.9)	57.3 (10.0) ^E	11	<3
	>70	74	31.9 (2.3)	25.2 (3.9)	26.7 (3.7)	29.4 (3.4)	32.7 (3.3)	36.4 (3.7)	40.1 (4.6)	42.4 (5.4)	11	0.0 (0.0)
	19+	530	32.1 (1.1)	21.1 (1.7)	23.2 (1.6)	27.1 (1.5)	32.0 (1.5)	37.8 (1.7)	44.0 (2.5)	48.3 (3.2)	11	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.2 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	58	24.3 (1.7)	20.3 (1.9)	21.2 (2.0)	22.8 (2.2)	24.7 (2.3)	26.7 (2.5)	28.6 (2.6)	29.8 (2.7)	5	0.0 (0.0)	
	4-8	110	28.1 (1.2)	21.5 (1.3)	23.0 (1.3)	25.4 (1.4)	28.1 (1.5)	30.9 (1.6)	33.7 (1.8)	35.5 (2.0)	6	0.0 (0.0)	
Male													
	9-13	95	36.8 (2.1)	31.5 (4.7)	32.5 (4.3)	34.3 (3.8)	36.3 (3.3)	38.5 (3.3)	40.4 (3.9)	41.7 (4.5)	9	0.0 (0.0)	
	14-18	87	48.7 (3.0)	34.7 (5.7)	37.9 (5.1)	43.6 (4.2)	50.6 (4.0)	58.2 (5.2)	65.5 (7.3)	70.2 (8.9)	12	<3	
	19-30	70	50.1 (2.6)	29.8 (5.1) ^E	34.0 (4.7)	42.2 (3.8)	51.8 (3.3)	61.1 (3.7)	69.7 (5.1)	75.4 (6.3)	12	<3	
	31-50	109	45.7 (2.5)	31.0 (3.7)	33.6 (3.4)	38.6 (3.0)	45.0 (3.0)	52.3 (4.0)	59.8 (5.5)	64.8 (6.8)	12	<3	
	51-70	128	41.3 (2.1)	25.7 (3.1)	28.3 (2.9)	33.4 (2.6)	40.0 (2.6)	47.9 (3.3)	56.3 (4.6)	61.9 (5.9)	12	<3	
	>70	65	31.2 (1.5)	24.2 (2.5)	25.5 (2.3)	27.7 (2.0)	30.3 (2.0)	32.8 (2.7)	35.3 (3.6)	36.8 (4.2)	12	<3	
	19+	372	43.9 (1.4)	27.1 (1.5)	30.1 (1.5)	35.6 (1.5)	42.9 (1.7)	51.7 (2.1)	60.9 (2.7)	67.1 (3.3)	12	<3	
Female													
	9-13	75	32.4 (3.1)	20.0 (4.1) ^E	22.7 (4.0) ^E	27.3 (4.0)	32.5 (4.5)	38.5 (5.6)	45.8 (7.4)	51.0 (9.0) ^E	9	<3	
	14-18	81	31.2 (1.7)	22.7 (3.4)	24.5 (3.0)	27.6 (2.5)	31.1 (2.3)	34.6 (2.7)	38.0 (3.4)	40.0 (4.0)	11	<3	
	19-30	101	36.8 (3.6)	22.0 (3.7) ^E	25.3 (3.6)	31.4 (3.7)	39.1 (4.7)	47.6 (6.6)	55.8 (8.6)	60.7 (9.7)	11	<3	
	31-50	116	31.7 (1.9)	23.0 (3.3)	24.8 (3.0)	27.9 (2.5)	31.5 (2.4)	35.3 (2.7)	38.9 (3.4)	41.1 (4.0)	11	<3	
	51-70	146	33.5 (1.5)	21.5 (2.4)	23.8 (2.3)	28.2 (2.1)	33.7 (2.0)	40.0 (2.3)	46.5 (3.0)	50.7 (3.7)	11	<3	
	>70	94	25.8 (1.4)	17.6 (2.5)	19.1 (2.2)	21.7 (1.9)	24.9 (1.7)	28.4 (2.0)	32.1 (2.8)	34.4 (3.4)	11	<3	
	19+	457	32.4 (1.1)	21.6 (1.5)	23.9 (1.4)	28.0 (1.3)	32.9 (1.5)	38.0 (1.8)	43.2 (2.4)	46.5 (2.8)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.3 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	112	24.0 (1.7)	17.5 (2.7)	18.6 (2.5)	20.8 (2.1)	23.5 (1.9)	26.7 (2.3)	30.0 (3.3)	32.3 (4.2)	5	0.0	(0.0)
	4-8	177	29.1 (1.1)	20.4 (1.6)	22.1 (1.5)	24.9 (1.5)	28.3 (1.5)	32.0 (1.5)	35.7 (1.7)	38.0 (1.7)	6	<3	
Male													
	9-13	111	38.6 (2.0)	24.7 (3.2)	26.9 (3.0)	31.0 (2.6)	36.5 (2.3)	43.0 (2.9)	50.0 (4.4)	54.8 (5.6)	9	<3	
	14-18	113	44.4 (4.6)	28.9 (5.4) ^E	31.4 (5.7) ^E	36.1 (6.2) ^E	42.2 (6.7)	49.4 (7.1)	57.0 (7.6)	62.1 (8.1)	12	0.0	(0.0)
	19-30	91	48.1 (3.1)	28.1 (6.3) ^E	31.7 (5.6) ^E	38.3 (4.5)	46.3 (3.9)	54.9 (4.6)	63.2 (6.5)	68.5 (8.1)	12	<3	
	31-50	101	47.7 (2.5)	33.7 (5.1)	36.6 (4.5)	41.7 (3.7)	48.0 (3.4)	54.8 (4.3)	61.3 (5.8)	65.5 (7.0)	12	<3	
	51-70	134	45.1 (2.8)	35.1 (2.4)	37.1 (2.4)	40.6 (2.5)	44.5 (2.8)	49.0 (3.3)	53.6 (3.7)	56.8 (3.9)	12	0.0	(0.0)
	>70	56	35.8 (2.6)	24.6 (4.3) ^E	26.6 (4.0)	30.4 (3.5)	35.3 (3.1)	41.2 (4.1)	47.4 (6.6)	51.6 (8.8) ^E	12	<3	
	19+	382	45.8 (1.5)	30.7 (2.5)	33.6 (2.3)	38.8 (1.9)	45.1 (1.9)	52.4 (2.4)	59.8 (3.3)	64.5 (4.0)	12	<3	
Female													
	9-13	105	32.5 (2.6)	23.9 (3.6)	25.6 (3.4)	28.6 (3.0)	32.4 (2.9)	36.7 (4.0)	41.2 (5.3)	44.0 (6.2)	9	<3	
	14-18	120	28.6 (2.7)	17.3 (4.5) ^E	19.6 (4.0) ^E	23.7 (3.3)	28.6 (2.8)	34.2 (3.2)	39.8 (4.5)	43.6 (5.6)	11	^F	
	19-30	91	33.5 (2.0)	25.4 (1.9)	27.1 (2.0)	30.2 (2.1)	33.6 (2.2)	36.9 (2.2)	40.0 (2.2)	41.8 (2.3)	11	0.0	(0.0)
	31-50	159	35.0 (2.1)	20.4 (3.2)	23.4 (3.0)	28.8 (2.8)	35.1 (2.7)	41.4 (2.9)	47.3 (3.3)	51.1 (3.7)	11	<3	
	51-70	174	32.8 (2.0)	21.9 (1.6)	23.8 (1.7)	27.2 (1.9)	31.5 (2.2)	36.5 (2.5)	41.4 (2.7)	44.4 (2.9)	11	0.0	(0.0)
	>70	80	29.2 (2.0)	19.8 (3.3) ^E	22.1 (2.9)	25.9 (2.5)	29.9 (2.4)	34.4 (3.1)	39.8 (4.7)	43.8 (6.0)	11	<3	
	19+	504	33.3 (1.1)	21.2 (1.8)	23.7 (1.6)	28.0 (1.4)	33.2 (1.4)	39.0 (1.8)	44.7 (2.5)	48.3 (2.9)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.4 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	99	26.0 (1.9)	17.4 (2.2)	18.9 (2.1)	21.7 (2.0)	25.5 (2.1)	30.4 (2.8)	35.9 (4.2)	39.7 (5.4)	5	<3	
	4-8	140	32.1 (2.4)	22.9 (3.2)	24.6 (3.0)	27.8 (2.6)	31.9 (2.5)	36.9 (3.1)	42.3 (4.6)	46.1 (5.9)	6	0.0 (0.0)	
Male													
	9-13	92	38.2 (2.5)	26.7 (3.7)	28.6 (3.5)	32.2 (3.1)	36.6 (3.0)	41.8 (3.5)	47.4 (4.9)	51.4 (6.3)	9	0.0 (0.0)	
	14-18	107	50.5 (3.0)	33.0 (4.2)	36.3 (3.9)	42.8 (3.5)	51.4 (3.7)	61.2 (4.9)	72.1 (7.2)	79.8 (9.2)	12	<3	
	19-30	73	56.8 (5.5)	33.6 (5.8) ^E	37.9 (5.4)	45.6 (5.0)	56.0 (5.6)	69.6 (8.4)	85.1 (13.0)	95.7 (16.4) ^E	12	<3	
	31-50	134	48.0 (2.9)	32.5 (2.7)	35.0 (2.9)	39.6 (3.1)	45.6 (3.4)	52.6 (3.9)	59.9 (4.6)	64.8 (5.2)	12	0.0 (0.0)	
	51-70	131	43.4 (2.7)	31.7 (2.7)	33.8 (2.7)	37.6 (2.8)	42.3 (3.0)	47.5 (3.4)	52.6 (3.8)	55.9 (4.1)	12	0.0 (0.0)	
	>70	55	35.3 (2.1)	22.1 (3.6)	24.5 (3.3)	28.8 (2.9)	33.8 (2.7)	39.2 (3.0)	44.2 (3.8)	47.4 (4.6)	12	<3	
	19+	393	47.3 (1.7)	31.5 (3.3)	34.3 (3.0)	39.5 (2.4)	46.3 (2.2)	54.6 (3.0)	63.6 (4.9)	69.6 (6.4)	12	0.0 (0.0)	
Female													
	9-13	79	33.7 (2.9)	24.0 (4.0) ^E	26.1 (3.9)	30.1 (3.6)	35.1 (3.7)	40.8 (4.8)	46.7 (7.1)	50.7 (9.1) ^E	9	<3	
	14-18	104	32.6 (1.8)	24.2 (4.2) ^E	26.6 (3.6)	30.2 (2.8)	33.8 (2.4)	37.3 (2.9)	41.2 (4.2)	44.0 (5.4)	11	<3	
	19-30	101	33.2 (2.0)	24.9 (2.9)	27.0 (2.7)	30.9 (2.4)	35.8 (2.4)	41.6 (3.1)	47.5 (4.4)	51.3 (5.3)	11	<3	
	31-50	143	31.2 (1.9)	17.4 (3.4) ^E	20.0 (3.0)	24.5 (2.3)	29.7 (2.0)	35.4 (2.7)	41.5 (4.2)	45.8 (5.6)	11	<3	
	51-70	193	31.4 (1.9)	22.3 (3.4)	23.9 (3.1)	26.8 (2.6)	30.7 (2.4)	35.2 (3.1)	40.0 (4.8)	43.4 (6.5)	11	<3	
	>70	94	29.4 (2.0)	21.5 (3.1)	22.9 (2.9)	25.4 (2.7)	28.5 (2.7)	31.7 (3.1)	34.9 (3.8)	36.9 (4.5)	11	<3	
	19+	531	31.4 (0.9)	20.9 (1.7)	22.9 (1.5)	26.5 (1.2)	30.9 (1.0)	36.0 (1.5)	41.3 (2.4)	44.8 (3.3)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.5 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	311	25.8	(1.3)	19.3 (2.9)	20.7 (2.6)	23.1 (2.1)	26.0 (1.8)	29.2 (1.9)	32.3 (2.4)	34.4 (2.9)	5	0.0 (0.0)
	4-8	485	32.8	(1.2)	24.1 (2.7)	25.8 (2.3)	29.0 (1.8)	32.9 (1.5)	37.0 (1.8)	40.9 (2.7)	43.4 (3.4)	6	0.0 (0.0)
Male													
	9-13	277	45.6	(3.1)	32.4 (4.8)	35.4 (4.5)	40.7 (4.1)	47.5 (4.1)	55.4 (4.9)	63.9 (6.7)	69.7 (8.3)	9	0.0 (0.0)
	14-18	339	51.8	(2.4)	32.3 (3.0)	36.0 (2.9)	43.1 (2.7)	52.6 (3.0)	64.5 (4.0)	77.7 (5.7)	87.0 (7.2)	12	0.0 (0.0)
	19-30	237	48.8	(2.3)	38.8 (4.5)	41.2 (4.0)	45.3 (3.3)	50.3 (2.8)	55.8 (3.4)	61.1 (4.9)	64.4 (6.2)	12	0.0 (0.0)
	31-50	423	48.6	(2.1)	33.4 (4.7)	36.6 (4.1)	42.5 (3.0)	48.9 (2.5)	56.4 (3.5)	63.7 (5.6)	68.5 (7.1)	12	<3
	51-70	387	44.0	(1.6)	27.4 (3.4)	30.5 (3.1)	36.1 (2.5)	43.5 (1.9)	51.9 (2.2)	60.5 (3.4)	66.4 (4.5)	12	<3
	>70	132	37.0	(2.3)	21.1 (4.1) ^E	23.7 (4.0) ^E	28.8 (3.9)	35.8 (3.9)	44.4 (4.3)	54.0 (5.1)	60.6 (5.9)	12	<3
	19+	1179	46.4	(1.2)	29.3 (1.7)	32.3 (1.7)	38.6 (1.5)	46.4 (1.5)	55.0 (1.8)	64.1 (2.6)	70.3 (3.3)	12	0.0 (0.0)
Female													
	9-13	281	33.9	(1.6)	20.9 (2.5)	23.1 (2.3)	27.2 (2.0)	32.6 (1.9)	38.8 (2.3)	45.3 (3.3)	49.6 (4.1)	9	<3
	14-18	321	34.8	(1.1)	23.6 (1.8)	26.0 (1.6)	30.2 (1.5)	35.5 (1.6)	41.5 (2.0)	47.6 (2.6)	51.4 (3.1)	11	<3
	19-30	249	35.6	(1.7)	28.5 (3.2)	30.0 (2.8)	32.5 (2.3)	35.4 (2.0)	38.5 (2.5)	41.4 (3.5)	43.2 (4.3)	11	<3
	31-50	364	41.0	(2.8)	25.4 (4.4) ^E	28.0 (4.1)	33.3 (3.6)	40.5 (3.3)	49.0 (3.9)	58.3 (5.6)	64.8 (7.2)	11	<3
	51-70	467	36.4	(1.5)	24.6 (3.0)	26.9 (2.6)	31.0 (2.0)	36.1 (1.7)	42.0 (2.3)	48.0 (3.6)	52.1 (4.7)	11	<3
	>70	215	28.6	(1.2)	20.0 (1.8)	21.6 (1.7)	24.5 (1.6)	28.1 (1.6)	32.1 (2.0)	36.2 (2.6)	39.1 (3.3)	11	<3
	19+	1295	37.1	(1.2)	24.6 (1.5)	27.0 (1.5)	31.3 (1.4)	36.8 (1.4)	43.3 (1.7)	50.2 (2.3)	54.9 (2.9)	11	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.6 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	644	22.9	(0.6)	15.3 (1.6)	16.9 (1.4)	19.6 (1.0)	23.0 (0.6)	26.7 (1.0)	30.4 (1.8)	32.7 (2.3)	5	<3	
	4-8	956	28.8	(0.6)	20.7 (1.6)	22.3 (1.4)	25.1 (1.0)	28.5 (0.7)	32.4 (1.0)	36.3 (1.8)	38.9 (2.4)	6	0.0	(0.0)
Male														
	9-13	589	37.7	(0.9)	27.6 (3.3)	29.7 (2.8)	33.4 (1.9)	37.5 (1.1)	42.0 (1.7)	46.7 (3.2)	49.9 (4.4)	9	0.0	(0.0)
	14-18	639	46.8	(1.3)	30.6 (3.4)	33.7 (3.0)	39.5 (2.3)	46.6 (1.7)	54.9 (2.1)	63.5 (3.6)	69.3 (4.9)	12	<3	
	19-30	481	49.4	(1.9)	30.3 (5.6) ^E	33.4 (4.9)	39.3 (3.7)	47.3 (2.5)	56.9 (3.3)	66.8 (6.1)	73.4 (8.4)	12	<3	
	31-50	709	47.5	(1.4)	31.1 (4.5)	34.0 (4.0)	39.5 (2.9)	46.7 (1.7)	55.2 (2.4)	64.0 (4.7)	69.8 (6.5)	12	0.0	(0.0)
	51-70	758	42.9	(1.1)	30.3 (3.5)	32.6 (3.0)	36.8 (2.1)	42.1 (1.3)	48.0 (1.9)	54.0 (3.7)	58.0 (5.1)	12	0.0	(0.0)
	>70	734	34.2	(0.8)	22.2 (1.9)	24.4 (1.7)	28.4 (1.2)	33.4 (0.9)	39.0 (1.2)	44.8 (2.0)	48.7 (2.7)	12	<3	
	19+	2682	45.5	(0.8)	29.0 (2.0)	31.7 (1.8)	37.0 (1.5)	44.5 (1.0)	53.0 (1.3)	62.0 (2.4)	68.3 (3.5)	12	0.0	(0.0)
Female														
	9-13	585	31.0	(0.8)	20.9 (2.2)	22.8 (1.8)	26.1 (1.2)	30.0 (0.8)	34.4 (1.4)	39.1 (2.6)	42.2 (3.4)	9	<3	
	14-18	645	33.2	(0.8)	19.3 (2.2)	21.9 (1.9)	26.7 (1.4)	32.7 (0.9)	39.5 (1.4)	46.2 (2.5)	50.5 (3.3)	11	<3	
	19-30	514	32.0	(1.1)	19.0 (3.0)	21.3 (2.6)	25.6 (1.8)	30.9 (1.2)	36.9 (1.8)	43.0 (3.2)	47.1 (4.4)	11	<3	
	31-50	758	33.9	(0.8)	20.2 (1.5)	22.6 (1.3)	27.0 (1.1)	33.2 (1.0)	40.5 (1.4)	47.3 (2.1)	51.7 (2.7)	11	<3	
	51-70	955	32.8	(1.0)	23.9 (2.7)	25.6 (2.3)	28.7 (1.7)	32.4 (1.1)	36.4 (1.3)	40.4 (2.3)	43.0 (3.2)	11	<3	
	>70	1345	29.1	(0.7)	17.2 (1.0)	19.2 (0.9)	22.9 (0.8)	27.5 (0.7)	33.2 (0.9)	39.5 (1.4)	43.9 (1.9)	11	<3	
	19+	3572	32.7	(0.5)	19.8 (0.9)	22.1 (0.8)	26.4 (0.6)	31.8 (0.6)	38.0 (0.7)	44.5 (1.1)	48.9 (1.5)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.7 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	324	22.0	(0.9)	14.1 (1.4)	15.6 (1.3)	18.3 (1.1)	21.6 (1.0)	25.5 (1.3)	29.8 (2.1)	32.9 (2.8)	5	<3
	4-8	425	27.5	(1.3)	21.2 (2.8)	22.5 (2.4)	24.6 (1.9)	27.3 (1.4)	30.3 (1.7)	33.4 (2.8)	35.4 (3.8)	6	0.0 (0.0)
Male													
	9-13	274	36.3	(1.0)	25.5 (3.0)	27.8 (2.5)	31.8 (1.8)	36.5 (1.4)	41.6 (1.9)	46.9 (3.1)	50.3 (4.0)	9	<3
	14-18	297	47.5	(1.6)	26.4 (2.6)	30.2 (2.4)	37.1 (2.1)	46.0 (2.1)	57.2 (2.5)	69.1 (3.3)	76.9 (4.1)	12	<3
	19-30	249	51.3	(3.2)	36.8 (6.8) ^E	39.4 (6.1)	44.1 (4.7)	49.9 (3.4)	56.1 (4.6)	62.2 (8.0)	66.0 (10.6)	12	<3
	31-50	309	44.1	(2.5)	26.5 (4.1)	29.7 (3.6)	35.4 (2.7)	41.8 (2.4)	49.3 (3.3)	58.2 (5.4)	64.7 (7.3)	12	<3
	51-70	277	42.1	(1.3)	26.2 (2.2)	29.0 (2.0)	34.0 (1.7)	40.4 (1.6)	48.3 (2.0)	56.5 (2.9)	61.7 (3.6)	12	<3
	>70	136	37.5	(2.5)	16.9 (4.7) ^E	20.1 (4.2) ^E	26.6 (3.3)	35.2 (2.8)	45.8 (4.6)	57.7 (7.3)	65.3 (9.3)	12	^F
	19+	971	44.5	(1.2)	26.9 (2.2)	29.6 (2.1)	35.2 (1.8)	42.9 (1.3)	50.7 (1.7)	60.0 (2.8)	66.0 (3.7)	12	<3
Female													
	9-13	265	31.2	(1.6)	22.2 (3.4)	24.0 (2.9)	27.0 (2.3)	30.6 (1.9)	34.6 (2.3)	38.7 (3.5)	41.4 (4.6)	9	<3
	14-18	290	32.0	(1.2)	24.1 (2.9)	25.7 (2.4)	28.5 (1.7)	31.7 (1.4)	35.0 (2.2)	38.1 (3.3)	40.1 (4.1)	11	<3
	19-30	197	35.0	(2.5)	22.3 (2.0)	24.7 (2.1)	29.1 (2.3)	34.3 (2.5)	39.9 (2.7)	45.0 (3.0)	48.1 (3.2)	11	<3
	31-50	312	36.3	(1.6)	22.3 (3.2)	25.0 (2.8)	29.3 (2.2)	35.0 (1.9)	41.8 (2.6)	48.3 (4.0)	52.9 (5.2)	11	<3
	51-70	312	33.4	(1.7)	22.5 (2.9)	24.5 (2.6)	28.1 (2.2)	32.7 (1.9)	37.9 (2.3)	43.3 (3.4)	46.8 (4.4)	11	<3
	>70	239	26.4	(1.2)	16.5 (2.5)	18.3 (2.2)	21.6 (1.7)	26.0 (1.2)	31.0 (1.7)	36.3 (2.9)	39.7 (3.8)	11	<3
	19+	1060	33.8	(0.9)	19.9 (1.4)	22.3 (1.3)	26.8 (1.1)	32.7 (1.0)	39.5 (1.4)	46.5 (2.2)	51.2 (2.8)	11	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.8 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	129	21.4 (0.9)	14.8 (1.8)	16.1 (1.6)	18.4 (1.4)	21.3 (1.3)	24.5 (1.4)	27.7 (1.8)	29.7 (2.2)	5	<3	
	4-8	213	29.7 (1.5)	20.8 (2.0)	22.5 (1.8)	25.5 (1.5)	29.3 (1.4)	33.6 (2.0)	38.2 (3.2)	41.4 (4.1)	6	0.0	(0.0)
Male													
	9-13	122	39.4 (2.9)	29.9 (2.4)	31.6 (2.6)	34.9 (3.0)	39.1 (3.7)	44.3 (4.5)	49.8 (5.3)	53.4 (5.7)	9	0.0	(0.0)
	14-18	150	51.6 (2.8)	31.7 (3.8)	35.4 (3.6)	42.1 (3.4)	50.7 (3.5)	61.1 (4.3)	72.2 (5.8)	79.5 (7.2)	12	<3	
	19-30	106	48.6 (3.0)	22.0 (4.7) ^E	26.8 (4.2)	35.5 (3.6)	46.2 (3.5)	58.4 (4.5)	71.2 (6.5)	79.8 (8.3)	12	<3	
	31-50	155	44.8 (2.4)	23.3 (4.3) ^E	27.5 (3.8)	34.8 (3.2)	43.6 (3.0)	53.8 (3.4)	63.9 (4.5)	70.3 (5.5)	12	<3	
	51-70	122	42.1 (2.5)	27.4 (4.4)	30.6 (3.7)	36.2 (3.0)	42.9 (3.0)	50.5 (4.5)	58.5 (6.8)	64.1 (8.6)	12	<3	
	>70	88	38.2 (2.0)	23.4 (3.1)	25.8 (2.9)	30.3 (2.6)	36.6 (2.4)	44.4 (3.3)	53.0 (5.2)	59.1 (6.8)	12	<3	
	19+	471	44.2 (1.3)	23.2 (1.8)	27.1 (1.6)	34.1 (1.5)	42.8 (1.6)	53.3 (2.1)	64.3 (2.8)	71.6 (3.5)	12	<3	
Female													
	9-13	103	32.2 (1.9)	22.7 (2.8)	24.5 (2.5)	27.8 (2.3)	31.7 (2.3)	36.1 (2.7)	40.9 (3.7)	44.1 (4.5)	9	0.0	(0.0)
	14-18	142	32.8 (1.3)	21.0 (1.9)	23.4 (1.7)	27.6 (1.6)	32.5 (1.9)	38.0 (2.5)	43.6 (3.3)	47.4 (4.0)	11	<3	
	19-30	111	29.9 (1.3)	20.1 (2.9)	21.9 (2.5)	25.1 (2.0)	28.9 (1.8)	33.1 (2.2)	37.1 (3.2)	39.7 (3.9)	11	<3	
	31-50	146	35.6 (1.9)	17.6 (3.0) ^E	20.7 (2.9)	26.7 (2.7)	33.8 (2.5)	40.6 (2.8)	46.9 (3.3)	51.0 (3.9)	11	<3	
	51-70	184	33.1 (1.6)	22.5 (1.2)	24.5 (1.3)	27.9 (1.6)	32.2 (1.9)	37.0 (2.4)	42.0 (2.9)	45.4 (3.3)	11	0.0	(0.0)
	>70	143	31.7 (2.0)	17.0 (2.3)	19.5 (2.2)	24.5 (2.2)	31.4 (2.4)	40.3 (3.0)	50.3 (4.5)	57.4 (6.0)	11	<3	
	19+	584	33.1 (1.0)	19.7 (1.4)	22.1 (1.4)	26.5 (1.3)	32.2 (1.2)	38.5 (1.5)	44.7 (2.0)	48.6 (2.5)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.9 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	169	20.7 (0.9)	12.2 (1.3)	13.7 (1.2)	16.5 (1.1)	20.0 (1.0)	24.1 (1.4)	28.4 (2.0)	31.4 (2.6)	5	<3	
	4-8	281	28.0 (0.9)	18.6 (2.4)	20.3 (2.1)	23.3 (1.5)	27.0 (1.0)	31.3 (1.5)	35.7 (2.6)	38.6 (3.4)	6	0.0 (0.0)	
Male	9-13	183	40.0 (2.3)	27.4 (4.0)	29.7 (3.7)	34.1 (3.1)	39.8 (2.7)	46.7 (3.3)	54.1 (5.1)	59.2 (6.7)	9	0.0 (0.0)	
	14-18	187	49.0 (2.3)	37.4 (5.7)	40.1 (5.0)	44.8 (3.8)	50.6 (3.0)	57.0 (4.0)	63.3 (6.5)	67.3 (8.5)	12	<3	
	19-30	223	49.0 (2.4)	36.9 (5.5)	39.3 (4.7)	43.6 (3.5)	48.6 (2.8)	53.9 (3.7)	59.2 (5.8)	62.5 (7.4)	12	<3	
	31-50	229	46.9 (2.7)	29.8 (2.2)	33.1 (2.2)	39.1 (2.4)	46.8 (3.1)	56.1 (4.1)	66.3 (5.2)	73.5 (5.9)	12	<3	
	51-70	197	42.0 (2.3)	22.7 (4.2) ^E	26.3 (3.8)	33.1 (3.2)	41.3 (2.8)	50.0 (3.2)	58.0 (4.3)	63.1 (5.3)	12	<3	
	>70	72	39.2 (3.6)	30.6 (5.0)	32.2 (4.7)	35.1 (4.1)	38.5 (3.9)	41.9 (4.1)	45.3 (5.0)	47.3 (5.7)	12	<3	
	19+	721	45.7 (1.5)	29.7 (2.9)	32.9 (2.5)	38.7 (1.9)	45.5 (1.7)	53.5 (2.4)	62.0 (3.8)	67.8 (5.0)	12	<3	
	Female	9-13	165	31.5 (1.7)	22.7 (3.7)	24.6 (3.2)	28.0 (2.5)	32.3 (2.1)	37.2 (2.9)	42.0 (4.5)	45.2 (5.8)	9	<3
14-18	206	31.5 (1.5)	21.7 (2.7)	23.6 (2.3)	27.0 (1.8)	30.8 (1.7)	35.2 (2.4)	39.8 (3.5)	42.8 (4.3)	11	<3		
19-30	191	30.3 (1.7)	18.0 (2.8)	20.4 (2.6)	24.6 (2.2)	29.5 (2.2)	35.0 (2.6)	40.7 (3.7)	44.5 (4.6)	11	<3		
31-50	258	32.6 (1.4)	22.1 (1.6)	24.4 (1.6)	28.4 (1.8)	33.2 (2.0)	38.7 (2.5)	44.2 (3.0)	47.8 (3.4)	11	<3		
51-70	249	31.3 (1.4)	19.1 (2.4)	21.3 (2.2)	25.4 (1.8)	30.6 (1.6)	36.8 (2.1)	43.4 (3.2)	47.9 (4.2)	11	<3		
>70	128	34.7 (3.5)	21.4 (2.9)	24.3 (2.9)	29.3 (3.0)	34.1 (3.3)	39.8 (4.3)	47.1 (5.5)	51.9 (6.3)	11	<3		
19+	826	32.0 (0.8)	20.6 (1.5)	22.9 (1.4)	27.0 (1.2)	32.0 (1.1)	37.7 (1.3)	43.6 (1.7)	47.5 (2.1)	11	<3		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.10 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR (SE)	
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	24.4 (1.0)	19.8 (2.1)	20.7 (1.9)	22.2 (1.4)	23.9 (1.3)	26.1 (1.5)	28.3 (2.1)	29.7 (2.6)	5	0.0 (0.0)	
	4-8	321	29.3 (0.8)	20.1 (1.7)	21.9 (1.5)	25.2 (1.1)	29.1 (1.0)	33.3 (1.4)	37.4 (2.1)	40.0 (2.5)	6	0.0 (0.0)	
Male													
	9-13	226	36.8 (1.3)	23.2 (2.0)	25.8 (1.9)	30.5 (1.7)	36.2 (1.7)	42.6 (2.1)	49.8 (2.9)	55.1 (3.8)	9	<3	
	14-18	262	50.8 (3.8)	29.8 (5.2) ^E	33.7 (4.7)	40.6 (3.9)	49.2 (3.6)	59.3 (4.6)	70.1 (7.0)	77.6 (9.2)	12	<3	
	19-30	197	51.7 (2.5)	36.1 (4.4)	39.0 (3.9)	44.3 (3.2)	51.1 (3.0)	59.0 (4.3)	67.3 (6.7)	73.0 (8.5)	12	0.0 (0.0)	
	31-50	282	53.7 (2.5)	31.6 (3.9)	35.7 (3.5)	43.2 (3.1)	52.8 (2.9)	63.4 (3.5)	73.6 (5.0)	80.2 (6.2)	12	<3	
	51-70	234	49.0 (2.3)	29.7 (3.3)	32.7 (3.1)	38.6 (2.8)	46.3 (2.8)	55.6 (3.6)	65.6 (5.4)	72.5 (6.9)	12	0.0 (0.0)	
	>70	119	38.7 (2.7)	22.6 (3.9) ^E	25.5 (3.7)	30.8 (3.5)	37.6 (3.7)	45.3 (4.3)	53.0 (5.5)	58.1 (6.5)	12	<3	
	19+	832	50.5 (1.3)	30.1 (1.5)	33.6 (1.5)	40.3 (1.4)	49.2 (1.6)	59.5 (2.1)	70.1 (2.9)	77.0 (3.5)	12	<3	
Female													
	9-13	226	32.3 (2.0)	21.2 (2.2)	23.2 (2.0)	26.9 (1.9)	31.6 (2.2)	37.5 (3.0)	43.8 (4.2)	48.0 (5.2)	9	<3	
	14-18	242	32.8 (1.5)	18.2 (2.3)	20.6 (2.3)	25.4 (2.0)	31.5 (1.9)	38.0 (2.3)	44.8 (3.1)	49.4 (3.8)	11	<3	
	19-30	208	34.6 (1.9)	24.1 (3.6)	26.3 (3.2)	30.3 (2.7)	35.1 (2.5)	40.1 (3.1)	44.8 (4.3)	47.8 (5.2)	11	<3	
	31-50	263	37.2 (1.8)	21.3 (2.2)	24.1 (2.0)	29.0 (1.8)	35.3 (2.0)	43.1 (2.9)	51.8 (4.3)	57.8 (5.6)	11	<3	
	51-70	322	34.2 (2.1)	21.6 (3.4)	23.9 (3.1)	28.2 (2.6)	33.7 (2.2)	39.9 (2.9)	46.5 (4.4)	51.0 (5.6)	11	<3	
	>70	198	31.8 (2.0)	20.4 (3.5) ^E	22.5 (3.3)	26.4 (2.9)	31.3 (2.7)	37.1 (3.1)	43.3 (4.5)	47.6 (5.9)	11	<3	
	19+	991	35.2 (0.9)	21.3 (1.2)	23.7 (1.2)	28.3 (1.1)	34.7 (1.3)	41.5 (1.6)	49.1 (2.3)	54.3 (2.9)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.11 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	348	24.4	(0.9)	19.3 (1.8)	20.3 (1.6)	22.2 (1.3)	24.5 (1.1)	27.1 (1.4)	29.7 (2.2)	31.4 (2.8)	5	0.0 (0.0)
	4-8	554	30.6	(1.1)	22.3 (1.8)	24.0 (1.7)	27.0 (1.4)	30.8 (1.3)	35.2 (1.7)	39.9 (2.7)	43.2 (3.6)	6	0.0 (0.0)
Male													
	9-13	409	39.4	(1.4)	29.1 (2.2)	30.9 (2.0)	34.1 (1.7)	38.3 (1.6)	43.0 (2.0)	47.7 (2.8)	50.9 (3.5)	9	0.0 (0.0)
	14-18	414	46.6	(2.1)	31.4 (3.3)	34.9 (3.0)	40.1 (2.8)	46.9 (2.8)	54.5 (3.3)	61.8 (4.6)	66.9 (5.5)	12	0.0 (0.0)
	19-30	311	51.1	(2.3)	32.4 (2.8)	36.1 (2.6)	42.6 (2.3)	50.7 (2.4)	59.9 (3.3)	69.8 (4.9)	76.7 (6.3)	12	<3
	31-50	489	48.4	(1.8)	32.3 (2.8)	35.1 (2.6)	40.4 (2.3)	47.4 (2.2)	55.6 (3.0)	64.2 (4.5)	69.9 (5.7)	12	<3
	51-70	575	43.3	(1.5)	31.8 (3.2)	34.1 (2.8)	37.9 (2.0)	42.2 (1.6)	47.0 (2.0)	51.9 (3.2)	55.1 (4.1)	12	0.0 (0.0)
	>70	239	37.2	(1.4)	25.7 (2.1)	28.0 (2.0)	32.2 (1.9)	37.4 (1.9)	43.2 (2.3)	49.2 (3.2)	53.3 (3.9)	12	0.0 (0.0)
	19+	1614	46.4	(1.0)	29.9 (1.2)	32.9 (1.1)	38.4 (1.1)	45.4 (1.2)	53.9 (1.6)	63.0 (2.2)	69.2 (2.8)	12	0.0 (0.0)
Female													
	9-13	355	32.5	(1.5)	22.1 (2.3)	24.1 (2.2)	27.9 (2.0)	32.7 (1.9)	38.4 (2.2)	44.4 (2.9)	48.4 (3.6)	9	0.0 (0.0)
	14-18	410	31.7	(1.4)	20.6 (2.3)	22.9 (2.1)	26.9 (1.7)	31.7 (1.6)	37.2 (2.0)	42.8 (2.8)	46.4 (3.5)	11	<3
	19-30	384	32.8	(1.1)	24.2 (1.6)	26.1 (1.5)	29.7 (1.3)	34.0 (1.4)	38.5 (1.7)	42.8 (2.3)	45.6 (2.8)	11	0.0 (0.0)
	31-50	585	33.0	(1.1)	20.2 (1.7)	22.9 (1.6)	27.4 (1.5)	32.8 (1.5)	38.9 (1.8)	45.1 (2.2)	49.0 (2.6)	11	<3
	51-70	711	32.5	(1.2)	21.0 (1.8)	23.0 (1.6)	26.5 (1.3)	31.2 (1.3)	37.0 (1.8)	43.5 (2.9)	48.0 (3.9)	11	<3
	>70	342	29.6	(1.2)	19.6 (1.7)	21.6 (1.6)	25.1 (1.5)	29.5 (1.6)	34.5 (1.8)	39.5 (2.3)	42.8 (2.7)	11	<3
	19+	2022	32.4	(0.6)	21.3 (0.9)	23.5 (0.8)	27.4 (0.8)	32.2 (0.8)	37.8 (1.0)	43.7 (1.3)	47.7 (1.6)	11	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.12 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	622	21.1	(0.5)	13.6 (0.9)	15.1 (0.8)	17.6 (0.8)	20.7 (0.7)	24.3 (0.8)	28.1 (1.2)	30.6 (1.5)	5	0.0 (0.0)
	4-8	919	28.2	(0.7)	19.1 (1.4)	20.7 (1.2)	23.7 (0.9)	27.4 (0.7)	31.7 (1.0)	36.1 (1.7)	38.9 (2.2)	6	0.0 (0.0)
Male													
	9-13	579	39.1	(1.5)	29.6 (3.0)	31.5 (2.7)	34.8 (2.1)	39.2 (1.7)	44.3 (2.1)	49.7 (3.4)	53.3 (4.4)	9	0.0 (0.0)
	14-18	634	49.2	(1.6)	29.5 (2.1)	33.2 (2.0)	39.9 (1.8)	48.8 (2.0)	59.5 (2.5)	70.8 (3.6)	78.3 (4.5)	12	<3
	19-30	578	49.4	(1.7)	29.9 (3.7)	33.4 (3.2)	39.6 (2.5)	47.6 (2.0)	57.0 (2.8)	66.8 (4.6)	73.3 (6.0)	12	<3
	31-50	693	46.0	(1.8)	28.3 (3.3)	31.7 (2.9)	37.6 (2.4)	45.1 (2.0)	54.2 (2.6)	64.0 (4.2)	70.7 (5.5)	12	<3
	51-70	596	42.0	(1.5)	23.5 (2.1)	27.1 (2.0)	33.5 (1.9)	41.4 (1.8)	50.2 (1.9)	59.0 (2.4)	64.7 (2.9)	12	<3
	>70	296	38.5	(2.0)	22.6 (2.4)	25.3 (2.3)	30.4 (2.2)	37.1 (2.3)	44.7 (2.8)	52.8 (3.8)	58.3 (4.6)	12	<3
	19+	2163	45.2	(1.0)	26.4 (1.3)	29.9 (1.2)	36.3 (1.1)	44.2 (1.1)	53.5 (1.4)	63.6 (2.0)	70.7 (2.6)	12	<3
Female													
	9-13	533	31.6	(1.2)	23.8 (2.5)	25.5 (2.2)	28.4 (1.8)	31.9 (1.5)	35.8 (1.8)	39.7 (2.6)	42.3 (3.3)	9	0.0 (0.0)
	14-18	638	31.8	(1.0)	21.1 (1.6)	23.1 (1.4)	26.8 (1.2)	31.2 (1.2)	36.2 (1.5)	41.1 (2.0)	44.3 (2.5)	11	<3
	19-30	499	31.2	(1.2)	20.1 (2.4)	22.2 (2.1)	26.0 (1.6)	30.4 (1.4)	35.2 (1.8)	40.0 (2.6)	43.1 (3.2)	11	<3
	31-50	716	33.9	(1.0)	21.9 (2.3)	24.4 (2.1)	28.7 (1.7)	33.9 (1.4)	39.7 (1.6)	45.4 (2.2)	49.0 (2.7)	11	<3
	51-70	745	32.1	(0.9)	21.8 (1.8)	23.7 (1.7)	27.3 (1.3)	31.6 (1.1)	36.6 (1.4)	41.8 (2.2)	45.3 (2.9)	11	<3
	>70	510	31.8	(1.8)	17.1 (1.3)	19.8 (1.3)	24.8 (1.4)	31.0 (1.7)	38.7 (2.3)	47.8 (3.2)	54.2 (3.8)	11	<3
	19+	2470	32.6	(0.6)	19.9 (0.9)	22.3 (0.9)	26.7 (0.8)	32.1 (0.8)	38.4 (0.9)	44.9 (1.1)	49.2 (1.4)	11	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 19.13 Niacin (NE/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	23.5 (0.4)	16.0 (0.8)	17.5 (0.7)	20.1 (0.6)	23.4 (0.5)	27.1 (0.6)	30.9 (0.9)	33.5 (1.1)	5	0.0 (0.0)	
	4-8	3235	29.8 (0.4)	20.1 (0.6)	21.9 (0.5)	25.2 (0.5)	29.4 (0.5)	34.1 (0.7)	38.8 (0.9)	42.0 (1.2)	6	0.0 (0.0)	
Male													
	9-13	2080	39.8 (0.9)	28.2 (1.5)	30.4 (1.4)	34.5 (1.2)	39.5 (1.0)	45.4 (1.2)	51.6 (1.7)	55.7 (2.1)	9	0.0 (0.0)	
	14-18	2288	48.8 (1.0)	30.2 (1.3)	33.8 (1.2)	40.4 (1.1)	48.7 (1.1)	58.5 (1.4)	69.3 (2.1)	76.7 (2.7)	12	0.0 (0.0)	
	19-30	1804	49.7 (1.1)	30.8 (1.6)	34.3 (1.5)	40.5 (1.3)	48.6 (1.3)	58.0 (1.6)	67.8 (2.3)	74.6 (3.0)	12	<3	
	31-50	2596	48.4 (0.9)	30.5 (1.3)	33.8 (1.2)	39.7 (1.1)	47.8 (1.1)	56.8 (1.4)	66.3 (2.1)	72.6 (2.7)	12	0.0 (0.0)	
	51-70	2550	43.9 (0.7)	27.6 (1.2)	30.6 (1.1)	36.0 (0.9)	42.8 (0.8)	50.7 (1.0)	58.9 (1.5)	64.6 (2.0)	12	0.0 (0.0)	
	>70	1520	36.4 (0.8)	22.6 (1.0)	25.0 (1.0)	29.5 (1.0)	35.4 (1.1)	42.3 (1.3)	49.5 (1.7)	54.4 (2.0)	12	<3	
	19+	8470	46.4 (0.5)	28.3 (0.6)	31.4 (0.6)	37.4 (0.6)	45.5 (0.6)	54.5 (0.8)	64.4 (1.2)	71.1 (1.5)	12	0.0 (0.0)	
Female													
	9-13	1980	32.0 (0.6)	20.9 (0.8)	23.0 (0.7)	26.7 (0.6)	31.3 (0.6)	36.5 (0.9)	42.1 (1.3)	45.9 (1.6)	9	0.0 (0.0)	
	14-18	2256	33.2 (0.5)	20.5 (0.7)	23.0 (0.7)	27.4 (0.6)	32.8 (0.6)	38.8 (0.8)	44.9 (1.0)	48.9 (1.3)	11	<3	
	19-30	1854	33.1 (0.7)	22.0 (1.1)	24.1 (1.0)	27.9 (0.9)	32.4 (0.8)	37.5 (1.0)	42.5 (1.4)	45.8 (1.8)	11	<3	
	31-50	2686	36.0 (0.8)	21.4 (0.9)	24.0 (0.8)	28.6 (0.8)	35.3 (1.0)	42.9 (1.2)	51.1 (1.6)	56.9 (2.1)	11	<3	
	51-70	3200	33.8 (0.6)	22.7 (0.9)	24.7 (0.9)	28.6 (0.8)	33.5 (0.7)	38.8 (0.8)	44.3 (1.2)	48.0 (1.6)	11	<3	
	>70	2610	29.8 (0.6)	18.5 (0.7)	20.5 (0.7)	24.2 (0.7)	29.0 (0.7)	34.6 (0.8)	40.6 (1.1)	44.8 (1.4)	11	<3	
	19+	10350	34.1 (0.4)	21.3 (0.4)	23.6 (0.4)	27.9 (0.4)	33.4 (0.5)	39.8 (0.6)	46.6 (0.8)	51.3 (1.0)	11	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

20. Phosphorus (mg/d): Usual intakes from food

Table 20.1 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	79	1077	(68)	778 (122)	849 (109)	963 (85)	1081 (70)	1206 (97)	1339 (148)	1431 (184)	380	<3		3000	<3
	4-8	127	1300	(71)	961 (131)	1035 (119)	1164 (103)	1318 (96)	1490 (110)	1669 (154)	1793 (196)	405	<3		3000	<3
Male																
	9-13	111	1601	(115)	1106 (154)	1202 (139)	1361 (118)	1547 (114)	1771 (155)	2025 (243)	2206 (319)	1055	F		4000	<3
	14-18	107	1555	(91)	1024 (164)	1131 (147)	1332 (119)	1594 (113)	1902 (160)	2225 (245)	2442 (313)	1055	F		4000	<3
	19-30	77	1510	(89)	979 (90)	1078 (100)	1281 (117)	1548 (125)	1805 (122)	2022 (125)	2168 (143)	580	<3		4000	<3
	31-50	145	1538	(181)	905 (184) ^E	999 (171) ^E	1176 (152)	1408 (147)	1698 (182)	2028 (265)	2259 (338)	580	<3		4000	<3
	51-70	182	1260	(72)	740 (71)	830 (72)	1002 (75)	1214 (83)	1437 (96)	1654 (122)	1799 (151)	580	<3		4000	<3
	>70	63	1542	(154)	1130 (208) ^E	1222 (207) ^E	1395 (211)	1622 (232)	1893 (287)	2182 (380) ^E	2378 (459) ^E	580	<3		3000	F
	19+	467	1449	(80)	966 (104)	1052 (96)	1213 (83)	1415 (76)	1644 (87)	1878 (123)	2034 (156)	580	<3			
Female																
	9-13	96	1223	(67)	791 (109)	873 (104)	1026 (95)	1221 (92)	1445 (105)	1674 (136)	1824 (164)	1055	F		4000	0.0 (0.0)
	14-18	105	1192	(116)	629 (137) ^E	714 (135) ^E	877 (132)	1106 (136)	1418 (169)	1818 (256)	2132 (340)	1055	44.6 (14.6) ^E		4000	<3
	19-30	91	1030	(58)	712 (97)	765 (90)	861 (79)	983 (76)	1125 (92)	1276 (127)	1379 (158)	580	F		4000	0.0 (0.0)
	31-50	167	1164	(85)	822 (164) ^E	889 (145)	1009 (121)	1157 (112)	1324 (128)	1487 (157)	1587 (180)	580	F		4000	<3
	51-70	198	1093	(61)	628 (128) ^E	721 (108)	880 (79)	1063 (69)	1289 (100)	1550 (173)	1728 (231)	580	F		4000	0.0 (0.0)
	>70	74	1125	(101)	833 (157) ^E	881 (145)	965 (126)	1065 (111)	1172 (110)	1274 (128)	1338 (149)	580	F		3000	0.0 (0.0)
	19+	530	1115	(38)	712 (65)	793 (60)	934 (54)	1105 (52)	1296 (60)	1488 (76)	1611 (88)	580	F			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.2 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	% UL ³	>UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	58	1190	(110)	896 (129)	962 (121)	1076 (115)	1211 (121)	1355 (144)	1491 (178)	1575 (203)	380	<3		3000	<3
	4-8	110	1323	(102)	965 (126)	1035 (119)	1161 (114)	1317 (122)	1484 (142)	1635 (166)	1725 (182)	405	<3		3000	0.0 (0.0)
Male																
	9-13	95	1467	(87)	1075 (164)	1155 (153)	1296 (138)	1462 (129)	1636 (135)	1801 (156)	1903 (174)	1055	F		4000	0.0 (0.0)
	14-18	87	1716	(104)	918 (171) ^E	1093 (157)	1402 (134)	1772 (133)	2183 (177)	2606 (250)	2888 (313)	1055	F		4000	<3
	19-30	70	2079	(197)	1064 (226) ^E	1266 (222) ^E	1650 (222)	2144 (254)	2709 (314)	3278 (396)	3647 (473)	580	<3		4000	F
	31-50	109	1463	(99)	825 (137) ^E	939 (125)	1149 (109)	1416 (116)	1741 (164)	2105 (240)	2367 (304)	580	<3		4000	<3
	51-70	128	1477	(68)	884 (102)	986 (95)	1173 (87)	1410 (86)	1695 (101)	2014 (144)	2242 (189)	580	<3		4000	<3
	>70	65	1097	(65)	716 (97)	782 (91)	906 (81)	1058 (83)	1223 (109)	1391 (156)	1506 (194)	580	<3		3000	<3
	19+	372	1559	(63)	842 (68)	970 (65)	1206 (62)	1506 (69)	1890 (102)	2358 (160)	2696 (207)	580	<3			
Female																
	9-13	75	1307	(125)	905 (188) ^E	992 (179) ^E	1148 (171)	1338 (180)	1545 (215)	1747 (271)	1875 (313) ^E	1055	F		4000	<3
	14-18	81	1234	(96)	644 (118) ^E	752 (118)	948 (115)	1189 (123)	1453 (154)	1709 (190)	1871 (215)	1055	F		4000	0.0 (0.0)
	19-30	101	1301	(106)	665 (110)	794 (106)	1036 (110)	1343 (136)	1691 (178)	2038 (232)	2262 (277)	580	F		4000	<3
	31-50	116	1106	(76)	594 (86)	690 (81)	866 (82)	1081 (97)	1320 (126)	1553 (161)	1701 (186)	580	F		4000	0.0 (0.0)
	51-70	146	1130	(47)	642 (55)	723 (55)	884 (55)	1102 (59)	1356 (73)	1608 (89)	1767 (101)	580	F		4000	0.0 (0.0)
	>70	94	951	(61)	577 (84)	635 (77)	746 (68)	899 (70)	1088 (100)	1296 (156)	1442 (204)	580	F		3000	<3
19+	457	1130	(41)	628 (40)	719 (40)	886 (43)	1100 (52)	1354 (69)	1611 (86)	1772 (95)	580	F				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.3 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	112	1092	(52)	904 (133)	943 (115)	1012 (87)	1093 (67)	1179 (90)	1261 (147)	1313 (194)	380	<3		3000	<3
	4-8	177	1305	(66)	853 (123)	943 (108)	1104 (85)	1302 (74)	1519 (100)	1732 (150)	1867 (188)	405	<3		3000	<3
Male																
	9-13	111	1540	(68)	935 (111)	1041 (102)	1236 (87)	1470 (80)	1734 (102)	2014 (157)	2206 (205)	1055	F		4000	<3
	14-18	113	1672	(177)	1008 (218) ^E	1126 (219) ^E	1350 (222)	1642 (234)	1953 (283)	2251 (377) ^E	2438 (468) ^E	1055	F		4000	<3
	19-30	91	1794	(143)	840 (239) ^E	986 (219) ^E	1272 (183)	1693 (152)	2145 (209)	2587 (309)	2906 (400)	580	F		4000	<3
	31-50	101	1502	(75)	1002 (84)	1104 (85)	1282 (90)	1489 (92)	1706 (94)	1918 (102)	2056 (111)	580	<3		4000	0.0 (0.0)
	51-70	134	1537	(63)	1037 (150)	1134 (131)	1319 (97)	1544 (78)	1774 (102)	1991 (152)	2134 (194)	580	<3		4000	<3
	>70	56	1271	(103)	814 (107)	896 (113)	1047 (122)	1241 (129)	1465 (140)	1695 (173)	1848 (204)	580	<3		3000	<3
	19+	382	1549	(49)	969 (77)	1076 (69)	1269 (59)	1513 (55)	1789 (68)	2063 (99)	2244 (127)	580	<3			
Female																
	9-13	105	1199	(97)	801 (129)	863 (122)	975 (111)	1133 (108)	1352 (128)	1591 (181)	1741 (227)	1055	F		4000	<3
	14-18	120	1157	(103)	536 (158) ^E	656 (143) ^E	874 (121)	1135 (108)	1418 (125)	1712 (171)	1917 (210)	1055	42.0 (11.9) ^E		4000	<3
	19-30	91	1249	(71)	903 (140)	980 (124)	1116 (100)	1276 (89)	1445 (107)	1605 (148)	1704 (182)	580	<3		4000	<3
	31-50	159	1252	(63)	653 (94)	777 (92)	1009 (83)	1282 (80)	1554 (92)	1802 (111)	1957 (126)	580	F		4000	0.0 (0.0)
	51-70	174	1155	(58)	674 (105)	768 (93)	934 (76)	1132 (67)	1348 (85)	1563 (118)	1702 (145)	580	F		4000	0.0 (0.0)
	>70	80	1101	(87)	582 (93)	669 (88)	823 (87)	1035 (106)	1344 (157)	1746 (236)	2058 (314)	580	F		3000	<3
	19+	504	1203	(35)	665 (51)	779 (52)	973 (49)	1202 (45)	1470 (60)	1732 (82)	1890 (102)	580	F			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.4 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	99	1275	(86)	932 (113)	995 (101)	1108 (89)	1260 (95)	1453 (129)	1663 (184)	1802 (226)	380	<3		3000	<3
	4-8	140	1286	(53)	931 (95)	1003 (84)	1132 (68)	1287 (59)	1458 (72)	1627 (102)	1736 (128)	405	0.0 (0.0)		3000	<3
Male																
	9-13	92	1442	(95)	1048 (138)	1122 (129)	1252 (118)	1406 (119)	1571 (138)	1721 (167)	1812 (189)	1055	F		4000	0.0 (0.0)
	14-18	107	1839	(136)	1199 (179)	1318 (169)	1542 (157)	1838 (164)	2190 (210)	2564 (293)	2819 (362)	1055	F		4000	<3
	19-30	73	1923	(166)	1092 (168)	1237 (169)	1529 (170)	1950 (196)	2504 (293)	3153 (459)	3627 (606) ^E	580	<3		4000	F
	31-50	134	1539	(96)	1028 (158)	1112 (144)	1267 (125)	1471 (116)	1709 (141)	1953 (204)	2115 (259)	580	<3		4000	<3
	51-70	131	1419	(86)	1037 (156)	1111 (139)	1243 (114)	1405 (103)	1586 (130)	1771 (196)	1892 (252)	580	<3		4000	<3
	>70	55	1227	(73)	750 (113)	830 (102)	977 (87)	1165 (84)	1383 (101)	1606 (139)	1752 (174)	580	F		3000	<3
	19+	393	1555	(56)	891 (70)	1003 (69)	1219 (70)	1510 (75)	1873 (93)	2276 (129)	2548 (163)	580	<3			
Female																
	9-13	79	1361	(107)	827 (140) ^E	925 (143)	1120 (148)	1382 (153)	1682 (167)	1972 (199)	2150 (230)	1055	F		4000	<3
	14-18	104	1237	(70)	777 (73)	906 (71)	1103 (74)	1267 (82)	1449 (110)	1699 (147)	1878 (171)	1055	F		4000	0.0 (0.0)
	19-30	101	1112	(83)	739 (138) ^E	818 (128)	967 (111)	1161 (102)	1390 (119)	1629 (163)	1790 (201)	580	F		4000	0.0 (0.0)
	31-50	143	1143	(71)	632 (109) ^E	723 (96)	877 (77)	1063 (71)	1281 (102)	1541 (207)	1741 (283)	580	F		4000	<3
	51-70	193	1060	(44)	642 (101)	713 (91)	844 (73)	1010 (61)	1196 (75)	1385 (110)	1507 (137)	580	F		4000	0.0 (0.0)
	>70	94	1062	(71)	680 (129) ^E	750 (121)	877 (107)	1037 (99)	1216 (111)	1397 (144)	1513 (172)	580	F		3000	<3
	19+	531	1102	(33)	648 (49)	732 (45)	887 (39)	1079 (37)	1294 (48)	1518 (73)	1674 (97)	580	F			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.5 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)								
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)							95th (SE)							
Both																								
	1-3	311	1181	(44)	727	(77)	831	(73)	1004	(66)	1194	(61)	1413	(67)	1633	(81)	1769	(92)	380	<3		3000	0.0	(0.0)
	4-8	485	1300	(45)	837	(89)	922	(76)	1074	(58)	1267	(52)	1493	(81)	1724	(131)	1872	(166)	405	<3		3000	<3	
Male																								
	9-13	277	1681	(78)	1056	(119)	1175	(110)	1396	(99)	1692	(102)	2047	(142)	2413	(207)	2653	(257)	1055	F		4000	<3	
	14-18	339	1846	(87)	1187	(136)	1321	(125)	1564	(109)	1872	(109)	2253	(136)	2671	(190)	2952	(237)	1055	F		4000	<3	
	19-30	237	1685	(91)	1171	(155)	1269	(142)	1448	(122)	1673	(115)	1928	(142)	2185	(199)	2353	(247)	580	<3		4000	<3	
	31-50	423	1619	(66)	1005	(116)	1121	(105)	1334	(87)	1604	(78)	1915	(99)	2236	(146)	2450	(186)	580	<3		4000	<3	
	51-70	387	1397	(48)	869	(79)	969	(73)	1156	(63)	1391	(59)	1654	(70)	1922	(101)	2102	(130)	580	<3		4000	<3	
	>70	132	1241	(101)	598	(126) ^E	704	(126) ^E	911	(126)	1190	(136)	1528	(160)	1888	(194)	2131	(219)	580	F		3000	<3	
	19+	1179	1538	(38)	900	(53)	1017	(51)	1236	(48)	1515	(47)	1837	(55)	2176	(75)	2409	(93)	580	<3				
Female																								
	9-13	281	1284	(59)	744	(98)	838	(90)	1011	(77)	1228	(69)	1490	(84)	1772	(131)	1952	(170)	1055	29.9	(8.8) ^E	4000	0.0	(0.0)
	14-18	321	1205	(38)	843	(67)	917	(62)	1054	(55)	1229	(51)	1427	(63)	1623	(90)	1751	(114)	1055	25.2	(7.2) ^E	4000	0.0	(0.0)
	19-30	249	1313	(74)	899	(111)	983	(97)	1132	(79)	1312	(80)	1512	(113)	1712	(165)	1844	(204)	580	<3		4000	0.0	(0.0)
	31-50	364	1340	(72)	891	(125)	971	(117)	1116	(107)	1317	(99)	1583	(114)	1888	(179)	2109	(247)	580	<3		4000	<3	
	51-70	467	1243	(44)	805	(89)	893	(77)	1050	(59)	1240	(50)	1450	(69)	1662	(106)	1802	(135)	580	<3		4000	0.0	(0.0)
	>70	215	1018	(51)	685	(67)	753	(64)	867	(63)	1012	(66)	1199	(79)	1396	(105)	1523	(126)	580	F		3000	<3	
	19+	1295	1266	(34)	792	(41)	879	(39)	1039	(38)	1251	(43)	1516	(55)	1807	(81)	2008	(107)	580	<3				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.6 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																
	1-3	644	1118 (27)	659 (55)	758 (46)	927 (35)	1124 (30)	1342 (39)	1568 (62)	1721 (81)	380	<3	3000	<3		
	4-8	956	1229 (22)	780 (47)	865 (41)	1020 (32)	1214 (26)	1435 (35)	1662 (56)	1812 (74)	405	<3	3000	0.0	(0.0)	
Male																
	9-13	589	1497 (43)	888 (73)	993 (66)	1191 (55)	1450 (49)	1758 (67)	2081 (108)	2298 (142)	1055	14.0 (4.3) ^E	4000	<3		
	14-18	639	1700 (45)	1085 (107)	1205 (94)	1424 (73)	1700 (56)	2018 (71)	2347 (119)	2568 (158)	1055	F	4000	<3		
	19-30	481	1580 (53)	917 (119)	1021 (106)	1218 (88)	1506 (67)	1864 (97)	2226 (165)	2465 (221)	580	<3	4000	<3		
	31-50	709	1465 (40)	935 (133)	1032 (115)	1214 (82)	1444 (49)	1709 (72)	1979 (138)	2156 (186)	580	<3	4000	0.0	(0.0)	
	51-70	758	1364 (34)	807 (72)	899 (65)	1074 (52)	1303 (40)	1575 (49)	1861 (86)	2054 (118)	580	<3	4000	<3		
	>70	734	1157 (28)	641 (39)	725 (37)	892 (34)	1110 (32)	1363 (41)	1622 (57)	1786 (71)	580	F	3000	<3		
	19+	2682	1436 (23)	803 (39)	909 (37)	1115 (31)	1381 (28)	1703 (36)	2041 (54)	2269 (71)	580	<3				
Female																
	9-13	585	1220 (32)	734 (61)	824 (53)	986 (40)	1185 (34)	1402 (49)	1613 (75)	1747 (95)	1055	33.2 (5.0)	4000	0.0	(0.0)	
	14-18	645	1261 (40)	665 (59)	770 (54)	971 (45)	1234 (42)	1545 (61)	1868 (98)	2083 (129)	1055	32.8 (4.3)	4000	<3		
	19-30	514	1099 (34)	630 (69)	715 (61)	868 (47)	1059 (38)	1275 (49)	1496 (78)	1640 (100)	580	F	4000	0.0	(0.0)	
	31-50	758	1168 (29)	654 (45)	747 (41)	919 (34)	1137 (31)	1396 (46)	1671 (71)	1850 (87)	580	F	4000	0.0	(0.0)	
	51-70	955	1133 (28)	606 (51)	703 (44)	877 (35)	1089 (31)	1336 (39)	1604 (63)	1792 (86)	580	F	4000	0.0	(0.0)	
	>70	1345	1049 (22)	577 (30)	658 (28)	812 (26)	1010 (25)	1239 (32)	1474 (46)	1629 (58)	580	5.2 (1.4) ^E	3000	0.0	(0.0)	
	19+	3572	1131 (16)	617 (20)	710 (19)	879 (17)	1094 (17)	1345 (23)	1612 (33)	1789 (41)	580	3.6 (0.7) ^E				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.7 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																
	1-3	324	1062 (43)	629 (80)	709 (70)	860 (54)	1053 (52)	1276 (80)	1504 (122)	1653 (152)	380	<3	3000	0.0	(0.0)	
	4-8	425	1194 (44)	877 (93)	938 (80)	1046 (59)	1176 (49)	1317 (69)	1453 (106)	1538 (134)	405	<3	3000	0.0	(0.0)	
Male																
	9-13	274	1494 (41)	933 (108)	1041 (93)	1231 (69)	1465 (52)	1731 (72)	1996 (118)	2163 (152)	1055	F	4000	0.0	(0.0)	
	14-18	297	1900 (97)	868 (112)	1040 (116)	1378 (119)	1837 (126)	2394 (180)	2989 (253)	3390 (287)	1055	F	4000	F		
	19-30	249	1831 (161)	1055 (225) ^E	1183 (199) ^E	1430 (156)	1726 (130)	2069 (228)	2422 (330)	2662 (414)	580	<3	4000	F		
	31-50	309	1418 (58)	860 (59)	961 (63)	1147 (66)	1378 (69)	1635 (74)	1891 (80)	2055 (85)	580	<3	4000	0.0	(0.0)	
	51-70	277	1308 (55)	693 (86)	793 (77)	974 (68)	1222 (66)	1556 (84)	1931 (129)	2177 (166)	580	F	4000	<3		
	>70	136	1214 (67)	600 (67)	695 (68)	878 (68)	1131 (77)	1450 (99)	1775 (119)	1970 (130)	580	F	3000	<3		
	19+	971	1462 (47)	791 (59)	894 (55)	1091 (49)	1368 (46)	1712 (65)	2079 (103)	2332 (139)	580	<3				
Female																
	9-13	265	1283 (55)	707 (84)	805 (75)	989 (66)	1221 (65)	1489 (78)	1787 (116)	2000 (157)	1055	31.8 (7.7) ^E	4000	<3		
	14-18	290	1189 (54)	738 (107)	822 (92)	970 (71)	1151 (65)	1356 (94)	1568 (145)	1712 (185)	1055	36.4 (10.3) ^E	4000	0.0	(0.0)	
	19-30	197	1289 (77)	777 (157) ^E	876 (137)	1057 (106)	1284 (90)	1541 (124)	1801 (194)	1972 (248)	580	F	4000	<3		
	31-50	312	1192 (51)	668 (60)	755 (59)	920 (57)	1139 (59)	1392 (73)	1654 (99)	1833 (123)	580	F	4000	0.0	(0.0)	
	51-70	312	1208 (79)	710 (97)	791 (89)	945 (75)	1145 (69)	1382 (95)	1638 (150)	1817 (199)	580	F	4000	<3		
	>70	239	1003 (51)	531 (75)	606 (70)	756 (59)	963 (52)	1223 (74)	1515 (124)	1721 (166)	580	F	3000	<3		
	19+	1060	1188 (33)	631 (38)	725 (36)	904 (33)	1134 (34)	1415 (46)	1725 (73)	1943 (98)	580	F				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.8 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	% >UL ³	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)			
Both															
	1-3	129	1134	(69)	638	733	912	1137	1393	1649	1815	380	<3	3000	<3
	4-8	213	1206	(41)	776	859	1003	1176	1375	1589	1739	405	<3	3000	<3
Male															
	9-13	122	1509	(88)	1165	1235	1361	1514	1685	1855	1964	1055	F	4000	0.0 (0.0)
	14-18	150	1877	(94)	1158	1295	1546	1860	2212	2562	2788	1055	F	4000	<3
	19-30	106	1688	(96)	864	1017	1299	1653	2049	2440	2689	580	F	4000	<3
	31-50	155	1584	(91)	856	1007	1279	1603	1946	2270	2470	580	F	4000	<3
	51-70	122	1396	(80)	840	960	1177	1441	1730	2010	2186	580	<3	4000	<3
	>70	88	1450	(93)	782	885	1092	1382	1754	2179	2484	580	<3	3000	F
	19+	471	1542	(48)	781	928	1194	1534	1922	2313	2574	580	<3		
Female															
	9-13	103	1311	(90)	871	950	1097	1283	1503	1742	1912	1055	F	4000	<3
	14-18	142	1150	(71)	633	730	899	1115	1380	1661	1843	1055	43.0 (9.9) ^E	4000	0.0 (0.0)
	19-30	111	1161	(79)	671	756	915	1117	1350	1586	1741	580	F	4000	0.0 (0.0)
	31-50	146	1232	(73)	647	739	904	1107	1335	1565	1715	580	F	4000	0.0 (0.0)
	51-70	184	1183	(65)	772	840	967	1141	1347	1557	1693	580	<3	4000	0.0 (0.0)
	>70	143	1111	(50)	669	756	915	1118	1363	1640	1838	580	F	3000	<3
	19+	584	1185	(37)	687	770	930	1132	1357	1590	1751	580	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.9 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	% UL ³	>UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	169	1054	(47)	580	665	826	1035	1277	1527	1691	380	<3		3000	<3
	4-8	281	1229	(39)	848	919	1046	1205	1381	1557	1671	405	<3		3000	<3
Male																
	9-13	183	1597	(99)	1129	1221	1391	1610	1871	2156	2355	1055	F		4000	<3
	14-18	187	1788	(80)	1011	1159	1435	1784	2178	2572	2827	1055	F		4000	<3
	19-30	223	1605	(65)	985	1103	1321	1587	1866	2119	2272	580	<3		4000	<3
	31-50	229	1526	(84)	990	1082	1260	1500	1797	2127	2360	580	<3		4000	<3
	51-70	197	1399	(69)	732	852	1073	1353	1680	2026	2260	580	F		4000	<3
	>70	72	1321	(69)	1013	1080	1199	1340	1493	1640	1732	580	<3		3000	<3
	19+	721	1500	(44)	932	1040	1235	1483	1776	2084	2288	580	<3			
Female																
	9-13	165	1222	(57)	913	984	1107	1255	1419	1585	1695	1055	F		4000	0.0 (0.0)
	14-18	206	1157	(46)	714	792	929	1109	1333	1567	1715	1055	42.6 (9.5) ^E		4000	0.0 (0.0)
	19-30	191	1156	(73)	755	828	961	1128	1313	1496	1611	580	F		4000	0.0 (0.0)
	31-50	258	1205	(62)	882	951	1072	1220	1382	1543	1647	580	<3		4000	0.0 (0.0)
	51-70	249	1082	(52)	657	735	872	1040	1237	1451	1602	580	F		4000	<3
	>70	128	1107	(78)	666	753	923	1116	1350	1630	1804	580	F		3000	<3
	19+	826	1152	(33)	706	792	943	1131	1359	1606	1770	580	F			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.10 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	192	1093	(43)	734 (64)	801 (60)	925 (54)	1071 (54)	1221 (63)	1366 (80)	1461 (95)	380	<3		3000	0.0 (0.0)
	4-8	321	1257	(31)	832 (53)	916 (48)	1066 (41)	1250 (39)	1462 (49)	1680 (68)	1824 (84)	405	<3		3000	<3
Male																
	9-13	226	1541	(57)	880 (70)	1002 (67)	1229 (66)	1522 (69)	1883 (88)	2296 (129)	2596 (165)	1055	12.9 (3.8) ^E		4000	<3
	14-18	262	1974	(127)	1215 (203) ^E	1360 (183)	1628 (150)	1973 (134)	2382 (178)	2820 (278)	3120 (360)	1055	^F		4000	<3
	19-30	197	1776	(73)	1122 (133)	1238 (119)	1454 (97)	1733 (88)	2058 (121)	2395 (187)	2618 (239)	580	<3		4000	<3
	31-50	282	1822	(89)	1111 (70)	1238 (77)	1488 (91)	1810 (103)	2155 (110)	2488 (121)	2707 (137)	580	<3		4000	<3
	51-70	234	1635	(72)	967 (86)	1081 (84)	1296 (82)	1574 (85)	1896 (106)	2237 (150)	2471 (190)	580	<3		4000	<3
	>70	119	1362	(97)	745 (118)	856 (116)	1063 (116)	1325 (123)	1620 (152)	1917 (182)	2109 (202)	580	^F		3000	<3
	19+	832	1714	(45)	976 (46)	1103 (45)	1347 (45)	1672 (51)	2049 (68)	2437 (92)	2697 (112)	580	<3			
Female																
	9-13	226	1315	(73)	772 (79)	870 (78)	1051 (78)	1284 (87)	1562 (108)	1868 (145)	2083 (177)	1055	25.4 (7.9) ^E		4000	<3
	14-18	242	1209	(55)	553 (73)	659 (71)	866 (69)	1142 (74)	1471 (86)	1815 (106)	2045 (128)	1055	42.1 (7.0)		4000	<3
	19-30	208	1279	(69)	919 (136)	998 (121)	1139 (99)	1307 (90)	1486 (113)	1657 (158)	1765 (193)	580	<3		4000	0.0 (0.0)
	31-50	263	1265	(56)	743 (69)	833 (62)	995 (58)	1197 (64)	1443 (88)	1723 (135)	1928 (178)	580	<3		4000	<3
	51-70	322	1148	(64)	660 (103)	751 (92)	915 (75)	1122 (71)	1376 (102)	1658 (164)	1850 (211)	580	^F		4000	0.0 (0.0)
	>70	198	1092	(57)	665 (94)	747 (89)	893 (81)	1065 (80)	1253 (98)	1440 (134)	1561 (158)	580	^F		3000	<3
	19+	991	1212	(30)	727 (41)	815 (38)	973 (36)	1176 (38)	1423 (52)	1691 (80)	1875 (104)	580	<3			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.11 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		UL ³	% >UL		
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)	(<i>SE</i>)		(<i>SE</i>)		
Both																	
	1-3	348	1155	(38)	812	883	1010	1159	1318	1483	1599	380	0.0	(0.0)	3000	0.0	(0.0)
	4-8	554	1299	(36)	912	991	1138	1319	1513	1704	1829	405	0.0	(0.0)	3000	0.0	(0.0)
Male																	
	9-13	409	1517	(48)	1007	1102	1272	1481	1706	1925	2072	1055	F		4000	0.0	(0.0)
	14-18	414	1700	(84)	984	1109	1360	1687	2056	2443	2710	1055	F		4000	<3	
	19-30	311	1788	(79)	967	1103	1378	1756	2197	2681	3031	580	<3		4000	<3	
	31-50	489	1520	(58)	936	1036	1224	1469	1752	2041	2235	580	<3		4000	<3	
	51-70	575	1432	(41)	930	1022	1193	1419	1669	1898	2047	580	<3		4000	0.0	(0.0)
	>70	239	1303	(58)	839	922	1078	1281	1518	1767	1934	580	<3		3000	<3	
	19+	1614	1529	(32)	901	1007	1211	1486	1807	2151	2391	580	<3				
Female																	
	9-13	355	1263	(55)	809	887	1040	1248	1492	1737	1892	1055	26.7	(8.5) ^E	4000	0.0	(0.0)
	14-18	410	1195	(53)	629	755	961	1190	1431	1696	1906	1055	34.6	(6.5) ^E	4000	0.0	(0.0)
	19-30	384	1161	(43)	763	843	989	1175	1387	1601	1738	580	<3		4000	0.0	(0.0)
	31-50	585	1189	(39)	623	742	957	1205	1464	1723	1904	580	F		4000	<3	
	51-70	711	1109	(31)	614	704	877	1079	1302	1542	1702	580	F		4000	0.0	(0.0)
	>70	342	1084	(48)	636	713	856	1046	1285	1551	1733	580	F		3000	<3	
	19+	2022	1147	(20)	663	762	934	1138	1377	1626	1801	580	2.4	(0.7) ^E			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.12 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		UL ³	% >UL		
					(SE)	(SE)	(SE)	(SE)	(SE)	(SE)		(SE)	(SE)		(SE)	(SE)	
Both																	
	1-3	622	1070	(32)	601	686	848	1056	1296	1542	1704	380	<3	3000	<3		
	4-8	919	1217	(27)	786	866	1011	1189	1392	1602	1742	405	<3	3000	<3		
Male																	
	9-13	579	1559	(61)	1088	1177	1345	1565	1821	2086	2262	1055	F	4000	<3		
	14-18	634	1828	(55)	1024	1179	1459	1808	2210	2631	2917	1055	F	4000	<3		
	19-30	578	1663	(54)	931	1056	1291	1601	1954	2300	2528	580	<3	4000	<3		
	31-50	693	1514	(57)	939	1048	1242	1489	1784	2097	2306	580	<3	4000	<3		
	51-70	596	1378	(45)	722	844	1066	1347	1674	2022	2258	580	F	4000	<3		
	>70	296	1325	(44)	762	865	1051	1283	1559	1863	2083	580	F	3000	<3		
	19+	2163	1500	(30)	836	955	1172	1458	1804	2171	2418	580	<3				
Female																	
	9-13	533	1251	(41)	820	901	1053	1246	1468	1703	1868	1055	25.3	(6.6) ^E	4000	0.0	(0.0)
	14-18	638	1163	(33)	688	775	926	1118	1357	1611	1775	1055	41.8	(4.8)	4000	0.0	(0.0)
	19-30	499	1185	(48)	701	788	949	1152	1382	1613	1764	580	F	4000	0.0	(0.0)	
	31-50	716	1207	(42)	721	808	970	1176	1411	1649	1804	580	F	4000	0.0	(0.0)	
	51-70	745	1130	(36)	712	789	927	1098	1296	1506	1651	580	<3	4000	0.0	(0.0)	
	>70	510	1080	(43)	602	692	866	1067	1321	1647	1868	580	F	3000	<3		
	19+	2470	1166	(22)	659	749	918	1129	1385	1667	1842	580	2.4	(0.7) ^E			

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 20.13 Phosphorus (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)		% UL ³ >UL (SE)			
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)					
Both																
	1-3	2117	1123 (17)	685 (28)	776 (26)	933 (22)	1121 (21)	1335 (24)	1553 (32)	1697 (38)	380	<3		3000	0.0	(0.0)
	4-8	3235	1250 (16)	788 (21)	876 (19)	1034 (17)	1231 (19)	1460 (25)	1698 (36)	1855 (45)	405	<3		3000	0.0	(0.0)
Male																
	9-13	2080	1557 (28)	971 (35)	1075 (34)	1275 (31)	1521 (33)	1827 (41)	2153 (58)	2365 (73)	1055	8.9 (1.8) ^E		4000	<3	
	14-18	2288	1790 (34)	1058 (45)	1199 (43)	1458 (39)	1786 (40)	2174 (50)	2595 (72)	2889 (90)	1055	4.9 (1.2) ^E		4000	<3	
	19-30	1804	1659 (34)	970 (43)	1085 (41)	1306 (39)	1602 (42)	1948 (54)	2304 (77)	2541 (96)	580	<3		4000	<3	
	31-50	2596	1560 (27)	928 (37)	1041 (35)	1253 (32)	1526 (32)	1849 (41)	2189 (59)	2416 (75)	580	<3		4000	<3	
	51-70	2550	1417 (22)	817 (32)	921 (31)	1118 (28)	1373 (27)	1669 (31)	1974 (44)	2184 (57)	580	<3		4000	0.0	(0.0)
	>70	1520	1244 (31)	686 (33)	782 (33)	965 (35)	1202 (39)	1478 (45)	1764 (56)	1952 (67)	580	^F		3000	<3	
	19+	8470	1515 (15)	861 (18)	974 (17)	1187 (17)	1467 (19)	1799 (24)	2155 (32)	2395 (40)	580	0.4 (0.1) ^E				
Female																
	9-13	1980	1255 (23)	755 (26)	846 (25)	1010 (24)	1220 (25)	1465 (33)	1714 (45)	1875 (55)	1055	30.2 (2.8)		4000	0.0	(0.0)
	14-18	2256	1219 (20)	678 (25)	775 (24)	956 (23)	1197 (25)	1480 (32)	1771 (43)	1966 (55)	1055	35.1 (2.5)		4000	0.0	(0.0)
	19-30	1854	1192 (25)	737 (34)	822 (32)	974 (29)	1163 (28)	1379 (35)	1598 (48)	1741 (60)	580	<3		4000	0.0	(0.0)
	31-50	2686	1229 (23)	689 (27)	787 (27)	964 (25)	1184 (27)	1469 (36)	1792 (58)	2013 (81)	580	1.8 (0.5) ^E		4000	<3	
	51-70	3200	1161 (18)	685 (25)	773 (23)	932 (20)	1131 (20)	1365 (26)	1618 (39)	1793 (51)	580	1.8 (0.5) ^E		4000	0.0	(0.0)
	>70	2610	1055 (19)	621 (24)	702 (24)	848 (23)	1032 (25)	1253 (29)	1494 (38)	1655 (44)	580	3.3 (0.9) ^E		3000	0.0	(0.0)
	19+	10350	1181 (12)	683 (13)	774 (13)	940 (13)	1147 (14)	1401 (18)	1679 (27)	1869 (37)	580	1.9 (0.3)				

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

21. Potassium (mg/d): Usual intakes from food

Table 21.1 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	2317 (115)	1695 (221)	1841 (187)	2069 (140)	2291 (139)	2532 (186)	2806 (251)	2992 (295)	3000	F	
	4-8	127	2570 (99)	1769 (167)	1916 (151)	2178 (127)	2507 (118)	2912 (147)	3379 (224)	3723 (300)	3800	F	
Male	9-13	111	3567 (256)	2370 (201)	2576 (216)	2965 (248)	3477 (298)	4090 (375)	4744 (468)	5190 (535)	4500	F	
	14-18	107	3246 (226)	1846 (263)	2067 (264)	2534 (266)	3233 (291)	4106 (419)	5125 (676)	5906 (926)	4700	F	
	19-30	77	3218 (216)	2090 (197)	2341 (216)	2779 (252)	3289 (287)	3823 (308)	4321 (322)	4626 (335)	4700	F	
	31-50	145	3386 (283)	2523 (366)	2697 (335)	2993 (295)	3340 (277)	3722 (302)	4110 (383)	4370 (460)	4700	F	
	51-70	182	3198 (162)	1895 (313)	2108 (281)	2507 (225)	3017 (184)	3607 (229)	4213 (358)	4613 (465)	4700	F	
	>70	63	3563 (271)	2498 (363)	2697 (354)	3079 (352)	3603 (386)	4251 (494)	4962 (686)	5466 (847)	4700	F	
	19+	467	3310 (136)	2251 (201)	2455 (180)	2814 (153)	3258 (143)	3772 (170)	4296 (236)	4637 (292)	4700	F	
	Female	9-13	96	2541 (125)	1816 (279)	1979 (243)	2265 (195)	2602 (179)	2959 (223)	3297 (299)	3507 (356)	4500	<3
14-18	105	2653 (195)	1516 (290) ^E	1716 (280)	2102 (267)	2623 (268)	3282 (315)	4040 (438)	4588 (560)	4700	F		
19-30	91	2470 (192)	1683 (246)	1815 (229)	2054 (209)	2350 (217)	2681 (278)	3011 (385)	3224 (472)	4700	<3		
31-50	167	2407 (132)	1531 (237)	1722 (219)	2058 (190)	2463 (182)	2913 (214)	3369 (287)	3671 (345)	4700	<3		
51-70	198	2671 (126)	1509 (190)	1735 (165)	2130 (135)	2605 (138)	3149 (192)	3738 (282)	4149 (362)	4700	F		
>70	74	2613 (191)	1583 (268) ^E	1761 (256)	2083 (237)	2482 (223)	2924 (230)	3360 (274)	3639 (324)	4700	<3		
19+	530	2520 (73)	1462 (105)	1672 (100)	2046 (93)	2504 (98)	3026 (117)	3570 (152)	3938 (182)	4700	<3		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.2 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	58	2376 (190)	1673 (237)	1822 (221)	2086 (205)	2402 (215)	2741 (259)	3067 (326)	3272 (377)	3000	F	
	4-8	110	2473 (157)	1804 (278)	1952 (259)	2210 (233)	2508 (221)	2813 (234)	3085 (262)	3244 (283)	3800	<3	
Male													
	9-13	95	2988 (162)	2354 (304)	2486 (280)	2712 (248)	2971 (230)	3241 (246)	3495 (292)	3654 (333)	4500	<3	
	14-18	87	3416 (189)	1743 (330) ^E	2074 (297)	2678 (243)	3433 (234)	4317 (348)	5275 (557)	5941 (728)	4700	F	
	19-30	70	4130 (295)	2585 (443) ^E	2911 (432)	3480 (412)	4316 (403)	5240 (465)	5962 (529)	6384 (589)	4700	F	
	31-50	109	3497 (208)	2174 (341)	2415 (301)	2864 (245)	3441 (242)	4110 (340)	4796 (500)	5248 (621)	4700	F	
	51-70	128	3342 (128)	2276 (258)	2501 (225)	2898 (178)	3367 (157)	3863 (188)	4332 (252)	4624 (309)	4700	F	
	>70	65	2769 (112)	2039 (195)	2189 (168)	2442 (134)	2736 (132)	3059 (180)	3392 (260)	3615 (324)	4700	<3	
	19+	372	3512 (109)	2253 (161)	2503 (145)	2950 (126)	3514 (132)	4182 (171)	4892 (243)	5370 (306)	4700	13.0 (4.0) ^E	
Female													
	9-13	75	2746 (208)	2423 (380)	2506 (357)	2651 (328)	2820 (319)	2998 (350)	3166 (420)	3271 (482)	4500	<3	
	14-18	81	2577 (141)	1577 (278) ^E	1770 (242)	2113 (190)	2500 (168)	2896 (201)	3258 (265)	3478 (313)	4700	<3	
	19-30	101	2878 (172)	1687 (276)	1935 (260)	2410 (235)	2992 (230)	3591 (274)	4137 (359)	4469 (424)	4700	F	
	31-50	116	2636 (149)	1494 (194)	1713 (186)	2107 (180)	2588 (193)	3115 (229)	3628 (281)	3952 (320)	4700	<3	
	51-70	146	2921 (110)	1672 (180)	1910 (169)	2350 (153)	2903 (143)	3527 (164)	4150 (230)	4552 (286)	4700	F	
	>70	94	2252 (133)	1276 (183)	1429 (171)	1730 (156)	2130 (164)	2586 (217)	3024 (292)	3289 (343)	4700	<3	
	19+	457	2715 (74)	1578 (87)	1802 (86)	2208 (90)	2701 (103)	3235 (123)	3749 (146)	4072 (164)	4700	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.3 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI	(SE)						
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)									
Both	1-3	112	2167	(99)	1389	(221)	1528	(196)	1780	(156)	2093	(131)	2440	(158)	2785	(230)	3006	(289)	3000	F
	4-8	177	2584	(129)	1666	(287) ^E	1845	(251)	2191	(183)	2613	(152)	3045	(207)	3468	(310)	3752	(395)	3800	F
Male	9-13	111	3091	(153)	1757	(258)	2000	(236)	2447	(203)	3007	(197)	3636	(254)	4261	(366)	4663	(474)	4500	F
	14-18	113	3210	(245)	2536	(426) ^E	2639	(395)	2846	(370)	3144	(385)	3486	(469)	3793	(626)	3973	(753) ^E	4700	F
	19-30	91	3669	(253)	2309	(522) ^E	2597	(447) ^E	3085	(337)	3636	(283)	4212	(345)	4775	(488)	5141	(601)	4700	F
	31-50	101	3692	(185)	2222	(349)	2499	(309)	2997	(256)	3599	(241)	4251	(295)	4880	(409)	5275	(506)	4700	F
	51-70	134	3440	(130)	2532	(285)	2722	(261)	3064	(219)	3463	(178)	3860	(184)	4221	(260)	4447	(339)	4700	F
	>70	56	2988	(218)	2252	(281)	2386	(271)	2634	(262)	2944	(264)	3290	(285)	3628	(347)	3842	(417)	4700	F
	19+	382	3544	(106)	2200	(175)	2444	(162)	2885	(148)	3455	(139)	4077	(152)	4643	(185)	4992	(217)	4700	F
	Female	9-13	105	2692	(218)	1739	(308) ^E	1917	(282)	2248	(235)	2659	(218)	3106	(271)	3528	(357)	3785	(418)	4500
14-18		120	2290	(170)	1150	(256) ^E	1346	(249) ^E	1738	(222)	2217	(210)	2750	(262)	3355	(372)	3786	(469)	4700	F
19-30		91	2723	(209)	1849	(266)	2022	(247)	2341	(232)	2728	(256)	3138	(328)	3542	(443)	3814	(539)	4700	F
31-50		159	2928	(124)	1739	(238)	2011	(225)	2484	(199)	3004	(186)	3536	(212)	4056	(263)	4386	(306)	4700	F
51-70		174	2695	(115)	1755	(244)	1939	(210)	2258	(161)	2631	(139)	3038	(179)	3446	(264)	3715	(335)	4700	<3
>70		80	2817	(173)	1734	(205)	1979	(189)	2375	(184)	2808	(201)	3329	(259)	3993	(368)	4515	(477)	4700	F
19+		504	2808	(72)	1682	(118)	1913	(114)	2336	(109)	2835	(107)	3348	(116)	3830	(142)	4137	(170)	4700	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.4 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	99	2666 (155)	1842 (238)	2007 (222)	2321 (201)	2731 (201)	3204 (249)	3679 (347)	3985 (432)	3000	F	
	4-8	140	2821 (215)	2009 (264)	2174 (239)	2490 (209)	2865 (232)	3247 (323)	3646 (487)	3913 (630)	3800	F	
Male													
	9-13	92	2729 (179)	1914 (202)	2047 (209)	2294 (220)	2609 (232)	2974 (251)	3353 (278)	3604 (300)	4500	<3	
	14-18	107	3636 (251)	2589 (333)	2814 (301)	3200 (267)	3663 (289)	4190 (383)	4738 (521)	5096 (619)	4700	F	
	19-30	73	3923 (286)	2102 (372) ^E	2490 (338)	3198 (312)	4024 (351)	4916 (463)	6025 (777)	6974 (1153)	4700	F	
	31-50	134	3594 (248)	2444 (364)	2629 (336)	2974 (296)	3420 (272)	3940 (304)	4487 (424)	4860 (554)	4700	F	
	51-70	131	3296 (235)	2028 (347) ^E	2251 (325)	2673 (291)	3215 (285)	3848 (345)	4524 (481)	4996 (610)	4700	F	
	>70	55	3089 (209)	1676 (324) ^E	1922 (304)	2382 (273)	2973 (248)	3654 (279)	4348 (380)	4802 (468)	4700	F	
	19+	393	3529 (133)	1913 (180)	2190 (179)	2733 (175)	3444 (173)	4268 (203)	5195 (283)	5828 (361)	4700	16.5 (4.1) ^E	
Female													
	9-13	79	2662 (168)	1844 (288)	2003 (276)	2314 (255)	2729 (243)	3175 (272)	3572 (346)	3812 (416)	4500	F	
	14-18	104	2596 (117)	1769 (143)	1965 (142)	2290 (142)	2654 (144)	3028 (156)	3376 (179)	3592 (200)	4700	<3	
	19-30	101	2497 (181)	1520 (287) ^E	1726 (275)	2116 (260)	2595 (276)	3097 (338)	3572 (454)	3878 (536)	4700	F	
	31-50	143	2803 (159)	1471 (282) ^E	1739 (261)	2213 (219)	2737 (190)	3262 (272)	3813 (367)	4203 (450)	4700	F	
	51-70	193	2607 (121)	1551 (221)	1722 (197)	2041 (159)	2451 (143)	2925 (194)	3414 (298)	3736 (382)	4700	<3	
	>70	94	2481 (150)	1398 (192)	1559 (189)	1867 (181)	2276 (181)	2768 (224)	3294 (295)	3651 (354)	4700	<3	
	19+	531	2645 (77)	1385 (119)	1618 (107)	2042 (92)	2596 (96)	3180 (110)	3722 (136)	4110 (172)	4700	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.5 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI	(SE)								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)				95th (SE)							
Both	1-3	311	2441	(88)	1455	(139)	1656	(133)	2003	(122)	2414	(117)	2872	(128)	3346	(158)	3663	(190)	3000	19.9	(4.8) ^E
	4-8	485	2740	(87)	1708	(185)	1917	(152)	2279	(108)	2713	(96)	3192	(151)	3659	(235)	3950	(293)	3800	F	
Male	9-13	277	3408	(165)	2238	(272)	2479	(252)	2915	(226)	3449	(224)	4038	(270)	4616	(351)	4983	(413)	4500	F	
	14-18	339	3805	(154)	2387	(165)	2699	(165)	3255	(173)	3939	(195)	4737	(238)	5602	(309)	6209	(383)	4700	25.9	(6.1) ^E
	19-30	237	3752	(182)	2549	(288)	2781	(270)	3219	(241)	3789	(233)	4462	(295)	5170	(426)	5648	(539)	4700	F	
	31-50	423	3687	(138)	2410	(251)	2668	(229)	3128	(193)	3688	(170)	4305	(191)	4916	(256)	5310	(312)	4700	F	
	51-70	387	3383	(110)	2070	(154)	2332	(144)	2808	(133)	3386	(141)	4043	(173)	4727	(227)	5167	(269)	4700	10.4	(3.4) ^E
	>70	132	2931	(192)	1612	(305) ^E	1865	(282)	2331	(251)	2917	(249)	3574	(282)	4227	(339)	4647	(396)	4700	F	
	19+	1179	3550	(80)	2160	(107)	2429	(108)	2927	(107)	3561	(104)	4284	(118)	5018	(150)	5496	(177)	4700	15.2	(2.6) ^E
	Female	9-13	281	2756	(124)	1568	(135)	1744	(135)	2112	(131)	2599	(131)	3105	(157)	3598	(219)	3939	(280)	4500	F
14-18	321	2746	(91)	1933	(171)	2126	(152)	2432	(127)	2783	(117)	3213	(136)	3650	(194)	3945	(247)	4700	<3		
19-30	249	2976	(133)	2065	(253)	2238	(230)	2551	(191)	2937	(168)	3366	(201)	3791	(287)	4064	(363)	4700	F		
31-50	364	3089	(116)	1915	(152)	2141	(141)	2549	(127)	3016	(141)	3648	(177)	4362	(253)	4830	(330)	4700	F		
51-70	467	3001	(93)	1906	(175)	2134	(151)	2530	(119)	2995	(109)	3501	(139)	4006	(198)	4337	(246)	4700	F		
>70	215	2523	(106)	1611	(177)	1786	(169)	2114	(156)	2521	(153)	2967	(175)	3411	(224)	3704	(271)	4700	<3		
19+	1295	2971	(63)	1877	(87)	2094	(80)	2481	(74)	2957	(78)	3514	(99)	4104	(136)	4495	(171)	4700	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.6 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI	(SE)	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)				95th (SE)
Both														
	1-3	644	2337	(51)	1401 (111)	1599 (95)	1940 (72)	2339 (59)	2785 (73)	3248 (115)	3559 (153)	3000	16.7	(3.1) ^E
	4-8	956	2591	(48)	1707 (111)	1881 (94)	2179 (70)	2551 (58)	2994 (82)	3445 (138)	3733 (180)	3800	F	
Male														
	9-13	589	3088	(91)	1920 (185)	2122 (169)	2521 (132)	3026 (103)	3573 (141)	4142 (242)	4545 (322)	4500	F	
	14-18	639	3491	(85)	2194 (209)	2452 (184)	2919 (140)	3499 (109)	4155 (138)	4826 (223)	5270 (292)	4700	12.0	(3.6) ^E
	19-30	481	3397	(123)	2025 (353) ^E	2256 (312)	2707 (227)	3305 (151)	3993 (238)	4691 (425)	5153 (571)	4700	F	
	31-50	709	3317	(92)	2020 (218)	2256 (195)	2692 (150)	3244 (109)	3874 (141)	4513 (240)	4930 (321)	4700	F	
	51-70	758	3409	(72)	2117 (144)	2348 (128)	2772 (99)	3306 (81)	3911 (112)	4519 (182)	4914 (238)	4700	F	
	>70	734	2982	(67)	1809 (140)	2029 (123)	2428 (93)	2921 (72)	3466 (100)	4003 (162)	4346 (213)	4700	F	
	19+	2682	3328	(51)	1988 (97)	2227 (89)	2682 (73)	3248 (61)	3894 (76)	4570 (120)	5016 (159)	4700	8.2	(1.6) ^E
Female														
	9-13	585	2646	(68)	1504 (130)	1714 (111)	2097 (83)	2572 (73)	3104 (110)	3638 (173)	3984 (219)	4500	F	
	14-18	645	2758	(76)	1429 (101)	1661 (94)	2102 (82)	2679 (83)	3354 (113)	4051 (164)	4512 (206)	4700	F	
	19-30	514	2511	(73)	1442 (141)	1633 (127)	1987 (100)	2424 (81)	2905 (97)	3382 (143)	3690 (180)	4700	<3	
	31-50	758	2779	(60)	1591 (98)	1805 (88)	2207 (76)	2718 (75)	3301 (91)	3911 (139)	4330 (189)	4700	F	
	51-70	955	2827	(67)	1612 (124)	1840 (111)	2247 (92)	2754 (80)	3341 (95)	3945 (142)	4338 (181)	4700	F	
	>70	1345	2651	(48)	1500 (74)	1711 (66)	2092 (56)	2558 (53)	3081 (67)	3615 (95)	3967 (118)	4700	<3	
	19+	3572	2725	(36)	1550 (50)	1764 (46)	2155 (41)	2651 (42)	3218 (51)	3799 (71)	4187 (90)	4700	1.9	(0.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.7 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both	1-3	324	2233	(63)	1170	(122)	1378	(113)	1747	(99)	2198	(82)	2717	(134)	3262	(163)	3630	(196)	3000	15.8	(3.6) ^E
	4-8	425	2334	(87)	1658	(186)	1781	(162)	2006	(124)	2288	(103)	2608	(140)	2932	(218)	3144	(279)	3800	<3	
Male	9-13	274	2841	(119)	1880	(257)	2072	(219)	2410	(159)	2805	(126)	3218	(173)	3611	(260)	3859	(323)	4500	<3	
	14-18	297	3697	(142)	1982	(225)	2309	(210)	2900	(183)	3649	(168)	4543	(207)	5500	(302)	6148	(398)	4700	21.7	(4.4) ^E
	19-30	249	3695	(322)	2093	(158)	2351	(173)	2817	(204)	3403	(250)	4110	(326)	4892	(437)	5429	(523)	4700	F	
	31-50	309	3190	(137)	2068	(269)	2288	(236)	2670	(188)	3139	(166)	3673	(194)	4201	(274)	4534	(352)	4700	F	
	51-70	277	3204	(108)	2030	(232)	2238	(209)	2635	(165)	3143	(130)	3690	(177)	4214	(303)	4554	(429)	4700	F	
	>70	136	3012	(190)	1936	(339) ^E	2133	(300)	2488	(225)	2933	(197)	3431	(329)	3917	(490)	4216	(577)	4700	F	
	19+	971	3291	(96)	2026	(177)	2254	(159)	2669	(125)	3180	(100)	3753	(125)	4346	(203)	4750	(277)	4700	F	
	Female	9-13	265	2435	(125)	1444	(206)	1641	(183)	1985	(153)	2376	(146)	2771	(175)	3141	(230)	3378	(278)	4500	<3
14-18	290	2464	(105)	1357	(162)	1561	(149)	1914	(133)	2332	(133)	2834	(159)	3416	(223)	3829	(286)	4700	<3		
19-30	197	2742	(158)	1922	(319) ^E	2092	(288)	2406	(239)	2785	(198)	3182	(277)	3556	(347)	3782	(403)	4700	F		
31-50	312	2725	(108)	1706	(129)	1892	(119)	2223	(113)	2635	(123)	3133	(159)	3707	(242)	4134	(334)	4700	F		
51-70	312	2944	(153)	1983	(262)	2149	(236)	2455	(192)	2838	(159)	3272	(189)	3710	(283)	3996	(358)	4700	F		
>70	239	2449	(102)	1308	(88)	1498	(91)	1869	(97)	2373	(108)	2991	(130)	3663	(169)	4126	(212)	4700	F		
19+	1060	2749	(69)	1618	(79)	1820	(72)	2188	(68)	2659	(75)	3221	(100)	3828	(146)	4248	(190)	4700	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.8 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	129	2292 (115)	1370 (170)	1566 (160)	1913 (145)	2325 (146)	2761 (171)	3173 (209)	3428 (245)	3000	F	
	4-8	213	2426 (81)	1804 (185)	1917 (161)	2115 (121)	2348 (99)	2598 (133)	2843 (216)	3001 (288)	3800	<3	
Male													
	9-13	122	2787 (140)	2081 (253)	2218 (229)	2458 (195)	2750 (187)	3083 (231)	3422 (319)	3645 (392)	4500	<3	
	14-18	150	3678 (296)	2301 (349)	2534 (330)	2954 (315)	3493 (345)	4135 (446)	4796 (595)	5220 (704)	4700	F	
	19-30	106	3546 (215)	1955 (374) ^E	2279 (328)	2850 (263)	3532 (266)	4286 (381)	5047 (562)	5545 (699)	4700	F	
	31-50	155	3585 (218)	1861 (377) ^E	2175 (347)	2780 (296)	3547 (268)	4378 (324)	5241 (471)	5860 (622)	4700	F	
	51-70	122	3335 (158)	2213 (342)	2454 (284)	2868 (210)	3355 (204)	3888 (311)	4425 (472)	4779 (591)	4700	F	
	>70	88	3453 (204)	2025 (213)	2226 (214)	2628 (216)	3205 (224)	3971 (278)	4869 (434)	5522 (600)	4700	F	
	19+	471	3494 (115)	1937 (146)	2237 (138)	2785 (132)	3457 (141)	4205 (177)	5004 (247)	5558 (312)	4700	14.4	(3.6) ^E
Female													
	9-13	103	2691 (219)	2030 (275)	2145 (254)	2351 (223)	2607 (212)	2901 (247)	3207 (334)	3414 (415)	4500	<3	
	14-18	142	2572 (148)	1529 (165)	1733 (154)	2095 (152)	2538 (183)	3045 (247)	3577 (329)	3936 (391)	4700	F	
	19-30	111	2589 (227)	1746 (338) ^E	1903 (317) ^E	2182 (296)	2518 (311)	2880 (377)	3230 (478)	3450 (554)	4700	<3	
	31-50	146	2858 (135)	1355 (295) ^E	1644 (256)	2116 (196)	2631 (152)	3243 (169)	3827 (238)	4164 (278)	4700	F	
	51-70	184	2924 (147)	2018 (116)	2210 (125)	2544 (151)	2945 (195)	3389 (252)	3840 (318)	4138 (365)	4700	F	
	>70	143	2724 (100)	1759 (166)	1972 (151)	2345 (135)	2776 (139)	3220 (168)	3652 (219)	3937 (262)	4700	<3	
	19+	584	2800 (77)	1629 (148)	1850 (137)	2244 (116)	2719 (97)	3242 (107)	3764 (151)	4104 (189)	4700	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.9 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	169	2148 (87)	1205 (135)	1387 (124)	1712 (111)	2109 (106)	2569 (124)	3067 (181)	3414 (240)	3000	11.4	(3.8) ^E
	4-8	281	2473 (76)	1561 (189)	1730 (161)	2035 (114)	2414 (86)	2845 (135)	3287 (230)	3581 (302)	3800		F
Male													
	9-13	183	3309 (220)	1971 (324)	2205 (299)	2656 (254)	3260 (231)	3994 (319)	4786 (528)	5329 (717)	4500		F
	14-18	187	3560 (165)	2130 (322)	2431 (281)	2964 (220)	3624 (198)	4378 (273)	5132 (411)	5609 (511)	4700		F
	19-30	223	3430 (161)	2116 (414) ^E	2378 (365)	2833 (274)	3358 (180)	3927 (364)	4509 (527)	4896 (646)	4700		F
	31-50	229	3458 (175)	2248 (316)	2480 (288)	2912 (246)	3486 (231)	4188 (304)	4938 (454)	5446 (577)	4700		F
	51-70	197	3208 (203)	1840 (267)	2073 (253)	2518 (232)	3075 (226)	3698 (280)	4387 (394)	4869 (487)	4700		F
	>70	72	2964 (174)	2090 (289)	2262 (265)	2582 (233)	2964 (221)	3354 (241)	3728 (307)	3982 (375)	4700		F
	19+	721	3355 (103)	2056 (163)	2307 (151)	2758 (134)	3320 (130)	3981 (159)	4692 (222)	5176 (280)	4700	9.9	(3.0) ^E
Female													
	9-13	165	2575 (138)	1615 (285) ^E	1805 (262)	2156 (221)	2597 (190)	3095 (220)	3595 (310)	3919 (382)	4500		F
	14-18	206	2425 (94)	1568 (244)	1730 (209)	2028 (151)	2392 (115)	2786 (161)	3190 (264)	3467 (349)	4700	<3	
	19-30	191	2573 (128)	1647 (258)	1818 (237)	2136 (203)	2541 (184)	3007 (216)	3483 (308)	3796 (389)	4700	<3	
	31-50	258	2713 (108)	2015 (283)	2168 (248)	2438 (190)	2763 (147)	3118 (171)	3476 (263)	3711 (346)	4700	<3	
	51-70	249	2683 (92)	1773 (202)	1959 (173)	2279 (130)	2655 (105)	3064 (131)	3471 (195)	3736 (248)	4700	<3	
	>70	128	2532 (131)	1584 (133)	1775 (127)	2096 (123)	2467 (139)	2898 (189)	3391 (262)	3751 (321)	4700	<3	
	19+	826	2657 (62)	1704 (110)	1889 (103)	2230 (90)	2657 (79)	3128 (87)	3599 (119)	3910 (150)	4700	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.10 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	2288 (71)	1570 (111)	1712 (103)	1967 (91)	2268 (92)	2563 (111)	2804 (133)	2943 (151)	3000	F	
	4-8	321	2521 (72)	1626 (125)	1793 (114)	2102 (96)	2496 (90)	2947 (114)	3409 (164)	3712 (206)	3800	F	
Male													
	9-13	226	2997 (113)	1794 (143)	2024 (135)	2437 (132)	2955 (144)	3589 (175)	4324 (258)	4867 (349)	4500	F	
	14-18	262	3979 (305)	2006 (311)	2389 (285)	3016 (277)	3816 (318)	4954 (408)	6252 (595)	7130 (764)	4700	29.4 (8.2) ^E	
	19-30	197	3649 (177)	2210 (307)	2452 (277)	2907 (234)	3519 (231)	4278 (314)	5076 (464)	5584 (572)	4700	F	
	31-50	282	4030 (182)	2306 (243)	2572 (244)	3156 (216)	3973 (200)	4805 (257)	5702 (405)	6382 (560)	4700	27.7 (6.9) ^E	
	51-70	234	3650 (143)	2132 (199)	2401 (190)	2891 (171)	3475 (172)	4128 (214)	4826 (300)	5305 (378)	4700	F	
	>70	119	3388 (247)	1688 (318) ^E	1965 (297)	2497 (267)	3200 (275)	4033 (378)	4904 (539)	5484 (663)	4700	F	
	19+	832	3778 (91)	2159 (107)	2451 (99)	2987 (94)	3664 (106)	4449 (140)	5272 (199)	5826 (251)	4700	19.2 (3.2) ^E	
Female													
	9-13	226	2693 (160)	1699 (190)	1884 (179)	2223 (165)	2653 (172)	3153 (225)	3681 (317)	4038 (390)	4500	F	
	14-18	242	2663 (133)	1282 (156)	1497 (162)	1940 (161)	2551 (166)	3246 (199)	4031 (276)	4595 (352)	4700	F	
	19-30	208	2726 (137)	1948 (276)	2125 (245)	2422 (201)	2797 (175)	3220 (213)	3588 (292)	3809 (365)	4700	<3	
	31-50	263	3005 (124)	1932 (207)	2125 (183)	2460 (155)	2867 (156)	3341 (216)	3860 (335)	4229 (439)	4700	F	
	51-70	322	2806 (119)	1570 (180)	1793 (167)	2211 (149)	2748 (144)	3366 (174)	3998 (241)	4411 (299)	4700	F	
	>70	198	2801 (121)	1670 (174)	1897 (175)	2317 (177)	2824 (176)	3337 (172)	3780 (179)	4044 (192)	4700	<3	
	19+	991	2872 (63)	1777 (91)	1990 (89)	2367 (83)	2826 (83)	3345 (104)	3888 (140)	4256 (178)	4700	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.11 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI	(SE)								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)				95th (SE)							
Both																					
	1-3	348	2377	(70)	1595	(117)	1751	(108)	2032	(93)	2377	(88)	2760	(108)	3137	(146)	3378	(174)	3000	14.3	(4.7) ^E
	4-8	554	2650	(90)	1740	(99)	1911	(96)	2231	(95)	2644	(104)	3104	(132)	3591	(188)	3938	(242)	3800	F	
Male																					
	9-13	409	3072	(102)	2027	(144)	2212	(136)	2554	(124)	2982	(123)	3467	(150)	3963	(206)	4293	(254)	4500	F	
	14-18	414	3364	(132)	2069	(188)	2308	(186)	2760	(180)	3342	(182)	4027	(211)	4754	(268)	5239	(317)	4700	F	
	19-30	311	3676	(142)	2184	(207)	2482	(186)	3020	(169)	3707	(172)	4469	(208)	5213	(292)	5713	(385)	4700	19.2	(5.1) ^E
	31-50	489	3579	(125)	2308	(194)	2539	(176)	2928	(157)	3463	(145)	4072	(178)	4673	(252)	5067	(313)	4700	F	
	51-70	575	3333	(98)	2078	(144)	2303	(135)	2722	(128)	3275	(125)	3888	(144)	4456	(183)	4822	(221)	4700	F	
	>70	239	3127	(126)	2030	(150)	2220	(152)	2587	(168)	3099	(180)	3688	(177)	4241	(213)	4604	(266)	4700	F	
	19+	1614	3484	(66)	2097	(86)	2344	(81)	2802	(79)	3410	(81)	4109	(96)	4795	(126)	5241	(154)	4700	11.5	(2.0) ^E
Female																					
	9-13	355	2654	(108)	1864	(151)	2025	(144)	2317	(135)	2682	(135)	3095	(157)	3505	(201)	3770	(241)	4500	<3	
	14-18	410	2485	(89)	1459	(168)	1673	(152)	2050	(121)	2475	(109)	2939	(134)	3434	(190)	3774	(236)	4700	<3	
	19-30	384	2604	(111)	1670	(145)	1848	(139)	2185	(134)	2617	(144)	3089	(172)	3528	(211)	3806	(246)	4700	<3	
	31-50	585	2751	(78)	1470	(113)	1732	(110)	2209	(109)	2781	(113)	3384	(127)	3970	(155)	4352	(180)	4700	F	
	51-70	711	2675	(70)	1533	(87)	1746	(82)	2117	(77)	2571	(84)	3106	(109)	3655	(142)	4007	(169)	4700	<3	
	>70	342	2634	(94)	1517	(109)	1720	(103)	2090	(101)	2552	(114)	3075	(136)	3617	(177)	3985	(213)	4700	<3	
	19+	2022	2687	(43)	1527	(58)	1753	(56)	2165	(56)	2675	(60)	3232	(69)	3772	(84)	4119	(98)	4700	1.4	(0.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.12 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI	(SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)										
Both																					
	1-3	622	2192	(57)	1218	(78)	1410	(74)	1753	(68)	2168	(69)	2632	(81)	3122	(110)	3462	(143)	3000	12.7	(2.5) ^E
	4-8	919	2435	(52)	1660	(120)	1805	(105)	2067	(79)	2388	(60)	2749	(81)	3116	(134)	3359	(178)	3800	F	
Male																					
	9-13	579	3112	(135)	2099	(218)	2297	(199)	2659	(167)	3111	(153)	3628	(198)	4171	(305)	4543	(401)	4500	F	
	14-18	634	3611	(116)	2055	(159)	2376	(144)	2934	(127)	3584	(138)	4350	(186)	5212	(257)	5787	(313)	4700	17.5	(4.1) ^E
	19-30	578	3502	(126)	2064	(268)	2338	(230)	2822	(172)	3396	(143)	4039	(196)	4718	(317)	5181	(418)	4700	F	
	31-50	693	3425	(120)	2063	(194)	2330	(179)	2807	(161)	3424	(159)	4136	(183)	4900	(247)	5419	(311)	4700	12.9	(3.6) ^E
	51-70	596	3231	(125)	1884	(163)	2133	(156)	2591	(143)	3151	(143)	3779	(174)	4452	(239)	4908	(294)	4700	F	
	>70	296	3092	(111)	1803	(152)	2032	(144)	2445	(141)	2965	(138)	3583	(144)	4262	(196)	4743	(260)	4700	F	
	19+	2163	3366	(69)	1959	(91)	2223	(87)	2712	(83)	3313	(87)	3999	(103)	4741	(135)	5260	(168)	4700	10.6	(1.9) ^E
Female																					
	9-13	533	2566	(98)	1639	(180)	1817	(165)	2139	(140)	2536	(123)	2986	(141)	3445	(196)	3748	(244)	4500	<3	
	14-18	638	2460	(67)	1472	(83)	1654	(78)	1982	(75)	2390	(79)	2859	(99)	3365	(140)	3720	(175)	4700	<3	
	19-30	499	2612	(94)	1720	(178)	1887	(163)	2196	(141)	2588	(130)	3031	(156)	3471	(217)	3756	(269)	4700	<3	
	31-50	716	2740	(75)	1697	(166)	1893	(152)	2259	(124)	2706	(101)	3193	(106)	3709	(154)	4074	(208)	4700	F	
	51-70	745	2788	(68)	1910	(149)	2083	(130)	2388	(101)	2757	(80)	3169	(97)	3588	(150)	3862	(195)	4700	<3	
	>70	510	2556	(72)	1454	(75)	1669	(71)	2053	(70)	2523	(82)	3048	(106)	3623	(138)	4015	(161)	4700	<3	
	19+	2470	2704	(43)	1649	(66)	1847	(63)	2210	(58)	2668	(54)	3181	(59)	3703	(77)	4056	(96)	4700	1.3	(0.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 21.13 Potassium (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										AI ²	% >AI	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)				
Both														
	1-3	2117	2331 (33)	1378 (50)	1568 (47)	1908 (42)	2321 (40)	2766 (45)	3224 (58)	3538 (71)	3000	15.9	(1.6)	
	4-8	3235	2591 (31)	1639 (47)	1824 (41)	2151 (36)	2549 (37)	3012 (47)	3488 (68)	3800 (87)	3800	5.0	(1.0) ^E	
Male														
	9-13	2080	3153 (59)	1955 (76)	2175 (73)	2583 (67)	3096 (69)	3685 (88)	4307 (124)	4730 (156)	4500	7.3	(1.7) ^E	
	14-18	2288	3635 (69)	2142 (92)	2438 (87)	2967 (80)	3637 (81)	4437 (99)	5289 (139)	5867 (181)	4700	19.2	(2.2)	
	19-30	1804	3552 (76)	2127 (104)	2388 (97)	2867 (89)	3469 (91)	4184 (117)	4958 (166)	5476 (206)	4700	13.8	(2.5) ^E	
	31-50	2596	3534 (58)	2093 (85)	2359 (81)	2862 (71)	3490 (68)	4193 (85)	4923 (123)	5421 (157)	4700	13.4	(1.9)	
	51-70	2550	3403 (50)	2023 (63)	2278 (60)	2741 (56)	3318 (61)	3977 (77)	4657 (105)	5111 (128)	4700	9.4	(1.5)	
	>70	1520	3059 (68)	1739 (83)	1977 (79)	2416 (80)	2984 (82)	3632 (93)	4285 (120)	4723 (147)	4700	5.2	(1.2) ^E	
	19+	8470	3460 (33)	2031 (44)	2296 (42)	2783 (39)	3394 (40)	4092 (49)	4826 (68)	5322 (86)	4700	11.8	(1.0)	
Female														
	9-13	1980	2664 (50)	1605 (56)	1796 (52)	2145 (48)	2578 (51)	3070 (67)	3579 (97)	3921 (124)	4500	<3		
	14-18	2256	2669 (40)	1560 (54)	1771 (51)	2151 (48)	2632 (49)	3195 (60)	3788 (81)	4186 (99)	4700	1.9	(0.5) ^E	
	19-30	1854	2674 (50)	1667 (75)	1852 (71)	2195 (64)	2627 (63)	3107 (76)	3578 (100)	3883 (122)	4700	<3		
	31-50	2686	2874 (41)	1677 (47)	1896 (44)	2296 (45)	2810 (49)	3398 (64)	4034 (91)	4469 (121)	4700	3.4	(0.8) ^E	
	51-70	3200	2851 (39)	1725 (60)	1941 (55)	2328 (49)	2804 (47)	3344 (54)	3894 (73)	4253 (88)	4700	1.9	(0.5) ^E	
	>70	2610	2624 (39)	1526 (51)	1732 (50)	2110 (49)	2582 (52)	3112 (58)	3642 (69)	3985 (79)	4700	1.0	(0.3) ^E	
	19+	10350	2798 (23)	1661 (27)	1877 (26)	2262 (25)	2750 (27)	3300 (35)	3870 (48)	4254 (61)	4700	2.1	(0.3)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

22. Riboflavin (mg/d): Usual intakes from food

Table 22.1 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	1.77 (0.17)	1.27 (0.21) ^E	1.38 (0.19)	1.55 (0.17)	1.76 (0.17)	2.00 (0.23)	2.24 (0.32)	2.40 (0.40) ^E	0.4	<3	
	4-8	127	2.01 (0.12)	1.42 (0.19)	1.54 (0.18)	1.74 (0.16)	2.00 (0.14)	2.29 (0.19)	2.58 (0.23)	2.78 (0.27)	0.5	<3	
Male	9-13	111	2.34 (0.18)	1.49 (0.22)	1.64 (0.20)	1.91 (0.17)	2.26 (0.18)	2.69 (0.27)	3.16 (0.41)	3.50 (0.52)	0.8	<3	
	14-18	107	2.25 (0.17)	1.47 (0.25) ^E	1.63 (0.23)	1.93 (0.20)	2.32 (0.19)	2.78 (0.26)	3.26 (0.38)	3.58 (0.48)	1.1	<3	
	19-30	77	2.03 (0.12)	1.33 (0.11)	1.46 (0.12)	1.72 (0.14)	2.05 (0.16)	2.41 (0.18)	2.79 (0.22)	3.04 (0.26)	1.1	<3	
	31-50	145	2.27 (0.25)	1.38 (0.23) ^E	1.51 (0.22)	1.76 (0.19)	2.09 (0.19)	2.51 (0.24)	3.00 (0.35)	3.36 (0.47)	1.1	F	
	51-70	182	1.79 (0.08)	1.11 (0.21) ^E	1.23 (0.19)	1.45 (0.14)	1.73 (0.10)	2.03 (0.13)	2.31 (0.19)	2.48 (0.25)	1.1	F	
	>70	63	2.05 (0.14)	1.60 (0.23)	1.70 (0.22)	1.87 (0.21)	2.09 (0.21)	2.32 (0.24)	2.56 (0.29)	2.71 (0.33)	1.1	<3	
	19+	467	2.06 (0.11)	1.36 (0.12)	1.48 (0.11)	1.70 (0.10)	1.99 (0.09)	2.33 (0.11)	2.70 (0.17)	2.96 (0.22)	1.1	<3	
Female	9-13	96	1.97 (0.17)	1.42 (0.24) ^E	1.53 (0.22)	1.73 (0.19)	1.97 (0.19)	2.24 (0.23)	2.53 (0.31)	2.72 (0.38)	0.8	<3	
	14-18	105	1.75 (0.16)	0.97 (0.20) ^E	1.10 (0.20) ^E	1.34 (0.19)	1.66 (0.19)	2.11 (0.29)	2.64 (0.41)	3.01 (0.49)	0.9	F	
	19-30	91	1.47 (0.10)	0.85 (0.12)	0.95 (0.12)	1.13 (0.12)	1.37 (0.13)	1.68 (0.17)	2.01 (0.26)	2.25 (0.35)	0.9	F	
	31-50	167	1.49 (0.08)	0.83 (0.12)	0.97 (0.11)	1.19 (0.10)	1.48 (0.11)	1.82 (0.14)	2.16 (0.16)	2.36 (0.18)	0.9	F	
	51-70	198	1.74 (0.17)	0.96 (0.18) ^E	1.09 (0.16)	1.33 (0.14)	1.60 (0.15)	1.99 (0.24)	2.66 (0.51) ^E	3.33 (0.86) ^E	0.9	F	
	>70	74	1.68 (0.15)	0.99 (0.11)	1.10 (0.12)	1.31 (0.14)	1.58 (0.16)	1.91 (0.19)	2.26 (0.23)	2.49 (0.26)	0.9	F	
	19+	530	1.58 (0.06)	0.94 (0.07)	1.06 (0.06)	1.26 (0.06)	1.52 (0.06)	1.87 (0.09)	2.29 (0.16)	2.61 (0.26)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.2 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both												
	1-3	58	1.91 (0.19)	1.10 (0.21) ^E	1.25 (0.20)	1.52 (0.20)	1.86 (0.22)	2.25 (0.27)	2.64 (0.35)	2.90 (0.40)	0.4	<3
	4-8	110	2.21 (0.26)	1.25 (0.23) ^E	1.43 (0.23)	1.75 (0.25)	2.17 (0.29)	2.64 (0.37)	3.12 (0.47)	3.42 (0.54)	0.5	<3
Male												
	9-13	95	2.14 (0.12)	1.69 (0.20)	1.76 (0.19)	1.90 (0.18)	2.07 (0.19)	2.25 (0.23)	2.43 (0.30)	2.55 (0.35)	0.8	<3
	14-18	87	2.51 (0.20)	1.29 (0.25) ^E	1.51 (0.25)	1.95 (0.24)	2.54 (0.26)	3.25 (0.33)	4.01 (0.45)	4.52 (0.56)	1.1	F
	19-30	70	3.11 (0.36)	1.66 (0.40) ^E	1.92 (0.40) ^E	2.45 (0.39)	3.21 (0.44)	4.12 (0.61)	5.08 (0.81)	5.70 (0.96) ^E	1.1	<3
	31-50	109	1.98 (0.15)	1.09 (0.19) ^E	1.23 (0.18)	1.50 (0.16)	1.83 (0.16)	2.21 (0.21)	2.65 (0.32)	2.96 (0.40)	1.1	F
	51-70	128	1.99 (0.10)	1.37 (0.17)	1.47 (0.15)	1.67 (0.12)	1.93 (0.11)	2.23 (0.14)	2.55 (0.22)	2.78 (0.30)	1.1	<3
	>70	65	1.76 (0.12)	1.03 (0.13)	1.15 (0.13)	1.38 (0.13)	1.70 (0.14)	2.10 (0.18)	2.55 (0.30)	2.87 (0.40)	1.1	F
	19+	372	2.20 (0.10)	1.21 (0.11)	1.36 (0.10)	1.67 (0.10)	2.08 (0.11)	2.57 (0.16)	3.19 (0.26)	3.68 (0.36)	1.1	F
Female												
	9-13	75	1.95 (0.15)	1.25 (0.24) ^E	1.40 (0.23)	1.68 (0.22)	2.01 (0.23)	2.37 (0.27)	2.72 (0.34)	2.93 (0.39)	0.8	<3
	14-18	81	1.74 (0.14)	0.76 (0.20) ^E	0.94 (0.19) ^E	1.27 (0.17)	1.68 (0.17)	2.14 (0.20)	2.58 (0.25)	2.86 (0.29)	0.9	F
	19-30	101	1.84 (0.12)	1.06 (0.14)	1.24 (0.14)	1.56 (0.14)	1.96 (0.16)	2.40 (0.19)	2.83 (0.24)	3.10 (0.27)	0.9	F
	31-50	116	1.68 (0.11)	1.01 (0.16)	1.14 (0.14)	1.39 (0.13)	1.70 (0.14)	2.04 (0.19)	2.38 (0.26)	2.60 (0.30)	0.9	F
	51-70	146	1.65 (0.07)	0.99 (0.09)	1.11 (0.09)	1.32 (0.09)	1.60 (0.09)	1.91 (0.10)	2.23 (0.12)	2.44 (0.15)	0.9	F
	>70	94	1.48 (0.11)	0.85 (0.14)	0.94 (0.13)	1.12 (0.12)	1.37 (0.12)	1.70 (0.16)	2.07 (0.24)	2.34 (0.31)	0.9	F
	19+	457	1.68 (0.06)	0.98 (0.06)	1.11 (0.06)	1.35 (0.07)	1.66 (0.08)	2.03 (0.10)	2.40 (0.12)	2.63 (0.14)	0.9	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.3 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	112	1.67 (0.08)	1.20 (0.14)	1.28 (0.13)	1.44 (0.11)	1.64 (0.09)	1.85 (0.11)	2.08 (0.16)	2.23 (0.21)	0.4	0.0 (0.0)
	4-8	177	2.04 (0.10)	1.29 (0.16)	1.44 (0.15)	1.72 (0.12)	2.05 (0.11)	2.43 (0.14)	2.81 (0.20)	3.06 (0.26)	0.5	<3
Male	9-13	111	2.33 (0.14)	1.51 (0.24)	1.64 (0.22)	1.89 (0.18)	2.21 (0.16)	2.58 (0.20)	2.99 (0.31)	3.27 (0.41)	0.8	<3
	14-18	113	2.32 (0.18)	1.40 (0.24) ^E	1.57 (0.23)	1.86 (0.24)	2.22 (0.27)	2.64 (0.31)	3.10 (0.37)	3.39 (0.41)	1.1	F
	19-30	91	2.51 (0.23)	1.20 (0.19)	1.41 (0.19)	1.81 (0.21)	2.33 (0.23)	2.95 (0.29)	3.62 (0.40)	4.08 (0.49)	1.1	F
	31-50	101	2.17 (0.12)	1.45 (0.20)	1.59 (0.17)	1.83 (0.13)	2.13 (0.12)	2.47 (0.17)	2.82 (0.27)	3.06 (0.34)	1.1	<3
	51-70	134	1.99 (0.07)	1.54 (0.16)	1.63 (0.14)	1.79 (0.10)	1.98 (0.08)	2.17 (0.10)	2.36 (0.15)	2.48 (0.20)	1.1	<3
	>70	56	1.85 (0.15)	1.40 (0.19)	1.47 (0.18)	1.60 (0.17)	1.76 (0.17)	1.93 (0.20)	2.10 (0.27)	2.21 (0.35)	1.1	F
	19+	382	2.15 (0.07)	1.38 (0.10)	1.52 (0.09)	1.77 (0.07)	2.08 (0.07)	2.43 (0.10)	2.80 (0.15)	3.06 (0.19)	1.1	<3
	Female	9-13	105	1.84 (0.15)	1.20 (0.20) ^E	1.30 (0.19)	1.50 (0.17)	1.78 (0.17)	2.09 (0.21)	2.39 (0.28)	2.58 (0.33)	0.8
14-18		120	1.77 (0.16)	0.84 (0.23) ^E	1.07 (0.22) ^E	1.46 (0.19)	1.82 (0.17)	2.18 (0.22)	2.62 (0.36)	2.95 (0.48)	0.9	F
19-30		91	1.70 (0.10)	1.25 (0.20)	1.35 (0.17)	1.53 (0.15)	1.71 (0.14)	1.92 (0.19)	2.17 (0.28)	2.34 (0.36)	0.9	<3
31-50		159	1.76 (0.09)	0.87 (0.11)	1.04 (0.11)	1.35 (0.11)	1.76 (0.11)	2.22 (0.14)	2.69 (0.20)	2.99 (0.25)	0.9	F
51-70		174	1.63 (0.13)	0.93 (0.13)	1.05 (0.12)	1.26 (0.11)	1.52 (0.11)	1.87 (0.16)	2.27 (0.23)	2.55 (0.29)	0.9	F
>70		80	1.72 (0.15)	0.90 (0.13)	1.03 (0.14)	1.28 (0.14)	1.64 (0.18)	2.14 (0.26)	2.77 (0.37)	3.25 (0.49)	0.9	F
19+		504	1.71 (0.06)	0.97 (0.07)	1.12 (0.07)	1.36 (0.07)	1.65 (0.07)	2.07 (0.10)	2.52 (0.14)	2.80 (0.16)	0.9	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.4 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both												
	1-3	99	2.07 (0.16)	1.42 (0.20)	1.56 (0.17)	1.79 (0.15)	2.02 (0.16)	2.31 (0.23)	2.69 (0.35)	2.97 (0.43)	0.4	0.0 (0.0)
	4-8	140	2.07 (0.12)	1.45 (0.17)	1.57 (0.15)	1.77 (0.12)	2.02 (0.11)	2.30 (0.15)	2.59 (0.23)	2.80 (0.32)	0.5	0.0 (0.0)
Male												
	9-13	92	2.23 (0.14)	1.43 (0.26) ^E	1.58 (0.23)	1.85 (0.19)	2.13 (0.18)	2.43 (0.20)	2.75 (0.25)	2.97 (0.31)	0.8	<3
	14-18	107	2.94 (0.27)	1.84 (0.28)	2.07 (0.27)	2.49 (0.26)	3.02 (0.29)	3.68 (0.40)	4.49 (0.59)	5.10 (0.77)	1.1	<3
	19-30	73	2.90 (0.30)	1.30 (0.28) ^E	1.57 (0.28) ^E	2.09 (0.28)	2.81 (0.34)	3.83 (0.52)	5.23 (0.88) ^E	6.40 (1.23) ^E	1.1	F
	31-50	134	2.24 (0.19)	1.49 (0.23)	1.60 (0.21)	1.80 (0.19)	2.08 (0.19)	2.42 (0.22)	2.77 (0.30)	2.99 (0.37)	1.1	F
	51-70	131	1.98 (0.11)	1.18 (0.16)	1.31 (0.15)	1.56 (0.14)	1.87 (0.14)	2.25 (0.16)	2.64 (0.22)	2.91 (0.27)	1.1	F
	>70	55	1.88 (0.13)	1.09 (0.20) ^E	1.23 (0.19)	1.47 (0.17)	1.76 (0.16)	2.11 (0.20)	2.47 (0.28)	2.72 (0.38)	1.1	F
	19+	393	2.27 (0.10)	1.20 (0.09)	1.36 (0.09)	1.69 (0.10)	2.13 (0.12)	2.72 (0.16)	3.44 (0.23)	3.95 (0.30)	1.1	F
Female												
	9-13	79	2.20 (0.17)	1.29 (0.19)	1.45 (0.19)	1.75 (0.19)	2.15 (0.22)	2.71 (0.29)	3.34 (0.40)	3.79 (0.51)	0.8	<3
	14-18	104	1.85 (0.11)	1.20 (0.15)	1.34 (0.13)	1.54 (0.11)	1.85 (0.14)	2.22 (0.18)	2.54 (0.22)	2.76 (0.26)	0.9	F
	19-30	101	1.51 (0.14)	0.74 (0.22) ^E	0.87 (0.22) ^E	1.13 (0.19) ^E	1.47 (0.18)	1.89 (0.20)	2.37 (0.28)	2.70 (0.35)	0.9	F
	31-50	143	1.70 (0.11)	0.93 (0.15)	1.05 (0.14)	1.29 (0.12)	1.58 (0.12)	1.95 (0.16)	2.39 (0.29)	2.75 (0.44)	0.9	F
	51-70	193	1.69 (0.08)	0.99 (0.13)	1.09 (0.12)	1.30 (0.11)	1.59 (0.10)	1.93 (0.13)	2.31 (0.19)	2.58 (0.26)	0.9	F
	>70	94	1.54 (0.11)	0.94 (0.14)	1.03 (0.14)	1.19 (0.14)	1.42 (0.15)	1.71 (0.17)	2.03 (0.23)	2.25 (0.28)	0.9	F
	19+	531	1.64 (0.06)	0.87 (0.07)	1.00 (0.06)	1.24 (0.06)	1.57 (0.07)	1.95 (0.08)	2.38 (0.12)	2.70 (0.16)	0.9	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.5 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	311	1.84	(0.06)	1.22 (0.12)	1.36 (0.11)	1.60 (0.10)	1.87 (0.09)	2.18 (0.10)	2.48 (0.12)	2.68 (0.14)	0.4	<3	
	4-8	485	2.06	(0.07)	1.30 (0.11)	1.43 (0.10)	1.68 (0.09)	2.01 (0.09)	2.40 (0.12)	2.80 (0.17)	3.06 (0.21)	0.5	0.0	(0.0)
Male														
	9-13	277	2.59	(0.11)	1.66 (0.18)	1.83 (0.17)	2.17 (0.15)	2.63 (0.15)	3.20 (0.22)	3.80 (0.32)	4.19 (0.40)	0.8	<3	
	14-18	339	2.85	(0.14)	1.69 (0.18)	1.90 (0.17)	2.29 (0.17)	2.84 (0.17)	3.53 (0.21)	4.31 (0.29)	4.87 (0.38)	1.1	<3	
	19-30	237	2.57	(0.15)	1.66 (0.21)	1.80 (0.20)	2.09 (0.18)	2.49 (0.18)	2.99 (0.26)	3.55 (0.41)	3.93 (0.54)	1.1	<3	
	31-50	423	2.37	(0.11)	1.57 (0.18)	1.72 (0.17)	2.00 (0.14)	2.36 (0.13)	2.77 (0.16)	3.17 (0.22)	3.43 (0.27)	1.1	<3	
	51-70	387	2.11	(0.08)	1.26 (0.12)	1.41 (0.11)	1.69 (0.09)	2.05 (0.09)	2.48 (0.13)	2.97 (0.20)	3.31 (0.26)	1.1	F	
	>70	132	1.94	(0.17)	0.91 (0.21) ^E	1.07 (0.21) ^E	1.38 (0.20)	1.81 (0.21)	2.34 (0.26)	2.93 (0.33)	3.34 (0.40)	1.1	F	
	19+	1179	2.30	(0.06)	1.36 (0.07)	1.52 (0.07)	1.82 (0.07)	2.23 (0.08)	2.73 (0.09)	3.26 (0.12)	3.61 (0.15)	1.1	<3	
Female														
	9-13	281	2.07	(0.10)	1.15 (0.16)	1.28 (0.16)	1.57 (0.14)	1.96 (0.11)	2.36 (0.13)	2.73 (0.18)	2.96 (0.23)	0.8	<3	
	14-18	321	1.88	(0.06)	1.23 (0.10)	1.36 (0.10)	1.60 (0.09)	1.92 (0.09)	2.26 (0.10)	2.57 (0.12)	2.77 (0.15)	0.9	<3	
	19-30	249	1.91	(0.09)	1.25 (0.15)	1.38 (0.13)	1.61 (0.11)	1.89 (0.10)	2.22 (0.15)	2.56 (0.24)	2.77 (0.31)	0.9	<3	
	31-50	364	1.86	(0.07)	1.24 (0.12)	1.37 (0.11)	1.59 (0.09)	1.87 (0.09)	2.20 (0.10)	2.56 (0.13)	2.79 (0.16)	0.9	<3	
	51-70	467	1.74	(0.05)	1.05 (0.09)	1.19 (0.08)	1.43 (0.07)	1.72 (0.06)	2.05 (0.08)	2.39 (0.11)	2.62 (0.15)	0.9	F	
	>70	215	1.59	(0.12)	1.08 (0.14)	1.19 (0.14)	1.39 (0.14)	1.63 (0.15)	1.89 (0.16)	2.14 (0.20)	2.31 (0.22)	0.9	F	
	19+	1295	1.80	(0.04)	1.16 (0.06)	1.29 (0.06)	1.52 (0.05)	1.81 (0.05)	2.13 (0.06)	2.47 (0.08)	2.69 (0.09)	0.9	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.6 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	644	1.84	(0.06)	1.27 (0.18)	1.39 (0.15)	1.61 (0.11)	1.88 (0.07)	2.18 (0.10)	2.51 (0.19)	2.74 (0.27)	0.4	<3	
	4-8	956	2.07	(0.11)	1.20 (0.08)	1.34 (0.07)	1.59 (0.06)	1.92 (0.06)	2.37 (0.09)	2.97 (0.22)	3.46 (0.38)	0.5	0.0	(0.0)
Male														
	9-13	589	2.39	(0.08)	1.31 (0.17)	1.46 (0.16)	1.77 (0.14)	2.21 (0.10)	2.79 (0.11)	3.48 (0.24)	4.04 (0.39)	0.8	<3	
	14-18	639	2.51	(0.07)	1.57 (0.15)	1.76 (0.13)	2.10 (0.10)	2.52 (0.08)	3.02 (0.11)	3.55 (0.18)	3.90 (0.24)	1.1	<3	
	19-30	481	2.30	(0.09)	1.27 (0.17)	1.46 (0.15)	1.79 (0.12)	2.21 (0.11)	2.74 (0.14)	3.34 (0.25)	3.78 (0.34)	1.1	F	
	31-50	709	2.05	(0.06)	1.12 (0.11)	1.27 (0.11)	1.57 (0.10)	1.99 (0.08)	2.51 (0.09)	3.08 (0.15)	3.48 (0.21)	1.1	F	
	51-70	758	1.93	(0.05)	1.16 (0.11)	1.29 (0.10)	1.51 (0.08)	1.82 (0.06)	2.24 (0.08)	2.67 (0.14)	2.94 (0.19)	1.1	F	
	>70	734	1.75	(0.04)	0.93 (0.06)	1.07 (0.06)	1.32 (0.05)	1.65 (0.05)	2.06 (0.06)	2.51 (0.10)	2.81 (0.13)	1.1	11.6	(2.6) ^E
	19+	2682	2.04	(0.04)	1.11 (0.06)	1.26 (0.06)	1.57 (0.05)	1.96 (0.04)	2.45 (0.05)	3.03 (0.08)	3.43 (0.12)	1.1	4.8	(1.3) ^E
Female														
	9-13	585	1.94	(0.06)	1.01 (0.08)	1.17 (0.07)	1.46 (0.06)	1.83 (0.06)	2.28 (0.09)	2.78 (0.13)	3.13 (0.18)	0.8	<3	
	14-18	645	1.90	(0.06)	1.02 (0.10)	1.18 (0.09)	1.48 (0.07)	1.86 (0.07)	2.31 (0.10)	2.79 (0.16)	3.12 (0.22)	0.9	F	
	19-30	514	1.61	(0.05)	1.03 (0.12)	1.13 (0.10)	1.32 (0.08)	1.56 (0.06)	1.84 (0.08)	2.12 (0.14)	2.30 (0.19)	0.9	F	
	31-50	758	1.67	(0.05)	0.94 (0.07)	1.07 (0.07)	1.31 (0.06)	1.62 (0.05)	2.00 (0.07)	2.42 (0.11)	2.71 (0.15)	0.9	F	
	51-70	955	1.61	(0.04)	0.91 (0.09)	1.04 (0.08)	1.26 (0.06)	1.54 (0.05)	1.87 (0.06)	2.23 (0.10)	2.48 (0.14)	0.9	F	
	>70	1345	1.54	(0.03)	0.90 (0.06)	1.01 (0.05)	1.21 (0.04)	1.47 (0.04)	1.77 (0.05)	2.10 (0.08)	2.34 (0.11)	0.9	F	
	19+	3572	1.63	(0.03)	0.97 (0.04)	1.09 (0.04)	1.30 (0.03)	1.57 (0.03)	1.90 (0.04)	2.26 (0.06)	2.51 (0.08)	0.9	3.2	(1.0) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.7 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	324	1.79	(0.07)	0.99 (0.11)	1.14 (0.10)	1.43 (0.08)	1.76 (0.08)	2.15 (0.13)	2.61 (0.21)	2.94 (0.27)	0.4	<3	
	4-8	425	1.99	(0.11)	1.24 (0.05)	1.36 (0.06)	1.59 (0.07)	1.89 (0.10)	2.28 (0.15)	2.72 (0.24)	3.06 (0.33)	0.5	0.0	(0.0)
Male														
	9-13	274	2.40	(0.09)	1.56 (0.08)	1.71 (0.08)	1.99 (0.09)	2.34 (0.11)	2.75 (0.14)	3.18 (0.18)	3.48 (0.22)	0.8	<3	
	14-18	297	2.91	(0.15)	1.22 (0.15)	1.47 (0.16)	1.99 (0.17)	2.77 (0.17)	3.72 (0.23)	4.89 (0.37)	5.79 (0.51)	1.1	F	
	19-30	249	2.55	(0.21)	1.19 (0.27) ^E	1.38 (0.25) ^E	1.76 (0.21)	2.29 (0.17)	2.97 (0.25)	3.76 (0.43)	4.33 (0.59)	1.1	F	
	31-50	309	2.07	(0.11)	1.29 (0.21) ^E	1.43 (0.19)	1.68 (0.15)	2.00 (0.12)	2.42 (0.19)	2.87 (0.32)	3.16 (0.42)	1.1	F	
	51-70	277	1.92	(0.09)	1.13 (0.13)	1.27 (0.12)	1.52 (0.10)	1.85 (0.10)	2.26 (0.12)	2.72 (0.20)	3.05 (0.28)	1.1	F	
	>70	136	1.88	(0.14)	0.85 (0.17) ^E	1.00 (0.16)	1.31 (0.13)	1.74 (0.14)	2.29 (0.22)	2.90 (0.34)	3.34 (0.43)	1.1	F	
	19+	971	2.12	(0.07)	1.11 (0.08)	1.27 (0.07)	1.56 (0.07)	1.98 (0.07)	2.51 (0.10)	3.11 (0.15)	3.54 (0.21)	1.1	F	
Female														
	9-13	265	2.03	(0.14)	1.25 (0.15)	1.37 (0.13)	1.59 (0.10)	1.86 (0.09)	2.19 (0.14)	2.54 (0.23)	2.80 (0.33)	0.8	<3	
	14-18	290	1.85	(0.09)	1.25 (0.19)	1.36 (0.17)	1.55 (0.13)	1.79 (0.10)	2.07 (0.13)	2.37 (0.22)	2.58 (0.30)	0.9	<3	
	19-30	197	1.91	(0.15)	1.08 (0.23) ^E	1.23 (0.21) ^E	1.50 (0.18)	1.88 (0.17)	2.30 (0.23)	2.70 (0.33)	2.97 (0.42)	0.9	F	
	31-50	312	1.68	(0.07)	0.95 (0.09)	1.08 (0.09)	1.34 (0.08)	1.61 (0.08)	1.91 (0.11)	2.27 (0.16)	2.53 (0.20)	0.9	F	
	51-70	312	1.77	(0.12)	1.15 (0.17)	1.25 (0.15)	1.44 (0.13)	1.70 (0.11)	2.01 (0.16)	2.37 (0.27)	2.62 (0.37)	0.9	<3	
	>70	239	1.42	(0.07)	0.79 (0.14) ^E	0.90 (0.13)	1.11 (0.10)	1.39 (0.07)	1.75 (0.11)	2.15 (0.20)	2.43 (0.29)	0.9	F	
	19+	1060	1.71	(0.05)	0.93 (0.06)	1.07 (0.06)	1.31 (0.05)	1.62 (0.05)	2.01 (0.07)	2.49 (0.13)	2.84 (0.18)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.8 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	1.83 (0.11)	0.92 (0.15) ^E	1.10 (0.14)	1.43 (0.14)	1.84 (0.15)	2.30 (0.18)	2.76 (0.21)	3.05 (0.24)	0.4	<3	
	4-8	213	1.86 (0.06)	1.25 (0.11)	1.36 (0.10)	1.56 (0.08)	1.81 (0.07)	2.09 (0.09)	2.36 (0.14)	2.54 (0.17)	0.5	0.0	(0.0)
Male	9-13	122	2.23 (0.16)	1.58 (0.21)	1.70 (0.19)	1.91 (0.19)	2.19 (0.20)	2.52 (0.25)	2.86 (0.31)	3.07 (0.36)	0.8	<3	
	14-18	150	2.66 (0.14)	1.63 (0.21)	1.81 (0.20)	2.15 (0.18)	2.59 (0.17)	3.10 (0.21)	3.63 (0.31)	3.98 (0.40)	1.1	<3	
	19-30	106	2.40 (0.14)	1.25 (0.25) ^E	1.47 (0.23)	1.86 (0.18)	2.33 (0.17)	2.87 (0.22)	3.45 (0.34)	3.86 (0.46)	1.1	F	
	31-50	155	2.30 (0.20)	1.06 (0.19) ^E	1.27 (0.18)	1.66 (0.18)	2.17 (0.21)	2.83 (0.27)	3.56 (0.39)	4.06 (0.50)	1.1	F	
	51-70	122	2.08 (0.16)	1.26 (0.20)	1.41 (0.18)	1.68 (0.16)	2.08 (0.17)	2.63 (0.29)	3.28 (0.53)	3.79 (0.76) ^E	1.1	F	
	>70	88	2.11 (0.12)	1.26 (0.19)	1.42 (0.19)	1.72 (0.17)	2.10 (0.17)	2.56 (0.20)	3.09 (0.27)	3.48 (0.35)	1.1	F	
	19+	471	2.24 (0.10)	1.09 (0.09)	1.30 (0.08)	1.66 (0.08)	2.15 (0.10)	2.79 (0.14)	3.56 (0.24)	4.16 (0.34)	1.1	5.2	(1.5) ^E
	Female	9-13	103	2.05 (0.18)	1.44 (0.19)	1.55 (0.18)	1.75 (0.17)	2.01 (0.19)	2.33 (0.25)	2.67 (0.36)	2.91 (0.44)	0.8	<3
14-18	142	1.78 (0.11)	0.99 (0.11)	1.11 (0.11)	1.33 (0.14)	1.65 (0.17)	2.08 (0.23)	2.56 (0.31)	2.88 (0.36)	0.9	F		
19-30	111	1.72 (0.15)	0.96 (0.18) ^E	1.10 (0.17)	1.35 (0.16)	1.67 (0.18)	2.04 (0.24)	2.40 (0.33)	2.64 (0.38)	0.9	F		
31-50	146	1.78 (0.13)	0.91 (0.17) ^E	1.04 (0.17)	1.30 (0.15)	1.64 (0.14)	2.03 (0.18)	2.43 (0.24)	2.70 (0.29)	0.9	F		
51-70	184	1.66 (0.07)	1.16 (0.07)	1.26 (0.07)	1.43 (0.09)	1.64 (0.11)	1.89 (0.13)	2.16 (0.16)	2.34 (0.18)	0.9	<3		
>70	143	1.50 (0.07)	0.90 (0.11)	0.99 (0.10)	1.19 (0.09)	1.46 (0.09)	1.82 (0.12)	2.24 (0.20)	2.54 (0.27)	0.9	F		
19+	584	1.69 (0.06)	1.00 (0.08)	1.12 (0.08)	1.35 (0.07)	1.63 (0.07)	1.98 (0.09)	2.34 (0.14)	2.59 (0.18)	0.9	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.9 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)				
Both														
	1-3	169	1.71 (0.08)	0.87 (0.11)	1.01 (0.10)	1.29 (0.09)	1.64 (0.09)	2.01 (0.12)	2.45 (0.18)	2.81 (0.25)	0.4	<3		
	4-8	281	1.91 (0.07)	1.22 (0.14)	1.35 (0.12)	1.57 (0.10)	1.84 (0.08)	2.16 (0.10)	2.53 (0.18)	2.81 (0.26)	0.5	<3		
Male														
	9-13	183	2.38 (0.14)	1.66 (0.26)	1.80 (0.24)	2.06 (0.19)	2.40 (0.16)	2.81 (0.20)	3.25 (0.34)	3.56 (0.46)	0.8	<3		
	14-18	187	2.57 (0.12)	1.51 (0.11)	1.71 (0.12)	2.08 (0.12)	2.54 (0.15)	3.10 (0.20)	3.73 (0.27)	4.20 (0.34)	1.1	<3		
	19-30	223	2.25 (0.09)	1.21 (0.20) ^E	1.41 (0.18)	1.76 (0.14)	2.19 (0.11)	2.66 (0.14)	3.13 (0.22)	3.42 (0.28)	1.1	F		
	31-50	229	2.26 (0.14)	1.47 (0.26) ^E	1.61 (0.23)	1.88 (0.19)	2.24 (0.17)	2.72 (0.23)	3.30 (0.41)	3.74 (0.59)	1.1	<3		
	51-70	197	1.91 (0.11)	0.94 (0.14)	1.13 (0.15)	1.47 (0.15)	1.89 (0.14)	2.37 (0.14)	2.83 (0.16)	3.12 (0.18)	1.1	F		
	>70	72	1.81 (0.09)	1.43 (0.15)	1.51 (0.13)	1.65 (0.11)	1.82 (0.12)	1.99 (0.16)	2.16 (0.22)	2.28 (0.27)	1.1	<3		
	19+	721	2.14 (0.07)	1.28 (0.12)	1.45 (0.10)	1.74 (0.09)	2.11 (0.08)	2.58 (0.10)	3.09 (0.16)	3.43 (0.21)	1.1	F		
Female														
	9-13	165	1.79 (0.10)	1.23 (0.18)	1.34 (0.16)	1.55 (0.13)	1.81 (0.12)	2.11 (0.16)	2.42 (0.24)	2.61 (0.30)	0.8	<3		
	14-18	206	2.08 (0.26)	1.08 (0.15)	1.23 (0.15)	1.52 (0.16)	1.91 (0.19)	2.43 (0.27)	3.04 (0.43)	3.50 (0.57)	0.9	F		
	19-30	191	1.64 (0.13)	1.07 (0.21) ^E	1.17 (0.19)	1.36 (0.17)	1.59 (0.15)	1.86 (0.19)	2.14 (0.29)	2.31 (0.36)	0.9	F		
	31-50	258	1.72 (0.09)	1.38 (0.25) ^E	1.45 (0.22)	1.58 (0.17)	1.73 (0.13)	1.90 (0.14)	2.07 (0.22)	2.18 (0.29)	0.9	F		
	51-70	249	1.62 (0.08)	1.04 (0.13)	1.15 (0.12)	1.35 (0.10)	1.59 (0.09)	1.87 (0.11)	2.17 (0.16)	2.36 (0.21)	0.9	F		
	>70	128	1.38 (0.09)	0.82 (0.09)	0.92 (0.09)	1.08 (0.08)	1.29 (0.09)	1.61 (0.14)	2.03 (0.23)	2.39 (0.33)	0.9	F		
	19+	826	1.64 (0.05)	1.01 (0.08)	1.14 (0.08)	1.35 (0.07)	1.61 (0.07)	1.93 (0.08)	2.28 (0.11)	2.52 (0.14)	0.9	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.10 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	1.79 (0.07)	1.12 (0.08)	1.25 (0.08)	1.48 (0.08)	1.77 (0.10)	2.09 (0.11)	2.42 (0.13)	2.63 (0.15)	0.4	0.0	(0.0)
	4-8	321	1.98 (0.08)	1.30 (0.09)	1.42 (0.09)	1.63 (0.07)	1.92 (0.07)	2.29 (0.11)	2.76 (0.21)	3.13 (0.33)	0.5	0.0	(0.0)
Male													
	9-13	226	2.32 (0.10)	1.44 (0.14)	1.61 (0.13)	1.91 (0.11)	2.30 (0.11)	2.78 (0.15)	3.36 (0.24)	3.80 (0.32)	0.8	<3	
	14-18	262	2.83 (0.21)	1.52 (0.23)	1.75 (0.21)	2.19 (0.19)	2.76 (0.19)	3.45 (0.25)	4.22 (0.38)	4.76 (0.49)	1.1	<3	
	19-30	197	2.51 (0.13)	1.59 (0.24)	1.75 (0.21)	2.04 (0.17)	2.41 (0.15)	2.84 (0.20)	3.29 (0.31)	3.59 (0.39)	1.1	<3	
	31-50	282	2.60 (0.16)	1.48 (0.20)	1.68 (0.19)	2.05 (0.17)	2.53 (0.17)	3.09 (0.22)	3.70 (0.36)	4.15 (0.50)	1.1	<3	
	51-70	234	2.15 (0.10)	1.25 (0.08)	1.39 (0.08)	1.66 (0.09)	2.02 (0.11)	2.44 (0.14)	2.92 (0.20)	3.27 (0.25)	1.1	F	
	>70	119	1.98 (0.15)	1.19 (0.22) ^E	1.33 (0.20)	1.61 (0.18)	1.95 (0.18)	2.32 (0.22)	2.66 (0.28)	2.86 (0.32)	1.1	F	
	19+	832	2.39 (0.08)	1.32 (0.08)	1.49 (0.07)	1.83 (0.07)	2.29 (0.08)	2.86 (0.11)	3.46 (0.17)	3.87 (0.23)	1.1	<3	
Female													
	9-13	226	2.00 (0.13)	1.09 (0.14)	1.25 (0.14)	1.54 (0.13)	1.91 (0.14)	2.36 (0.18)	2.91 (0.25)	3.31 (0.32)	0.8	<3	
	14-18	242	1.91 (0.12)	0.89 (0.10)	1.02 (0.11)	1.31 (0.13)	1.77 (0.14)	2.36 (0.20)	3.08 (0.30)	3.64 (0.40)	0.9	F	
	19-30	208	1.77 (0.10)	1.22 (0.21) ^E	1.35 (0.19)	1.58 (0.16)	1.85 (0.14)	2.13 (0.15)	2.39 (0.20)	2.56 (0.25)	0.9	F	
	31-50	263	1.78 (0.08)	1.15 (0.13)	1.27 (0.11)	1.48 (0.09)	1.73 (0.09)	2.00 (0.13)	2.27 (0.20)	2.45 (0.26)	0.9	<3	
	51-70	322	1.54 (0.07)	0.85 (0.09)	0.98 (0.08)	1.21 (0.07)	1.49 (0.07)	1.85 (0.10)	2.26 (0.15)	2.55 (0.19)	0.9	F	
	>70	198	1.47 (0.06)	0.84 (0.11)	0.96 (0.10)	1.18 (0.09)	1.42 (0.08)	1.70 (0.09)	2.01 (0.13)	2.22 (0.17)	0.9	F	
	19+	991	1.67 (0.04)	0.96 (0.05)	1.09 (0.05)	1.33 (0.05)	1.63 (0.05)	1.99 (0.06)	2.35 (0.09)	2.59 (0.12)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.11 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		(SE)	
Both														
	1-3	348	1.84	(0.07)	1.20 (0.10)	1.32 (0.10)	1.54 (0.09)	1.81 (0.08)	2.13 (0.10)	2.47 (0.15)	2.71 (0.19)	0.4	0.0	(0.0)
	4-8	554	2.05	(0.06)	1.39 (0.08)	1.53 (0.08)	1.77 (0.07)	2.07 (0.07)	2.40 (0.09)	2.73 (0.12)	2.96 (0.16)	0.5	0.0	(0.0)
Male														
	9-13	409	2.29	(0.08)	1.47 (0.12)	1.63 (0.11)	1.90 (0.09)	2.22 (0.09)	2.58 (0.11)	2.98 (0.16)	3.26 (0.21)	0.8	0.0	(0.0)
	14-18	414	2.51	(0.12)	1.44 (0.12)	1.63 (0.12)	2.01 (0.13)	2.51 (0.14)	3.10 (0.19)	3.74 (0.27)	4.19 (0.34)	1.1	<3	
	19-30	311	2.56	(0.13)	1.33 (0.13)	1.53 (0.13)	1.94 (0.13)	2.50 (0.15)	3.19 (0.20)	3.99 (0.30)	4.60 (0.40)	1.1	F	
	31-50	489	2.20	(0.10)	1.31 (0.09)	1.46 (0.09)	1.73 (0.09)	2.08 (0.09)	2.52 (0.12)	3.00 (0.17)	3.34 (0.22)	1.1	F	
	51-70	575	1.94	(0.05)	1.28 (0.08)	1.40 (0.07)	1.62 (0.07)	1.89 (0.06)	2.19 (0.07)	2.47 (0.09)	2.66 (0.11)	1.1	F	
	>70	239	1.90	(0.08)	1.27 (0.09)	1.37 (0.09)	1.58 (0.10)	1.83 (0.10)	2.12 (0.12)	2.42 (0.16)	2.62 (0.21)	1.1	F	
	19+	1614	2.17	(0.05)	1.26 (0.05)	1.41 (0.05)	1.69 (0.05)	2.08 (0.05)	2.54 (0.07)	3.05 (0.10)	3.43 (0.13)	1.1	1.9	(0.6) ^E
Female														
	9-13	355	1.99	(0.09)	1.29 (0.13)	1.42 (0.12)	1.68 (0.12)	2.00 (0.12)	2.37 (0.14)	2.75 (0.19)	3.00 (0.23)	0.8	<3	
	14-18	410	1.79	(0.08)	0.90 (0.12)	1.10 (0.11)	1.43 (0.08)	1.77 (0.08)	2.14 (0.10)	2.58 (0.15)	2.91 (0.18)	0.9	F	
	19-30	384	1.60	(0.07)	0.98 (0.08)	1.10 (0.08)	1.31 (0.07)	1.58 (0.08)	1.90 (0.10)	2.26 (0.13)	2.50 (0.16)	0.9	F	
	31-50	585	1.68	(0.05)	0.84 (0.07)	1.00 (0.06)	1.30 (0.06)	1.67 (0.07)	2.08 (0.09)	2.52 (0.12)	2.84 (0.17)	0.9	6.6	(2.0) ^E
	51-70	711	1.67	(0.07)	0.93 (0.06)	1.05 (0.06)	1.28 (0.06)	1.56 (0.06)	1.95 (0.09)	2.43 (0.17)	2.83 (0.27)	0.9	F	
	>70	342	1.64	(0.08)	0.93 (0.08)	1.04 (0.08)	1.25 (0.08)	1.55 (0.10)	1.96 (0.14)	2.43 (0.19)	2.76 (0.23)	0.9	F	
	19+	2022	1.66	(0.03)	0.92 (0.03)	1.06 (0.03)	1.30 (0.04)	1.61 (0.04)	2.01 (0.05)	2.44 (0.08)	2.76 (0.11)	0.9	4.3	(0.9) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.12 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	622	1.75	(0.06)	0.92 (0.07)	1.07 (0.07)	1.36 (0.06)	1.70 (0.07)	2.11 (0.08)	2.53 (0.12)	2.86 (0.17)	0.4	<3	
	4-8	919	1.92	(0.05)	1.26 (0.09)	1.37 (0.08)	1.58 (0.06)	1.85 (0.05)	2.17 (0.07)	2.52 (0.12)	2.78 (0.17)	0.5	0.0	(0.0)
Male														
	9-13	579	2.36	(0.09)	1.54 (0.11)	1.69 (0.10)	1.97 (0.10)	2.34 (0.10)	2.79 (0.14)	3.28 (0.21)	3.63 (0.27)	0.8	<3	
	14-18	634	2.66	(0.09)	1.50 (0.12)	1.72 (0.11)	2.12 (0.11)	2.61 (0.11)	3.21 (0.14)	3.90 (0.22)	4.41 (0.31)	1.1	<3	
	19-30	578	2.33	(0.07)	1.15 (0.10)	1.35 (0.10)	1.74 (0.09)	2.22 (0.08)	2.77 (0.10)	3.34 (0.14)	3.76 (0.19)	1.1	F	
	31-50	693	2.23	(0.10)	1.37 (0.15)	1.52 (0.14)	1.80 (0.13)	2.18 (0.13)	2.69 (0.15)	3.27 (0.24)	3.70 (0.33)	1.1	F	
	51-70	596	1.94	(0.08)	1.00 (0.12)	1.18 (0.11)	1.51 (0.09)	1.91 (0.09)	2.41 (0.11)	2.95 (0.16)	3.32 (0.22)	1.1	F	
	>70	296	1.90	(0.07)	1.11 (0.10)	1.26 (0.09)	1.52 (0.08)	1.83 (0.08)	2.22 (0.10)	2.66 (0.14)	2.99 (0.18)	1.1	F	
	19+	2163	2.15	(0.05)	1.17 (0.05)	1.35 (0.05)	1.67 (0.05)	2.09 (0.06)	2.62 (0.07)	3.21 (0.10)	3.63 (0.12)	1.1	3.7	(0.9) ^E
Female														
	9-13	533	1.89	(0.07)	1.26 (0.10)	1.38 (0.09)	1.60 (0.08)	1.85 (0.08)	2.15 (0.11)	2.49 (0.15)	2.72 (0.20)	0.8	<3	
	14-18	638	1.98	(0.16)	1.09 (0.12)	1.22 (0.12)	1.49 (0.12)	1.86 (0.13)	2.34 (0.26)	2.89 (0.45)	3.31 (0.52)	0.9	F	
	19-30	499	1.71	(0.09)	0.94 (0.13)	1.08 (0.12)	1.33 (0.10)	1.66 (0.10)	2.03 (0.14)	2.41 (0.20)	2.66 (0.24)	0.9	F	
	31-50	716	1.72	(0.06)	1.09 (0.14)	1.21 (0.12)	1.42 (0.10)	1.67 (0.09)	1.98 (0.09)	2.30 (0.13)	2.52 (0.16)	0.9	F	
	51-70	745	1.66	(0.06)	1.12 (0.11)	1.22 (0.10)	1.41 (0.08)	1.64 (0.06)	1.91 (0.08)	2.20 (0.13)	2.41 (0.17)	0.9	<3	
	>70	510	1.42	(0.05)	0.82 (0.06)	0.93 (0.05)	1.12 (0.05)	1.36 (0.05)	1.69 (0.08)	2.10 (0.13)	2.42 (0.19)	0.9	8.3	(2.7) ^E
	19+	2470	1.67	(0.04)	0.95 (0.05)	1.08 (0.04)	1.32 (0.04)	1.62 (0.04)	1.97 (0.05)	2.36 (0.07)	2.64 (0.09)	0.9	3.6	(1.0) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 22.13 Riboflavin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	1.82 (0.03)	1.09 (0.05)	1.24 (0.04)	1.50 (0.04)	1.81 (0.04)	2.18 (0.05)	2.60 (0.07)	2.91 (0.10)	0.4	<3	
	4-8	3235	2.03 (0.05)	1.22 (0.04)	1.36 (0.04)	1.62 (0.03)	1.95 (0.04)	2.32 (0.05)	2.84 (0.13)	3.33 (0.24)	0.5	0.0	(0.0)
Male													
	9-13	2080	2.41 (0.05)	1.46 (0.06)	1.63 (0.06)	1.94 (0.05)	2.34 (0.05)	2.85 (0.07)	3.43 (0.10)	3.85 (0.15)	0.8	<3	
	14-18	2288	2.65 (0.05)	1.50 (0.06)	1.70 (0.06)	2.10 (0.06)	2.63 (0.06)	3.27 (0.08)	3.97 (0.11)	4.48 (0.15)	1.1	<3	
	19-30	1804	2.41 (0.06)	1.33 (0.06)	1.51 (0.06)	1.85 (0.06)	2.29 (0.07)	2.84 (0.09)	3.46 (0.13)	3.88 (0.17)	1.1	<3	
	31-50	2596	2.24 (0.05)	1.29 (0.05)	1.46 (0.05)	1.76 (0.05)	2.17 (0.05)	2.68 (0.06)	3.25 (0.09)	3.66 (0.12)	1.1	1.7	(0.6) ^E
	51-70	2550	2.01 (0.03)	1.16 (0.04)	1.30 (0.04)	1.57 (0.04)	1.93 (0.04)	2.36 (0.05)	2.83 (0.07)	3.14 (0.09)	1.1	3.6	(0.9) ^E
	>70	1520	1.86 (0.05)	1.07 (0.05)	1.21 (0.05)	1.45 (0.05)	1.78 (0.06)	2.18 (0.07)	2.62 (0.10)	2.92 (0.12)	1.1	5.9	(1.4) ^E
	19+	8470	2.18 (0.03)	1.22 (0.03)	1.38 (0.03)	1.69 (0.03)	2.09 (0.03)	2.59 (0.04)	3.16 (0.05)	3.57 (0.07)	1.1	2.7	(0.4)
Female													
	9-13	1980	1.97 (0.04)	1.12 (0.04)	1.26 (0.04)	1.53 (0.04)	1.87 (0.04)	2.29 (0.05)	2.75 (0.08)	3.07 (0.10)	0.8	<3	
	14-18	2256	1.90 (0.04)	1.03 (0.04)	1.18 (0.04)	1.46 (0.04)	1.85 (0.05)	2.33 (0.06)	2.85 (0.10)	3.22 (0.13)	0.9	2.4	(0.6) ^E
	19-30	1854	1.72 (0.04)	1.02 (0.05)	1.14 (0.05)	1.38 (0.04)	1.68 (0.04)	2.02 (0.05)	2.37 (0.07)	2.60 (0.09)	0.9	F	
	31-50	2686	1.74 (0.03)	0.99 (0.04)	1.13 (0.03)	1.38 (0.03)	1.70 (0.04)	2.07 (0.05)	2.48 (0.07)	2.76 (0.08)	0.9	2.8	(0.7) ^E
	51-70	3200	1.65 (0.02)	0.98 (0.03)	1.10 (0.03)	1.32 (0.03)	1.59 (0.03)	1.93 (0.03)	2.31 (0.05)	2.58 (0.07)	0.9	3.0	(0.8) ^E
	>70	2610	1.53 (0.04)	0.92 (0.04)	1.03 (0.04)	1.23 (0.04)	1.49 (0.04)	1.81 (0.05)	2.16 (0.06)	2.40 (0.07)	0.9	4.4	(1.3) ^E
	19+	10350	1.68 (0.02)	0.99 (0.02)	1.12 (0.02)	1.35 (0.02)	1.64 (0.02)	1.99 (0.02)	2.37 (0.03)	2.64 (0.04)	0.9	2.8	(0.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

23. Thiamin (mg/d): Usual intakes from food

Table 23.1 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	1.11 (0.07)	0.73 (0.07)	0.79 (0.07)	0.90 (0.08)	1.04 (0.09)	1.19 (0.10)	1.35 (0.11)	1.45 (0.12)	0.4	<3	
	4-8	127	1.77 (0.10)	1.19 (0.23) ^E	1.31 (0.21)	1.52 (0.18)	1.79 (0.16)	2.09 (0.17)	2.40 (0.23)	2.59 (0.29)	0.5	<3	
Male	9-13	111	1.93 (0.12)	1.42 (0.18)	1.51 (0.17)	1.67 (0.15)	1.86 (0.15)	2.09 (0.19)	2.31 (0.25)	2.46 (0.30)	0.7	0.0 (0.0)	
	14-18	107	2.29 (0.24)	1.42 (0.30) ^E	1.59 (0.28) ^E	1.96 (0.26)	2.44 (0.28)	2.97 (0.37)	3.54 (0.52)	3.93 (0.65)	1.0	<3	
	19-30	77	1.90 (0.12)	1.32 (0.21)	1.43 (0.19)	1.64 (0.15)	1.89 (0.13)	2.17 (0.17)	2.43 (0.24)	2.59 (0.29)	1.0	F	
	31-50	145	1.69 (0.11)	1.09 (0.18)	1.19 (0.16)	1.38 (0.14)	1.62 (0.13)	1.91 (0.16)	2.22 (0.30)	2.44 (0.47) ^E	1.0	F	
	51-70	182	1.68 (0.07)	1.11 (0.08)	1.22 (0.08)	1.40 (0.08)	1.60 (0.08)	1.83 (0.09)	2.07 (0.11)	2.23 (0.12)	1.0	F	
	>70	63	2.09 (0.24)	1.39 (0.32) ^E	1.52 (0.32) ^E	1.76 (0.32) ^E	2.09 (0.34)	2.48 (0.39)	2.90 (0.50) ^E	3.18 (0.60) ^E	1.0	F	
	19+	467	1.76 (0.06)	1.11 (0.09)	1.23 (0.08)	1.44 (0.07)	1.71 (0.07)	2.03 (0.08)	2.38 (0.11)	2.62 (0.13)	1.0	F	
	Female	9-13	96	1.49 (0.09)	0.88 (0.13)	0.99 (0.12)	1.18 (0.11)	1.44 (0.11)	1.76 (0.14)	2.14 (0.22)	2.43 (0.29)	0.7	F
14-18	105	1.60 (0.13)	0.94 (0.19) ^E	1.05 (0.19) ^E	1.26 (0.18)	1.57 (0.17)	1.95 (0.21)	2.31 (0.28)	2.54 (0.33)	0.9	F		
19-30	91	1.23 (0.07)	0.88 (0.12)	0.94 (0.11)	1.04 (0.09)	1.17 (0.10)	1.30 (0.12)	1.43 (0.16)	1.51 (0.19)	0.9	F		
31-50	167	1.53 (0.12)	0.87 (0.16) ^E	1.01 (0.15)	1.27 (0.14)	1.60 (0.15)	1.98 (0.19)	2.35 (0.23)	2.58 (0.27)	0.9	F		
51-70	198	1.48 (0.08)	0.85 (0.17) ^E	0.95 (0.15)	1.16 (0.12)	1.45 (0.10)	1.76 (0.14)	2.04 (0.20)	2.20 (0.24)	0.9	F		
>70	74	1.62 (0.13)	1.07 (0.20) ^E	1.17 (0.19)	1.36 (0.17)	1.59 (0.17)	1.85 (0.19)	2.11 (0.25)	2.28 (0.30)	0.9	F		
19+	530	1.47 (0.06)	0.84 (0.08)	0.96 (0.07)	1.18 (0.07)	1.46 (0.07)	1.80 (0.09)	2.13 (0.11)	2.33 (0.13)	0.9	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.2 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	58	1.19 (0.07)	0.81 (0.12)	0.89 (0.12)	1.04 (0.10)	1.21 (0.10)	1.40 (0.11)	1.58 (0.13)	1.69 (0.16)	0.4	<3	
	4-8	110	1.51 (0.07)	1.03 (0.14)	1.13 (0.12)	1.29 (0.09)	1.48 (0.08)	1.68 (0.11)	1.86 (0.16)	1.97 (0.20)	0.5	<3	
Male													
	9-13	95	1.76 (0.08)	1.36 (0.14)	1.42 (0.13)	1.52 (0.12)	1.65 (0.13)	1.80 (0.15)	1.94 (0.20)	2.04 (0.24)	0.7	<3	
	14-18	87	2.16 (0.16)	1.47 (0.31) ^E	1.64 (0.29) ^E	1.94 (0.25)	2.32 (0.23)	2.74 (0.27)	3.17 (0.37)	3.45 (0.46)	1.0	F	
	19-30	70	2.41 (0.16)	1.66 (0.31) ^E	1.83 (0.28)	2.12 (0.23)	2.47 (0.20)	2.85 (0.24)	3.22 (0.34)	3.45 (0.41)	1.0	<3	
	31-50	109	1.83 (0.16)	1.06 (0.17)	1.17 (0.16)	1.37 (0.15)	1.64 (0.17)	1.98 (0.24)	2.37 (0.35)	2.65 (0.42)	1.0	F	
	51-70	128	1.67 (0.08)	1.16 (0.14)	1.25 (0.12)	1.41 (0.10)	1.60 (0.09)	1.82 (0.13)	2.07 (0.20)	2.26 (0.26)	1.0	F	
	>70	65	1.57 (0.11)	0.99 (0.14)	1.09 (0.13)	1.27 (0.12)	1.52 (0.12)	1.83 (0.16)	2.15 (0.22)	2.37 (0.27)	1.0	F	
	19+	372	1.88 (0.08)	1.11 (0.08)	1.23 (0.08)	1.46 (0.08)	1.76 (0.10)	2.13 (0.13)	2.55 (0.19)	2.85 (0.23)	1.0	F	
Female													
	9-13	75	1.47 (0.10)	1.12 (0.17)	1.21 (0.16)	1.36 (0.15)	1.55 (0.15)	1.75 (0.18)	1.95 (0.22)	2.07 (0.25)	0.7	<3	
	14-18	81	1.45 (0.11)	0.96 (0.19) ^E	1.05 (0.17)	1.22 (0.15)	1.41 (0.13)	1.62 (0.14)	1.81 (0.17)	1.93 (0.19)	0.9	F	
	19-30	101	1.50 (0.09)	1.11 (0.14)	1.21 (0.12)	1.38 (0.11)	1.58 (0.12)	1.79 (0.16)	1.99 (0.21)	2.12 (0.24)	0.9	<3	
	31-50	116	1.44 (0.08)	0.83 (0.09)	0.95 (0.09)	1.16 (0.09)	1.41 (0.10)	1.68 (0.13)	1.94 (0.17)	2.12 (0.20)	0.9	F	
	51-70	146	1.58 (0.09)	0.99 (0.08)	1.10 (0.08)	1.28 (0.09)	1.53 (0.12)	1.82 (0.14)	2.12 (0.17)	2.32 (0.18)	0.9	F	
	>70	94	1.27 (0.10)	0.64 (0.15) ^E	0.74 (0.14) ^E	0.92 (0.13)	1.17 (0.13)	1.47 (0.15)	1.79 (0.19)	2.01 (0.24)	0.9	F	
	19+	457	1.47 (0.05)	0.94 (0.06)	1.04 (0.06)	1.23 (0.06)	1.46 (0.06)	1.71 (0.08)	1.97 (0.10)	2.14 (0.12)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.3 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	112	1.20 (0.05)	0.89 (0.09)	0.95 (0.08)	1.05 (0.07)	1.17 (0.06)	1.31 (0.08)	1.45 (0.11)	1.55 (0.14)	0.4	0.0 (0.0)	
	4-8	177	1.55 (0.09)	1.03 (0.22) ^E	1.14 (0.19) ^E	1.33 (0.14)	1.57 (0.11)	1.83 (0.14)	2.09 (0.22)	2.27 (0.28)	0.5	<3	
Male													
	9-13	111	1.87 (0.08)	1.12 (0.15)	1.23 (0.13)	1.45 (0.11)	1.75 (0.10)	2.10 (0.14)	2.44 (0.21)	2.66 (0.26)	0.7	<3	
	14-18	113	1.84 (0.17)	1.11 (0.19) ^E	1.23 (0.21) ^E	1.46 (0.22)	1.76 (0.24)	2.10 (0.26)	2.45 (0.28)	2.69 (0.29)	1.0	F	
	19-30	91	2.23 (0.20)	1.37 (0.16)	1.51 (0.16)	1.78 (0.18)	2.12 (0.21)	2.52 (0.27)	2.94 (0.35)	3.21 (0.41)	1.0	<3	
	31-50	101	1.90 (0.14)	1.04 (0.12)	1.17 (0.12)	1.42 (0.12)	1.77 (0.15)	2.23 (0.20)	2.76 (0.27)	3.14 (0.34)	1.0	F	
	51-70	134	1.91 (0.14)	1.34 (0.14)	1.45 (0.15)	1.65 (0.16)	1.89 (0.18)	2.15 (0.21)	2.40 (0.24)	2.56 (0.26)	1.0	<3	
	>70	56	1.72 (0.13)	1.21 (0.16)	1.30 (0.17)	1.47 (0.18)	1.68 (0.19)	1.91 (0.18)	2.15 (0.17)	2.30 (0.17)	1.0	<3	
	19+	382	1.95 (0.09)	1.33 (0.14)	1.44 (0.12)	1.64 (0.11)	1.89 (0.10)	2.19 (0.14)	2.50 (0.20)	2.71 (0.25)	1.0	<3	
Female													
	9-13	105	1.49 (0.11)	1.04 (0.09)	1.13 (0.10)	1.29 (0.11)	1.49 (0.14)	1.74 (0.18)	2.00 (0.23)	2.18 (0.29)	0.7	<3	
	14-18	120	1.43 (0.15)	0.75 (0.24) ^E	0.88 (0.22) ^E	1.13 (0.19) ^E	1.44 (0.18)	1.79 (0.21)	2.19 (0.31)	2.49 (0.40)	0.9	F	
	19-30	91	1.47 (0.11)	1.05 (0.12)	1.12 (0.12)	1.25 (0.14)	1.42 (0.15)	1.61 (0.18)	1.82 (0.20)	1.97 (0.23)	0.9	F	
	31-50	159	1.49 (0.07)	0.75 (0.10)	0.89 (0.10)	1.15 (0.09)	1.46 (0.09)	1.76 (0.11)	2.07 (0.16)	2.31 (0.20)	0.9	F	
	51-70	174	1.42 (0.08)	0.91 (0.15)	1.01 (0.13)	1.18 (0.11)	1.39 (0.09)	1.62 (0.11)	1.86 (0.17)	2.00 (0.22)	0.9	F	
	>70	80	1.44 (0.13)	0.85 (0.20) ^E	0.95 (0.19) ^E	1.14 (0.18)	1.44 (0.17)	1.82 (0.23)	2.22 (0.33)	2.50 (0.42) ^E	0.9	F	
	19+	504	1.46 (0.04)	0.87 (0.08)	0.98 (0.08)	1.19 (0.07)	1.44 (0.06)	1.72 (0.07)	2.01 (0.10)	2.20 (0.12)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.4 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	99	1.25 (0.08)	0.97 (0.12)	1.03 (0.11)	1.13 (0.09)	1.26 (0.09)	1.41 (0.12)	1.57 (0.19)	1.68 (0.23)	0.4	<3	
	4-8	140	1.65 (0.09)	1.01 (0.14)	1.13 (0.13)	1.36 (0.11)	1.64 (0.10)	1.95 (0.13)	2.27 (0.19)	2.51 (0.25)	0.5	<3	
Male													
	9-13	92	1.87 (0.18)	1.19 (0.17)	1.29 (0.17)	1.48 (0.18)	1.75 (0.20)	2.10 (0.25)	2.50 (0.31)	2.79 (0.36)	0.7	<3	
	14-18	107	2.27 (0.18)	1.38 (0.28) ^E	1.57 (0.24)	1.89 (0.20)	2.28 (0.19)	2.76 (0.28)	3.35 (0.45)	3.80 (0.60)	1.0	<3	
	19-30	73	2.95 (0.52) ^E	1.35 (0.31) ^E	1.58 (0.31) ^E	2.06 (0.34)	2.76 (0.45)	3.70 (0.68) ^E	4.82 (1.04) ^E	5.65 (1.34) ^E	1.0	F	
	31-50	134	1.92 (0.14)	1.01 (0.25) ^E	1.14 (0.23) ^E	1.40 (0.20)	1.74 (0.16)	2.17 (0.19)	2.64 (0.32)	2.99 (0.45)	1.0	F	
	51-70	131	1.76 (0.10)	1.34 (0.19)	1.42 (0.17)	1.56 (0.13)	1.73 (0.12)	1.92 (0.15)	2.11 (0.23)	2.23 (0.30)	1.0	<3	
	>70	55	1.77 (0.15)	1.20 (0.19)	1.29 (0.18)	1.47 (0.16)	1.70 (0.17)	2.00 (0.22)	2.33 (0.31)	2.53 (0.38)	1.0	F	
	19+	393	2.08 (0.12)	1.05 (0.12)	1.20 (0.11)	1.49 (0.11)	1.91 (0.12)	2.50 (0.18)	3.27 (0.32)	3.87 (0.48)	1.0	F	
Female													
	9-13	79	1.83 (0.16)	1.32 (0.22) ^E	1.43 (0.20)	1.62 (0.18)	1.85 (0.18)	2.13 (0.23)	2.44 (0.33)	2.66 (0.43)	0.7	<3	
	14-18	104	1.59 (0.12)	1.06 (0.20) ^E	1.16 (0.18)	1.34 (0.15)	1.55 (0.14)	1.78 (0.17)	2.01 (0.23)	2.16 (0.29)	0.9	F	
	19-30	101	1.57 (0.16)	1.04 (0.25) ^E	1.16 (0.24) ^E	1.38 (0.22)	1.67 (0.21)	2.00 (0.25)	2.34 (0.32)	2.56 (0.39)	0.9	F	
	31-50	143	1.43 (0.10)	0.77 (0.16) ^E	0.88 (0.15)	1.10 (0.12)	1.38 (0.11)	1.67 (0.15)	1.97 (0.22)	2.18 (0.28)	0.9	F	
	51-70	193	1.50 (0.09)	0.96 (0.16)	1.05 (0.14)	1.21 (0.12)	1.41 (0.11)	1.64 (0.12)	1.90 (0.18)	2.08 (0.24)	0.9	F	
	>70	94	1.53 (0.10)	0.89 (0.11)	1.00 (0.12)	1.21 (0.12)	1.47 (0.13)	1.78 (0.15)	2.11 (0.17)	2.32 (0.19)	0.9	F	
	19+	531	1.49 (0.06)	0.89 (0.09)	1.00 (0.08)	1.20 (0.07)	1.45 (0.07)	1.74 (0.08)	2.04 (0.11)	2.24 (0.14)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.5 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both													
	1-3	311	1.26	(0.06)	0.80 (0.11)	0.89 (0.10)	1.06 (0.08)	1.25 (0.07)	1.47 (0.09)	1.69 (0.13)	1.84 (0.16)	0.4	<3
	4-8	485	1.76	(0.06)	1.21 (0.14)	1.33 (0.13)	1.54 (0.10)	1.79 (0.08)	2.07 (0.11)	2.36 (0.17)	2.56 (0.22)	0.5	<3
Male													
	9-13	277	2.31	(0.12)	1.51 (0.20)	1.67 (0.19)	1.97 (0.17)	2.38 (0.17)	2.86 (0.21)	3.41 (0.30)	3.79 (0.38)	0.7	<3
	14-18	339	2.66	(0.12)	1.48 (0.15)	1.70 (0.14)	2.13 (0.14)	2.70 (0.16)	3.40 (0.20)	4.15 (0.27)	4.66 (0.34)	1.0	<3
	19-30	237	2.14	(0.10)	1.38 (0.18)	1.50 (0.16)	1.74 (0.14)	2.05 (0.13)	2.41 (0.16)	2.77 (0.24)	3.01 (0.30)	1.0	<3
	31-50	423	2.10	(0.11)	1.48 (0.22)	1.60 (0.20)	1.83 (0.17)	2.10 (0.13)	2.39 (0.24)	2.66 (0.46) ^E	2.84 (0.63) ^E	1.0	^F
	51-70	387	1.91	(0.06)	1.26 (0.11)	1.39 (0.10)	1.60 (0.08)	1.87 (0.07)	2.17 (0.09)	2.49 (0.14)	2.70 (0.18)	1.0	<3
	>70	132	1.54	(0.10)	0.85 (0.17) ^E	0.96 (0.16) ^E	1.17 (0.15)	1.43 (0.15)	1.74 (0.19)	2.05 (0.27)	2.25 (0.33)	1.0	^F
	19+	1179	2.01	(0.05)	1.23 (0.08)	1.36 (0.07)	1.62 (0.07)	1.94 (0.07)	2.33 (0.08)	2.72 (0.11)	2.99 (0.14)	1.0	<3
Female													
	9-13	281	1.76	(0.07)	1.00 (0.10)	1.13 (0.10)	1.38 (0.09)	1.69 (0.09)	2.07 (0.11)	2.48 (0.16)	2.77 (0.22)	0.7	<3
	14-18	321	1.68	(0.07)	1.13 (0.12)	1.25 (0.11)	1.44 (0.09)	1.69 (0.09)	1.98 (0.12)	2.29 (0.17)	2.49 (0.22)	0.9	<3
	19-30	249	1.67	(0.09)	1.14 (0.16)	1.25 (0.14)	1.43 (0.11)	1.64 (0.10)	1.87 (0.13)	2.11 (0.20)	2.26 (0.25)	0.9	^F
	31-50	364	1.59	(0.06)	1.09 (0.13)	1.19 (0.12)	1.37 (0.10)	1.58 (0.09)	1.83 (0.10)	2.09 (0.14)	2.26 (0.17)	0.9	^F
	51-70	467	1.53	(0.05)	1.03 (0.11)	1.12 (0.10)	1.30 (0.07)	1.51 (0.06)	1.75 (0.08)	2.00 (0.13)	2.16 (0.17)	0.9	^F
	>70	215	1.37	(0.10)	0.82 (0.12)	0.93 (0.12)	1.12 (0.12)	1.35 (0.13)	1.63 (0.16)	1.93 (0.19)	2.13 (0.22)	0.9	^F
19+	1295	1.56	(0.04)	1.00 (0.06)	1.11 (0.05)	1.30 (0.05)	1.54 (0.05)	1.82 (0.06)	2.12 (0.08)	2.31 (0.10)	0.9	^F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.6 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both													
	1-3	644	1.22	(0.04)	0.73 (0.07)	0.82 (0.07)	0.98 (0.05)	1.19 (0.04)	1.46 (0.06)	1.75 (0.11)	1.95 (0.15)	0.4	<3
	4-8	956	1.56	(0.03)	0.96 (0.07)	1.07 (0.06)	1.27 (0.05)	1.53 (0.04)	1.84 (0.06)	2.17 (0.10)	2.40 (0.14)	0.5	<3
Male													
	9-13	589	2.03	(0.08)	1.01 (0.07)	1.16 (0.07)	1.44 (0.07)	1.85 (0.08)	2.40 (0.11)	3.09 (0.20)	3.62 (0.29)	0.7	<3
	14-18	639	2.26	(0.07)	1.51 (0.22)	1.66 (0.19)	1.93 (0.15)	2.27 (0.10)	2.67 (0.12)	3.10 (0.24)	3.39 (0.33)	1.0	<3
	19-30	481	2.02	(0.08)	1.11 (0.23) ^E	1.27 (0.20)	1.55 (0.15)	1.93 (0.10)	2.41 (0.14)	2.94 (0.29)	3.31 (0.41)	1.0	F
	31-50	709	1.91	(0.07)	1.15 (0.21) ^E	1.27 (0.19)	1.51 (0.15)	1.83 (0.09)	2.24 (0.12)	2.69 (0.24)	3.01 (0.35)	1.0	F
	51-70	758	1.85	(0.05)	0.95 (0.09)	1.10 (0.08)	1.36 (0.07)	1.73 (0.05)	2.19 (0.07)	2.67 (0.13)	3.01 (0.19)	1.0	F
	>70	734	1.67	(0.04)	0.91 (0.08)	1.04 (0.07)	1.28 (0.06)	1.61 (0.05)	2.00 (0.06)	2.41 (0.10)	2.70 (0.13)	1.0	F
	19+	2682	1.90	(0.04)	1.04 (0.07)	1.19 (0.07)	1.45 (0.06)	1.82 (0.05)	2.27 (0.06)	2.75 (0.10)	3.08 (0.15)	1.0	F
Female													
	9-13	585	1.59	(0.04)	1.05 (0.12)	1.15 (0.10)	1.33 (0.07)	1.56 (0.05)	1.82 (0.07)	2.07 (0.13)	2.23 (0.17)	0.7	<3
	14-18	645	1.69	(0.07)	0.98 (0.12)	1.10 (0.11)	1.32 (0.09)	1.61 (0.07)	2.00 (0.10)	2.45 (0.20)	2.78 (0.29)	0.9	F
	19-30	514	1.38	(0.05)	0.70 (0.08)	0.79 (0.08)	0.99 (0.07)	1.29 (0.06)	1.64 (0.07)	1.98 (0.12)	2.20 (0.16)	0.9	17.4 (5.3) ^E
	31-50	758	1.41	(0.04)	0.80 (0.07)	0.91 (0.06)	1.12 (0.05)	1.38 (0.04)	1.68 (0.06)	2.00 (0.09)	2.23 (0.12)	0.9	F
	51-70	955	1.42	(0.04)	0.78 (0.06)	0.88 (0.06)	1.08 (0.05)	1.34 (0.05)	1.66 (0.06)	2.00 (0.09)	2.22 (0.12)	0.9	11.1 (3.3) ^E
	>70	1345	1.41	(0.03)	0.85 (0.06)	0.94 (0.05)	1.11 (0.04)	1.34 (0.04)	1.60 (0.05)	1.88 (0.08)	2.07 (0.11)	0.9	F
	19+	3572	1.41	(0.02)	0.78 (0.03)	0.89 (0.03)	1.09 (0.03)	1.34 (0.02)	1.66 (0.03)	1.98 (0.05)	2.21 (0.07)	0.9	10.8 (1.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.7 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	324	1.18	(0.05)	0.71 (0.09)	0.80 (0.08)	0.95 (0.06)	1.16 (0.06)	1.40 (0.08)	1.66 (0.11)	1.82 (0.14)	0.4	<3	
	4-8	425	1.50	(0.06)	0.97 (0.05)	1.06 (0.05)	1.23 (0.06)	1.46 (0.07)	1.72 (0.08)	2.00 (0.11)	2.19 (0.14)	0.5	0.0	(0.0)
Male														
	9-13	274	1.95	(0.11)	1.44 (0.22)	1.53 (0.20)	1.71 (0.15)	1.93 (0.13)	2.17 (0.16)	2.42 (0.24)	2.57 (0.30)	0.7	<3	
	14-18	297	2.35	(0.12)	1.20 (0.19)	1.40 (0.18)	1.78 (0.16)	2.30 (0.15)	2.92 (0.18)	3.59 (0.26)	4.04 (0.33)	1.0	F	
	19-30	249	2.38	(0.19)	1.49 (0.29) ^E	1.63 (0.25)	1.90 (0.19)	2.25 (0.15)	2.65 (0.22)	3.07 (0.37)	3.34 (0.48)	1.0	<3	
	31-50	309	1.78	(0.12)	1.21 (0.18)	1.32 (0.16)	1.51 (0.13)	1.75 (0.13)	2.03 (0.17)	2.31 (0.26)	2.49 (0.32)	1.0	F	
	51-70	277	1.80	(0.08)	0.92 (0.13)	1.05 (0.12)	1.33 (0.10)	1.70 (0.09)	2.16 (0.12)	2.67 (0.20)	3.02 (0.26)	1.0	F	
	>70	136	1.81	(0.16)	0.95 (0.20) ^E	1.08 (0.19) ^E	1.33 (0.16)	1.68 (0.14)	2.12 (0.20)	2.61 (0.33)	2.95 (0.44)	1.0	F	
	19+	971	1.93	(0.07)	1.09 (0.13)	1.23 (0.12)	1.49 (0.09)	1.83 (0.07)	2.25 (0.10)	2.70 (0.18)	3.02 (0.25)	1.0	F	
Female														
	9-13	265	1.58	(0.08)	0.99 (0.18) ^E	1.09 (0.15)	1.28 (0.11)	1.51 (0.07)	1.77 (0.11)	2.03 (0.21)	2.20 (0.28)	0.7	<3	
	14-18	290	1.62	(0.13)	1.32 (0.30) ^E	1.39 (0.28) ^E	1.53 (0.23)	1.69 (0.18)	1.88 (0.20)	2.07 (0.34)	2.20 (0.49) ^E	0.9	F	
	19-30	197	1.60	(0.15)	0.92 (0.11)	1.04 (0.12)	1.27 (0.14)	1.55 (0.17)	1.89 (0.21)	2.26 (0.26)	2.50 (0.30)	0.9	F	
	31-50	312	1.41	(0.07)	0.89 (0.14)	0.97 (0.12)	1.14 (0.09)	1.36 (0.08)	1.63 (0.12)	1.92 (0.20)	2.13 (0.27)	0.9	F	
	51-70	312	1.41	(0.07)	0.90 (0.11)	0.98 (0.10)	1.14 (0.09)	1.35 (0.08)	1.61 (0.10)	1.86 (0.15)	2.03 (0.19)	0.9	F	
	>70	239	1.30	(0.07)	0.69 (0.13) ^E	0.78 (0.12)	0.96 (0.10)	1.22 (0.08)	1.56 (0.10)	1.95 (0.19)	2.23 (0.27)	0.9	F	
	19+	1060	1.43	(0.05)	0.84 (0.06)	0.94 (0.06)	1.13 (0.05)	1.38 (0.05)	1.67 (0.07)	1.99 (0.11)	2.22 (0.15)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.8 Thiamin (mg/d): Usual intakes from food, by DRI age-sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	129	1.12 (0.04)	0.77 (0.07)	0.84 (0.06)	0.96 (0.05)	1.11 (0.05)	1.29 (0.07)	1.48 (0.09)	1.61 (0.11)	0.4	<3	
	4-8	213	1.61 (0.06)	1.14 (0.06)	1.23 (0.07)	1.41 (0.07)	1.63 (0.08)	1.86 (0.09)	2.09 (0.09)	2.24 (0.10)	0.5	0.0	(0.0)
Male													
	9-13	122	2.17 (0.23)	1.45 (0.23)	1.58 (0.22)	1.83 (0.22)	2.17 (0.25)	2.59 (0.34)	3.07 (0.48)	3.40 (0.59) ^E	0.7	<3	
	14-18	150	2.43 (0.13)	1.55 (0.23)	1.73 (0.21)	2.04 (0.18)	2.47 (0.17)	2.98 (0.23)	3.52 (0.34)	3.88 (0.44)	1.0	<3	
	19-30	106	2.21 (0.18)	1.23 (0.14)	1.40 (0.15)	1.72 (0.16)	2.15 (0.19)	2.65 (0.26)	3.19 (0.35)	3.54 (0.44)	1.0	F	
	31-50	155	2.09 (0.13)	1.16 (0.25) ^E	1.36 (0.22)	1.72 (0.17)	2.13 (0.15)	2.54 (0.19)	2.92 (0.26)	3.16 (0.31)	1.0	F	
	51-70	122	1.67 (0.12)	0.96 (0.19) ^E	1.10 (0.16)	1.34 (0.13)	1.66 (0.14)	2.03 (0.23)	2.43 (0.38)	2.72 (0.49) ^E	1.0	F	
	>70	88	1.79 (0.10)	1.12 (0.17)	1.24 (0.16)	1.47 (0.15)	1.76 (0.14)	2.09 (0.15)	2.42 (0.19)	2.65 (0.23)	1.0	F	
	19+	471	1.97 (0.07)	1.09 (0.11)	1.25 (0.10)	1.56 (0.08)	1.95 (0.08)	2.41 (0.12)	2.90 (0.19)	3.23 (0.26)	1.0	F	
Female													
	9-13	103	1.88 (0.20)	1.52 (0.30) ^E	1.58 (0.27) ^E	1.68 (0.22)	1.80 (0.22)	1.95 (0.28)	2.10 (0.40) ^E	2.20 (0.49) ^E	0.7	<3	
	14-18	142	1.54 (0.09)	0.88 (0.10)	0.99 (0.10)	1.20 (0.11)	1.49 (0.13)	1.86 (0.17)	2.28 (0.24)	2.58 (0.29)	0.9	F	
	19-30	111	1.36 (0.08)	0.75 (0.13) ^E	0.85 (0.12)	1.04 (0.10)	1.29 (0.10)	1.59 (0.13)	1.90 (0.19)	2.11 (0.25)	0.9	F	
	31-50	146	1.56 (0.09)	1.27 (0.23) ^E	1.32 (0.20)	1.42 (0.15)	1.53 (0.12)	1.64 (0.18)	1.75 (0.28)	1.82 (0.36) ^E	0.9	F	
	51-70	184	1.45 (0.06)	0.96 (0.05)	1.04 (0.05)	1.19 (0.06)	1.37 (0.07)	1.58 (0.09)	1.81 (0.10)	1.96 (0.12)	0.9	F	
	>70	143	1.36 (0.06)	0.98 (0.11)	1.06 (0.10)	1.22 (0.08)	1.43 (0.08)	1.68 (0.11)	1.94 (0.16)	2.11 (0.20)	0.9	F	
	19+	584	1.46 (0.04)	0.91 (0.07)	1.01 (0.07)	1.19 (0.06)	1.42 (0.06)	1.68 (0.07)	1.95 (0.10)	2.13 (0.12)	0.9	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.9 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	169	1.13 (0.05)	0.73 (0.11)	0.80 (0.09)	0.94 (0.07)	1.11 (0.06)	1.30 (0.08)	1.48 (0.13)	1.61 (0.17)	0.4	<3	
	4-8	281	1.55 (0.06)	0.96 (0.13)	1.06 (0.12)	1.25 (0.09)	1.49 (0.06)	1.76 (0.10)	2.04 (0.17)	2.23 (0.21)	0.5	<3	
Male	9-13	183	1.95 (0.09)	1.44 (0.19)	1.54 (0.17)	1.72 (0.13)	1.94 (0.11)	2.20 (0.15)	2.46 (0.24)	2.63 (0.32)	0.7	<3	
	14-18	187	2.27 (0.12)	1.41 (0.24) ^E	1.57 (0.21)	1.85 (0.16)	2.22 (0.13)	2.66 (0.19)	3.10 (0.32)	3.40 (0.41)	1.0	<3	
	19-30	223	2.14 (0.10)	1.15 (0.20) ^E	1.33 (0.19)	1.67 (0.16)	2.10 (0.13)	2.59 (0.15)	3.09 (0.22)	3.42 (0.28)	1.0	F	
	31-50	229	2.18 (0.16)	1.27 (0.28) ^E	1.43 (0.26) ^E	1.75 (0.23)	2.20 (0.20)	2.78 (0.28)	3.46 (0.48)	3.95 (0.64)	1.0	F	
	51-70	197	1.76 (0.10)	0.97 (0.17) ^E	1.10 (0.16)	1.37 (0.13)	1.72 (0.12)	2.12 (0.15)	2.51 (0.21)	2.77 (0.27)	1.0	F	
	>70	72	1.78 (0.13)	1.12 (0.22) ^E	1.24 (0.20)	1.49 (0.18)	1.79 (0.16)	2.12 (0.19)	2.48 (0.27)	2.75 (0.35)	1.0	F	
	19+	721	2.04 (0.07)	1.08 (0.09)	1.25 (0.09)	1.57 (0.08)	2.01 (0.09)	2.57 (0.12)	3.20 (0.19)	3.66 (0.26)	1.0	F	
	Female	9-13	165	1.64 (0.10)	1.30 (0.08)	1.38 (0.09)	1.52 (0.10)	1.69 (0.11)	1.87 (0.13)	2.06 (0.14)	2.17 (0.15)	0.7	0.0 (0.0)
14-18	206	1.56 (0.08)	1.05 (0.15)	1.15 (0.13)	1.31 (0.11)	1.52 (0.10)	1.76 (0.13)	2.00 (0.19)	2.14 (0.23)	0.9	F		
19-30	191	1.40 (0.09)	0.80 (0.15) ^E	0.90 (0.14)	1.09 (0.13)	1.34 (0.12)	1.63 (0.15)	1.94 (0.21)	2.14 (0.26)	0.9	F		
31-50	258	1.42 (0.06)	1.13 (0.18)	1.19 (0.16)	1.29 (0.12)	1.42 (0.08)	1.55 (0.10)	1.69 (0.17)	1.77 (0.23)	0.9	F		
51-70	249	1.43 (0.09)	0.89 (0.14)	0.99 (0.12)	1.18 (0.10)	1.41 (0.10)	1.68 (0.14)	1.96 (0.23)	2.17 (0.30)	0.9	F		
>70	128	1.15 (0.06)	0.61 (0.08)	0.69 (0.08)	0.84 (0.07)	1.04 (0.07)	1.28 (0.08)	1.55 (0.12)	1.73 (0.16)	0.9	32.8 (8.7) ^E		
19+	826	1.39 (0.04)	0.83 (0.06)	0.93 (0.06)	1.12 (0.05)	1.35 (0.05)	1.63 (0.06)	1.93 (0.08)	2.14 (0.11)	0.9	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.10 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	192	1.24 (0.06)	0.84 (0.10)	0.91 (0.09)	1.05 (0.08)	1.23 (0.08)	1.42 (0.10)	1.62 (0.14)	1.75 (0.16)	0.4	0.0 (0.0)	
	4-8	321	1.61 (0.07)	1.07 (0.11)	1.17 (0.09)	1.35 (0.07)	1.57 (0.07)	1.82 (0.10)	2.09 (0.16)	2.27 (0.21)	0.5	<3	
Male	9-13	226	1.94 (0.09)	1.26 (0.17)	1.39 (0.15)	1.63 (0.13)	1.93 (0.11)	2.27 (0.14)	2.65 (0.21)	2.92 (0.29)	0.7	<3	
	14-18	262	2.59 (0.20)	1.38 (0.33) ^E	1.59 (0.31) ^E	1.98 (0.27)	2.49 (0.24)	3.12 (0.28)	3.81 (0.41)	4.30 (0.54)	1.0	<3	
	19-30	197	2.32 (0.15)	1.22 (0.15)	1.38 (0.14)	1.70 (0.14)	2.14 (0.16)	2.71 (0.22)	3.35 (0.35)	3.81 (0.47)	1.0	F	
	31-50	282	2.40 (0.17)	1.27 (0.19)	1.46 (0.17)	1.82 (0.15)	2.30 (0.15)	2.88 (0.27)	3.54 (0.44)	4.03 (0.55)	1.0	F	
	51-70	234	2.07 (0.08)	1.10 (0.12)	1.25 (0.11)	1.55 (0.11)	1.97 (0.10)	2.47 (0.13)	2.98 (0.19)	3.33 (0.25)	1.0	F	
	>70	119	1.76 (0.14)	0.91 (0.21) ^E	1.07 (0.19) ^E	1.36 (0.17)	1.70 (0.16)	2.07 (0.22)	2.48 (0.29)	2.77 (0.36)	1.0	F	
	19+	832	2.23 (0.08)	1.15 (0.07)	1.33 (0.07)	1.66 (0.07)	2.10 (0.08)	2.66 (0.11)	3.29 (0.17)	3.77 (0.23)	1.0	F	
	Female	9-13	226	1.67 (0.10)	0.95 (0.10)	1.07 (0.10)	1.30 (0.10)	1.60 (0.12)	1.96 (0.14)	2.34 (0.18)	2.60 (0.22)	0.7	<3
14-18		242	1.56 (0.08)	0.82 (0.11)	0.93 (0.11)	1.16 (0.10)	1.46 (0.09)	1.81 (0.12)	2.20 (0.17)	2.48 (0.23)	0.9	F	
19-30		208	1.48 (0.09)	1.07 (0.17)	1.16 (0.15)	1.31 (0.13)	1.49 (0.11)	1.69 (0.14)	1.89 (0.19)	2.02 (0.24)	0.9	F	
31-50		263	1.56 (0.08)	0.80 (0.08)	0.91 (0.08)	1.12 (0.07)	1.40 (0.08)	1.77 (0.13)	2.23 (0.22)	2.58 (0.31)	0.9	F	
51-70		322	1.50 (0.12)	0.73 (0.11)	0.84 (0.11)	1.06 (0.11)	1.38 (0.11)	1.80 (0.16)	2.32 (0.26)	2.71 (0.34)	0.9	F	
>70		198	1.32 (0.06)	0.70 (0.07)	0.81 (0.07)	1.00 (0.06)	1.25 (0.07)	1.55 (0.09)	1.89 (0.13)	2.13 (0.17)	0.9	16.3 (4.9) ^E	
19+		991	1.50 (0.05)	0.77 (0.05)	0.89 (0.05)	1.11 (0.05)	1.39 (0.06)	1.77 (0.08)	2.22 (0.13)	2.56 (0.18)	0.9	10.5 (2.6) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.11 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	348	1.19	(0.04)	0.82 (0.05)	0.89 (0.05)	1.01 (0.05)	1.16 (0.05)	1.34 (0.06)	1.53 (0.08)	1.66 (0.10)	0.4	0.0 (0.0)	
	4-8	554	1.63	(0.05)	1.07 (0.11)	1.18 (0.10)	1.39 (0.08)	1.65 (0.07)	1.93 (0.07)	2.21 (0.10)	2.39 (0.13)	0.5	<3	
Male														
	9-13	409	1.87	(0.07)	1.23 (0.11)	1.33 (0.10)	1.52 (0.09)	1.77 (0.08)	2.07 (0.10)	2.38 (0.15)	2.59 (0.19)	0.7	<3	
	14-18	414	2.10	(0.10)	1.30 (0.14)	1.45 (0.14)	1.74 (0.14)	2.11 (0.14)	2.57 (0.17)	3.06 (0.23)	3.39 (0.29)	1.0	<3	
	19-30	311	2.40	(0.19)	1.43 (0.16)	1.59 (0.16)	1.90 (0.15)	2.32 (0.18)	2.85 (0.25)	3.45 (0.38)	3.88 (0.49)	1.0	<3	
	31-50	489	1.86	(0.08)	1.01 (0.08)	1.14 (0.08)	1.38 (0.08)	1.71 (0.09)	2.14 (0.11)	2.63 (0.15)	2.98 (0.19)	1.0	F	
	51-70	575	1.79	(0.07)	1.24 (0.14)	1.34 (0.12)	1.53 (0.10)	1.76 (0.08)	2.01 (0.10)	2.26 (0.14)	2.43 (0.18)	1.0	<3	
	>70	239	1.81	(0.09)	1.20 (0.10)	1.31 (0.10)	1.50 (0.10)	1.76 (0.11)	2.09 (0.13)	2.44 (0.17)	2.69 (0.21)	1.0	<3	
	19+	1614	1.94	(0.06)	1.14 (0.05)	1.27 (0.05)	1.51 (0.05)	1.84 (0.06)	2.27 (0.08)	2.76 (0.13)	3.11 (0.17)	1.0	F	
Female														
	9-13	355	1.60	(0.07)	1.08 (0.09)	1.18 (0.09)	1.36 (0.08)	1.60 (0.09)	1.89 (0.11)	2.22 (0.17)	2.44 (0.21)	0.7	<3	
	14-18	410	1.52	(0.08)	0.83 (0.12)	0.96 (0.11)	1.19 (0.10)	1.48 (0.09)	1.81 (0.10)	2.17 (0.15)	2.42 (0.19)	0.9	F	
	19-30	384	1.45	(0.07)	1.04 (0.12)	1.13 (0.12)	1.28 (0.11)	1.48 (0.10)	1.70 (0.11)	1.93 (0.14)	2.08 (0.17)	0.9	F	
	31-50	585	1.48	(0.05)	0.74 (0.06)	0.88 (0.06)	1.13 (0.06)	1.45 (0.06)	1.80 (0.08)	2.17 (0.10)	2.43 (0.13)	0.9	11.1 (2.9) ^E	
	51-70	711	1.47	(0.05)	0.90 (0.07)	1.00 (0.07)	1.17 (0.06)	1.41 (0.05)	1.69 (0.07)	1.97 (0.11)	2.15 (0.13)	0.9	F	
	>70	342	1.50	(0.07)	0.85 (0.09)	0.95 (0.09)	1.16 (0.09)	1.46 (0.09)	1.81 (0.10)	2.17 (0.13)	2.42 (0.15)	0.9	F	
	19+	2022	1.47	(0.03)	0.87 (0.04)	0.98 (0.04)	1.19 (0.04)	1.45 (0.04)	1.76 (0.04)	2.07 (0.05)	2.27 (0.06)	0.9	6.3 (1.5) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.12 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both														
	1-3	622	1.14	(0.03)	0.67 (0.04)	0.76 (0.04)	0.91 (0.04)	1.11 (0.04)	1.34 (0.05)	1.58 (0.07)	1.74 (0.09)	0.4	<3	
	4-8	919	1.55	(0.04)	1.01 (0.09)	1.11 (0.08)	1.29 (0.06)	1.51 (0.04)	1.77 (0.06)	2.03 (0.11)	2.21 (0.15)	0.5	<3	
Male														
	9-13	579	1.99	(0.07)	1.50 (0.16)	1.60 (0.14)	1.78 (0.11)	2.00 (0.08)	2.25 (0.11)	2.50 (0.17)	2.67 (0.22)	0.7	0.0 (0.0)	
	14-18	634	2.32	(0.08)	1.30 (0.11)	1.48 (0.10)	1.82 (0.09)	2.27 (0.09)	2.81 (0.13)	3.39 (0.19)	3.78 (0.23)	1.0	<3	
	19-30	578	2.20	(0.08)	1.23 (0.16)	1.41 (0.15)	1.73 (0.12)	2.14 (0.10)	2.63 (0.13)	3.15 (0.21)	3.49 (0.28)	1.0	F	
	31-50	693	2.08	(0.10)	1.19 (0.19)	1.35 (0.18)	1.65 (0.15)	2.07 (0.13)	2.59 (0.17)	3.17 (0.27)	3.58 (0.37)	1.0	F	
	51-70	596	1.75	(0.06)	0.91 (0.08)	1.06 (0.08)	1.34 (0.08)	1.70 (0.08)	2.12 (0.10)	2.59 (0.15)	2.92 (0.19)	1.0	F	
	>70	296	1.79	(0.08)	1.09 (0.07)	1.22 (0.08)	1.45 (0.09)	1.75 (0.10)	2.10 (0.11)	2.48 (0.13)	2.75 (0.15)	1.0	F	
	19+	2163	2.00	(0.05)	1.08 (0.06)	1.24 (0.06)	1.55 (0.05)	1.96 (0.06)	2.48 (0.08)	3.04 (0.11)	3.45 (0.15)	1.0	3.2 (1.0) ^E	
Female														
	9-13	533	1.67	(0.07)	1.27 (0.16)	1.35 (0.14)	1.50 (0.11)	1.68 (0.09)	1.89 (0.11)	2.09 (0.18)	2.22 (0.23)	0.7	<3	
	14-18	638	1.57	(0.06)	0.98 (0.09)	1.07 (0.09)	1.25 (0.08)	1.53 (0.08)	1.87 (0.10)	2.24 (0.15)	2.49 (0.20)	0.9	F	
	19-30	499	1.44	(0.07)	0.81 (0.10)	0.92 (0.09)	1.12 (0.08)	1.38 (0.08)	1.69 (0.11)	2.00 (0.15)	2.21 (0.19)	0.9	F	
	31-50	716	1.44	(0.05)	1.00 (0.13)	1.08 (0.11)	1.23 (0.09)	1.41 (0.06)	1.63 (0.07)	1.85 (0.11)	2.00 (0.15)	0.9	F	
	51-70	745	1.43	(0.06)	0.92 (0.08)	1.01 (0.08)	1.19 (0.06)	1.40 (0.06)	1.64 (0.08)	1.90 (0.13)	2.08 (0.17)	0.9	F	
	>70	510	1.24	(0.04)	0.69 (0.05)	0.78 (0.05)	0.95 (0.04)	1.17 (0.04)	1.45 (0.06)	1.78 (0.09)	2.02 (0.11)	0.9	20.3 (4.1) ^E	
	19+	2470	1.41	(0.03)	0.82 (0.04)	0.93 (0.03)	1.12 (0.03)	1.37 (0.03)	1.65 (0.04)	1.97 (0.06)	2.19 (0.07)	0.9	8.5 (1.8) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 23.13 Thiamin (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	1.21 (0.02)	0.74 (0.03)	0.83 (0.03)	0.99 (0.02)	1.18 (0.03)	1.43 (0.03)	1.69 (0.05)	1.87 (0.07)	0.4	<3	
	4-8	3235	1.61 (0.02)	1.01 (0.03)	1.12 (0.03)	1.33 (0.03)	1.59 (0.03)	1.90 (0.04)	2.23 (0.06)	2.46 (0.08)	0.5	<3	
Male													
	9-13	2080	2.06 (0.05)	1.30 (0.06)	1.43 (0.06)	1.68 (0.05)	2.01 (0.05)	2.41 (0.07)	2.86 (0.10)	3.18 (0.14)	0.7	<3	
	14-18	2288	2.39 (0.05)	1.35 (0.07)	1.54 (0.07)	1.89 (0.06)	2.36 (0.06)	2.93 (0.08)	3.56 (0.11)	4.00 (0.14)	1.0	<3	
	19-30	1804	2.14 (0.05)	1.18 (0.06)	1.34 (0.06)	1.64 (0.06)	2.05 (0.06)	2.54 (0.08)	3.08 (0.12)	3.46 (0.16)	1.0	F	
	31-50	2596	2.04 (0.05)	1.14 (0.06)	1.29 (0.06)	1.57 (0.05)	1.96 (0.06)	2.45 (0.07)	2.99 (0.10)	3.37 (0.14)	1.0	F	
	51-70	2550	1.88 (0.03)	1.08 (0.04)	1.22 (0.04)	1.48 (0.03)	1.80 (0.03)	2.18 (0.04)	2.59 (0.07)	2.89 (0.09)	1.0	3.1 (0.8) ^E	
	>70	1520	1.68 (0.04)	0.97 (0.05)	1.10 (0.05)	1.32 (0.04)	1.61 (0.05)	1.96 (0.07)	2.32 (0.10)	2.57 (0.12)	1.0	5.9 (1.6) ^E	
	19+	8470	1.99 (0.02)	1.10 (0.03)	1.25 (0.03)	1.52 (0.03)	1.90 (0.03)	2.36 (0.04)	2.88 (0.06)	3.25 (0.08)	1.0	2.8 (0.5) ^E	
Female													
	9-13	1980	1.65 (0.03)	1.03 (0.05)	1.14 (0.04)	1.35 (0.04)	1.62 (0.04)	1.93 (0.04)	2.25 (0.07)	2.47 (0.09)	0.7	<3	
	14-18	2256	1.64 (0.03)	0.93 (0.04)	1.06 (0.04)	1.28 (0.03)	1.58 (0.04)	1.96 (0.05)	2.41 (0.09)	2.72 (0.13)	0.9	4.1 (1.0) ^E	
	19-30	1854	1.48 (0.03)	0.89 (0.05)	0.99 (0.04)	1.19 (0.04)	1.44 (0.04)	1.70 (0.05)	1.97 (0.07)	2.16 (0.08)	0.9	F	
	31-50	2686	1.48 (0.03)	0.83 (0.03)	0.94 (0.03)	1.15 (0.03)	1.43 (0.03)	1.75 (0.04)	2.11 (0.05)	2.36 (0.07)	0.9	7.9 (1.5) ^E	
	51-70	3200	1.46 (0.03)	0.89 (0.03)	0.98 (0.03)	1.16 (0.03)	1.40 (0.03)	1.70 (0.04)	2.03 (0.06)	2.25 (0.08)	0.9	5.6 (1.5) ^E	
	>70	2610	1.37 (0.03)	0.78 (0.03)	0.88 (0.03)	1.07 (0.03)	1.31 (0.04)	1.61 (0.04)	1.95 (0.06)	2.18 (0.07)	0.9	11.4 (2.2) ^E	
	19+	10350	1.46 (0.01)	0.84 (0.02)	0.95 (0.02)	1.15 (0.02)	1.41 (0.02)	1.72 (0.02)	2.06 (0.03)	2.29 (0.04)	0.9	7.6 (0.8)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

24. Vitamin B₆ (mg/d): Usual intakes from food

Table 24.1 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)		UL ³	% >UL (SE)									
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)												
Both																								
	1-3	79	1.24	(0.10)	0.86	(0.08)	0.93	(0.09)	1.04	(0.10)	1.18	(0.11)	1.34	(0.12)	1.49	(0.14)	1.59	(0.15)	0.4	0.0	(0.0)	30	0.0	(0.0)
	4-8	127	1.53	(0.10)	0.85	(0.11)	0.96	(0.11)	1.17	(0.10)	1.45	(0.12)	1.80	(0.15)	2.19	(0.22)	2.45	(0.28)	0.5	<3		40	0.0	(0.0)
Male																								
	9-13	111	1.98	(0.15)	1.37	(0.19)	1.48	(0.18)	1.68	(0.17)	1.93	(0.18)	2.22	(0.23)	2.50	(0.31)	2.68	(0.37)	0.8	<3		60	0.0	(0.0)
	14-18	107	2.02	(0.16)	1.40	(0.27) ^E	1.53	(0.26) ^E	1.77	(0.25)	2.10	(0.26)	2.49	(0.35)	2.92	(0.54) ^E	3.22	(0.73) ^E	1.1	F		80	0.0	(0.0)
	19-30	77	1.89	(0.10)	1.28	(0.12)	1.39	(0.12)	1.57	(0.12)	1.79	(0.14)	2.02	(0.16)	2.26	(0.18)	2.41	(0.19)	1.1	F		100	0.0	(0.0)
	31-50	145	2.28	(0.15)	1.49	(0.24)	1.65	(0.21)	1.93	(0.18)	2.27	(0.17)	2.64	(0.23)	3.02	(0.32)	3.27	(0.39)	1.1	F		100	0.0	(0.0)
	51-70	182	1.95	(0.11)	1.35	(0.23) ^E	1.46	(0.21)	1.64	(0.16)	1.87	(0.13)	2.12	(0.16)	2.37	(0.26)	2.53	(0.34)	1.4	F		100	0.0	(0.0)
	>70	63	2.14	(0.22)	1.28	(0.23) ^E	1.43	(0.22)	1.72	(0.23)	2.13	(0.27)	2.66	(0.39)	3.26	(0.58) ^E	3.70	(0.73) ^E	1.4	F		100	0.0	(0.0)
	19+	467	2.09	(0.08)	1.34	(0.14)	1.48	(0.12)	1.74	(0.10)	2.06	(0.09)	2.43	(0.11)	2.81	(0.17)	3.07	(0.21)				100	0.0	(0.0)
Female																								
	9-13	96	1.48	(0.12)	1.07	(0.17)	1.16	(0.16)	1.32	(0.15)	1.52	(0.15)	1.74	(0.17)	1.95	(0.22)	2.09	(0.25)	0.8	F		60	0.0	(0.0)
	14-18	105	1.58	(0.13)	1.08	(0.20) ^E	1.19	(0.20)	1.39	(0.18)	1.64	(0.18)	1.93	(0.21)	2.21	(0.28)	2.39	(0.33)	1.0	F		80	0.0	(0.0)
	19-30	91	1.40	(0.10)	0.82	(0.14) ^E	0.92	(0.13)	1.10	(0.12)	1.32	(0.13)	1.57	(0.16)	1.81	(0.22)	1.97	(0.27)	1.1	F		100	0.0	(0.0)
	31-50	167	1.42	(0.11)	0.70	(0.13) ^E	0.83	(0.13)	1.09	(0.12)	1.42	(0.12)	1.77	(0.18)	2.23	(0.28)	2.58	(0.36)	1.1	F		100	0.0	(0.0)
	51-70	198	1.68	(0.09)	1.07	(0.20) ^E	1.19	(0.17)	1.39	(0.12)	1.63	(0.10)	1.90	(0.13)	2.19	(0.22)	2.39	(0.29)	1.3	F		100	0.0	(0.0)
	>70	74	1.53	(0.12)	0.94	(0.19) ^E	1.05	(0.18) ^E	1.25	(0.16)	1.50	(0.15)	1.77	(0.17)	2.03	(0.22)	2.20	(0.27)	1.3	F		100	0.0	(0.0)
	19+	530	1.51	(0.05)	0.81	(0.08)	0.94	(0.07)	1.19	(0.07)	1.49	(0.07)	1.82	(0.08)	2.16	(0.11)	2.41	(0.14)				100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.2 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		UL ³	% >UL	
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)	(<i>SE</i>)		(<i>SE</i>)	
Both																
	1-3	58	1.31	(0.08)	0.93 (0.15)	1.01 (0.14)	1.16 (0.13)	1.34 (0.12)	1.53 (0.12)	1.71 (0.14)	1.82 (0.16)	0.4	<3	30	0.0	(0.0)
	4-8	110	1.34	(0.08)	0.92 (0.14)	1.01 (0.13)	1.15 (0.11)	1.33 (0.10)	1.51 (0.12)	1.69 (0.15)	1.80 (0.17)	0.5	<3	40	0.0	(0.0)
Male																
	9-13	95	1.72	(0.10)	1.22 (0.11)	1.32 (0.12)	1.49 (0.13)	1.69 (0.14)	1.91 (0.16)	2.12 (0.18)	2.26 (0.19)	0.8	<3	60	0.0	(0.0)
	14-18	87	2.04	(0.16)	1.03 (0.21) ^E	1.22 (0.20)	1.58 (0.19)	2.05 (0.20)	2.63 (0.29)	3.27 (0.43)	3.72 (0.55)	1.1	F	80	0.0	(0.0)
	19-30	70	2.14	(0.12)	1.31 (0.18)	1.48 (0.17)	1.79 (0.16)	2.15 (0.16)	2.56 (0.20)	3.00 (0.25)	3.29 (0.30)	1.1	F	100	0.0	(0.0)
	31-50	109	1.97	(0.13)	1.06 (0.15)	1.21 (0.15)	1.49 (0.15)	1.87 (0.15)	2.33 (0.19)	2.83 (0.27)	3.17 (0.34)	1.1	F	100	0.0	(0.0)
	51-70	128	1.97	(0.11)	1.06 (0.20) ^E	1.21 (0.19)	1.50 (0.16)	1.88 (0.13)	2.34 (0.15)	2.83 (0.23)	3.16 (0.31)	1.4	F	100	0.0	(0.0)
	>70	65	1.53	(0.08)	0.97 (0.11)	1.07 (0.10)	1.25 (0.08)	1.46 (0.09)	1.72 (0.13)	1.99 (0.20)	2.17 (0.24)	1.4	43.3 (11.3) ^E	100	0.0	(0.0)
	19+	372	1.96	(0.07)	1.09 (0.08)	1.24 (0.08)	1.53 (0.08)	1.90 (0.08)	2.33 (0.10)	2.81 (0.14)	3.15 (0.18)			100	0.0	(0.0)
Female																
	9-13	75	1.42	(0.11)	1.01 (0.14)	1.09 (0.14)	1.23 (0.16)	1.39 (0.18)	1.56 (0.20)	1.74 (0.23)	1.85 (0.24)	0.8	F	60	0.0	(0.0)
	14-18	81	1.50	(0.09)	0.88 (0.15) ^E	1.01 (0.14)	1.23 (0.13)	1.49 (0.13)	1.75 (0.14)	1.99 (0.16)	2.13 (0.18)	1.0	F	80	0.0	(0.0)
	19-30	101	1.75	(0.16)	1.25 (0.21) ^E	1.38 (0.20)	1.61 (0.20)	1.89 (0.22)	2.19 (0.28)	2.49 (0.34)	2.68 (0.39)	1.1	F	100	0.0	(0.0)
	31-50	116	1.52	(0.11)	0.86 (0.15) ^E	0.99 (0.15)	1.22 (0.14)	1.48 (0.14)	1.76 (0.16)	2.02 (0.20)	2.19 (0.23)	1.1	F	100	0.0	(0.0)
	51-70	146	1.65	(0.08)	1.02 (0.14)	1.15 (0.13)	1.38 (0.11)	1.66 (0.10)	1.97 (0.12)	2.28 (0.17)	2.48 (0.20)	1.3	F	100	0.0	(0.0)
	>70	94	1.15	(0.08)	0.59 (0.15) ^E	0.68 (0.14) ^E	0.83 (0.11)	1.03 (0.10)	1.25 (0.11)	1.47 (0.16)	1.60 (0.20)	1.3	79.4 (10.3)	100	0.0	(0.0)
	19+	457	1.56	(0.06)	0.94 (0.08)	1.06 (0.08)	1.28 (0.08)	1.54 (0.08)	1.83 (0.09)	2.12 (0.12)	2.29 (0.14)			100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.3 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		(SE)	UL ³	% >UL		(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)						
Both																		
	1-3	112	1.21	(0.07)	0.81 (0.09)	0.88 (0.10)	1.01 (0.10)	1.18 (0.10)	1.36 (0.11)	1.54 (0.12)	1.65 (0.12)	0.4	0.0	(0.0)	30	0.0	(0.0)	
	4-8	177	1.41	(0.06)	0.93 (0.06)	1.03 (0.07)	1.20 (0.07)	1.42 (0.07)	1.66 (0.08)	1.89 (0.09)	2.05 (0.10)	0.5	<3		40	0.0	(0.0)	
Male																		
	9-13	111	1.73	(0.10)	0.91 (0.12)	1.04 (0.11)	1.29 (0.11)	1.61 (0.11)	1.97 (0.12)	2.35 (0.17)	2.60 (0.21)	0.8	F		60	0.0	(0.0)	
	14-18	113	1.96	(0.16)	1.24 (0.14)	1.37 (0.15)	1.60 (0.18)	1.91 (0.22)	2.26 (0.26)	2.62 (0.29)	2.84 (0.30)	1.1	F		80	0.0	(0.0)	
	19-30	91	2.31	(0.22)	1.31 (0.18)	1.48 (0.19)	1.80 (0.21)	2.20 (0.24)	2.67 (0.28)	3.14 (0.34)	3.45 (0.38)	1.1	F		100	0.0	(0.0)	
	31-50	101	2.01	(0.11)	1.24 (0.23) ^E	1.40 (0.20)	1.67 (0.16)	1.99 (0.16)	2.34 (0.20)	2.68 (0.27)	2.89 (0.33)	1.1	F		100	0.0	(0.0)	
	51-70	134	1.98	(0.10)	1.45 (0.19)	1.55 (0.17)	1.72 (0.13)	1.93 (0.11)	2.16 (0.12)	2.38 (0.19)	2.52 (0.25)	1.4	F		100	0.0	(0.0)	
	>70	56	1.72	(0.15)	1.11 (0.22) ^E	1.21 (0.21) ^E	1.39 (0.21)	1.63 (0.21)	1.93 (0.24)	2.26 (0.33)	2.49 (0.42) ^E	1.4	F		100	0.0	(0.0)	
	19+	382	2.03	(0.07)	1.32 (0.12)	1.45 (0.11)	1.68 (0.10)	1.97 (0.10)	2.30 (0.12)	2.61 (0.15)	2.81 (0.18)				100	0.0	(0.0)	
Female																		
	9-13	105	1.50	(0.14)	1.07 (0.11)	1.15 (0.11)	1.28 (0.12)	1.44 (0.13)	1.61 (0.15)	1.79 (0.17)	1.91 (0.18)	0.8	<3		60	0.0	(0.0)	
	14-18	120	1.26	(0.15)	0.63 (0.17) ^E	0.73 (0.16) ^E	0.93 (0.16) ^E	1.19 (0.17)	1.54 (0.23)	1.96 (0.32)	2.27 (0.41) ^E	1.0	F		80	0.0	(0.0)	
	19-30	91	1.53	(0.11)	1.16 (0.23) ^E	1.23 (0.20)	1.36 (0.16)	1.51 (0.14)	1.66 (0.17)	1.81 (0.24)	1.90 (0.30)	1.1	F		100	0.0	(0.0)	
	31-50	159	1.54	(0.07)	0.76 (0.12)	0.92 (0.11)	1.20 (0.10)	1.52 (0.10)	1.85 (0.12)	2.20 (0.16)	2.43 (0.20)	1.1	18.7 (6.1) ^E		100	0.0	(0.0)	
	51-70	174	1.54	(0.09)	0.97 (0.17) ^E	1.07 (0.16)	1.25 (0.13)	1.48 (0.11)	1.74 (0.14)	2.01 (0.22)	2.19 (0.30)	1.3	F		100	0.0	(0.0)	
	>70	80	1.71	(0.11)	1.15 (0.16)	1.28 (0.14)	1.51 (0.12)	1.78 (0.12)	2.09 (0.15)	2.45 (0.23)	2.72 (0.30)	1.3	F		100	0.0	(0.0)	
	19+	504	1.56	(0.05)	0.90 (0.08)	1.02 (0.07)	1.25 (0.06)	1.54 (0.06)	1.85 (0.08)	2.17 (0.10)	2.38 (0.12)				100	0.0	(0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.4 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)								
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)							95th (SE)							
Both																								
	1-3	99	1.40	(0.11)	0.89	(0.14)	0.99	(0.13)	1.18	(0.12)	1.43	(0.13)	1.74	(0.17)	2.07	(0.23)	2.29	(0.29)	0.4	<3		30	0.0	(0.0)
	4-8	140	1.69	(0.16)	1.02	(0.17)	1.14	(0.16)	1.35	(0.15)	1.65	(0.17)	2.03	(0.24)	2.46	(0.37)	2.78	(0.49) ^E	0.5	<3		40	0.0	(0.0)
Male																								
	9-13	92	1.63	(0.12)	1.10	(0.13)	1.18	(0.13)	1.33	(0.13)	1.53	(0.15)	1.78	(0.20)	2.05	(0.28)	2.23	(0.36)	0.8	<3		60	0.0	(0.0)
	14-18	107	2.07	(0.14)	1.34	(0.24) ^E	1.48	(0.22)	1.72	(0.18)	2.02	(0.17)	2.37	(0.21)	2.72	(0.29)	2.95	(0.35)	1.1	F		80	0.0	(0.0)
	19-30	73	2.57	(0.30)	1.28	(0.25) ^E	1.50	(0.24)	1.91	(0.25)	2.44	(0.29)	3.10	(0.39)	3.88	(0.57)	4.46	(0.74) ^E	1.1	F		100	0.0	(0.0)
	31-50	134	2.11	(0.18)	1.26	(0.13)	1.37	(0.15)	1.59	(0.18)	1.94	(0.21)	2.39	(0.24)	2.85	(0.28)	3.17	(0.31)	1.1	F		100	0.0	(0.0)
	51-70	131	2.00	(0.17)	1.09	(0.22) ^E	1.25	(0.21) ^E	1.55	(0.19)	1.94	(0.19)	2.37	(0.23)	2.81	(0.31)	3.09	(0.37)	1.4	F		100	0.0	(0.0)
	>70	55	1.83	(0.14)	0.98	(0.24) ^E	1.13	(0.22) ^E	1.40	(0.20)	1.74	(0.18)	2.13	(0.21)	2.55	(0.28)	2.84	(0.34)	1.4	F		100	0.0	(0.0)
	19+	393	2.15	(0.11)	1.09	(0.15)	1.27	(0.14)	1.61	(0.12)	2.07	(0.12)	2.62	(0.16)	3.21	(0.24)	3.61	(0.31)				100	0.0	(0.0)
Female																								
	9-13	79	1.59	(0.14)	1.23	(0.22) ^E	1.31	(0.21)	1.46	(0.20)	1.65	(0.20)	1.87	(0.24)	2.09	(0.32)	2.23	(0.39) ^E	0.8	<3		60	0.0	(0.0)
	14-18	104	1.46	(0.08)	1.01	(0.08)	1.10	(0.08)	1.25	(0.09)	1.44	(0.09)	1.64	(0.11)	1.83	(0.13)	1.95	(0.15)	1.0	F		80	0.0	(0.0)
	19-30	101	1.44	(0.10)	1.07	(0.19) ^E	1.17	(0.17)	1.35	(0.15)	1.57	(0.14)	1.81	(0.17)	2.04	(0.23)	2.19	(0.29)	1.1	F		100	0.0	(0.0)
	31-50	143	1.48	(0.09)	0.72	(0.18) ^E	0.86	(0.16) ^E	1.11	(0.12)	1.41	(0.11)	1.73	(0.15)	2.05	(0.23)	2.29	(0.29)	1.1	F		100	0.0	(0.0)
	51-70	193	1.46	(0.08)	0.85	(0.13)	0.95	(0.12)	1.13	(0.10)	1.37	(0.10)	1.64	(0.13)	1.92	(0.19)	2.10	(0.24)	1.3	42.9	(11.7) ^E	100	0.0	(0.0)
	>70	94	1.53	(0.11)	0.92	(0.13)	1.02	(0.12)	1.19	(0.11)	1.41	(0.12)	1.66	(0.17)	1.90	(0.23)	2.06	(0.28)	1.3	F		100	0.0	(0.0)
19+	531	1.47	(0.05)	0.77	(0.07)	0.89	(0.06)	1.13	(0.06)	1.42	(0.06)	1.75	(0.07)	2.07	(0.10)	2.29	(0.12)				100	0.0	(0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.5 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	311	1.36	(0.07)	0.84 (0.11)	0.93 (0.11)	1.10 (0.10)	1.33 (0.09)	1.62 (0.12)	1.91 (0.16)	2.11 (0.21)	0.4	<3		30	0.0 (0.0)	
	4-8	485	1.55	(0.05)	0.98 (0.05)	1.08 (0.05)	1.28 (0.05)	1.52 (0.07)	1.80 (0.08)	2.06 (0.10)	2.22 (0.11)	0.5	<3		40	0.0 (0.0)	
Male																	
	9-13	277	1.98	(0.12)	1.30 (0.18)	1.43 (0.17)	1.67 (0.16)	1.97 (0.16)	2.33 (0.19)	2.70 (0.25)	2.95 (0.30)	0.8	<3		60	0.0 (0.0)	
	14-18	339	2.33	(0.10)	1.43 (0.13)	1.61 (0.12)	1.96 (0.12)	2.41 (0.13)	2.96 (0.17)	3.55 (0.25)	3.94 (0.32)	1.1	<3		80	0.0 (0.0)	
	19-30	237	2.33	(0.15)	1.66 (0.26)	1.80 (0.24)	2.08 (0.22)	2.44 (0.23)	2.89 (0.33)	3.37 (0.51)	3.70 (0.68) ^E	1.1	<3		100	0.0 (0.0)	
	31-50	423	2.25	(0.10)	1.55 (0.18)	1.69 (0.16)	1.94 (0.13)	2.23 (0.11)	2.56 (0.14)	2.89 (0.20)	3.10 (0.25)	1.1	<3		100	0.0 (0.0)	
	51-70	387	2.04	(0.09)	1.34 (0.20)	1.48 (0.18)	1.75 (0.15)	2.08 (0.12)	2.45 (0.13)	2.83 (0.21)	3.09 (0.27)	1.4	^F		100	0.0 (0.0)	
	>70	132	1.80	(0.13)	0.89 (0.26) ^E	1.06 (0.25) ^E	1.38 (0.23)	1.81 (0.21)	2.27 (0.22)	2.75 (0.28)	3.09 (0.34)	1.4	^F		100	0.0 (0.0)	
	19+	1179	2.17	(0.06)	1.38 (0.10)	1.53 (0.09)	1.80 (0.08)	2.17 (0.07)	2.62 (0.10)	3.08 (0.15)	3.41 (0.20)				100	0.0 (0.0)	
Female																	
	9-13	281	1.55	(0.07)	0.80 (0.08)	0.92 (0.08)	1.14 (0.08)	1.44 (0.08)	1.78 (0.11)	2.14 (0.15)	2.38 (0.18)	0.8	^F		60	0.0 (0.0)	
	14-18	321	1.51	(0.06)	1.03 (0.11)	1.13 (0.10)	1.31 (0.08)	1.53 (0.08)	1.77 (0.10)	2.01 (0.14)	2.17 (0.18)	1.0	^F		80	0.0 (0.0)	
	19-30	249	1.73	(0.09)	1.26 (0.18)	1.37 (0.16)	1.54 (0.13)	1.75 (0.11)	1.98 (0.20)	2.18 (0.32)	2.31 (0.40) ^E	1.1	^F		100	0.0 (0.0)	
	31-50	364	1.75	(0.09)	1.05 (0.15)	1.18 (0.14)	1.42 (0.13)	1.74 (0.12)	2.08 (0.14)	2.44 (0.20)	2.70 (0.26)	1.1	^F		100	0.0 (0.0)	
	51-70	467	1.73	(0.07)	1.08 (0.15)	1.21 (0.13)	1.44 (0.10)	1.72 (0.08)	2.04 (0.11)	2.37 (0.18)	2.58 (0.23)	1.3	^F		100	0.0 (0.0)	
	>70	215	1.48	(0.07)	0.93 (0.12)	1.03 (0.12)	1.21 (0.11)	1.45 (0.10)	1.72 (0.11)	2.00 (0.14)	2.19 (0.17)	1.3	^F		100	0.0 (0.0)	
	19+	1295	1.70	(0.05)	1.07 (0.07)	1.20 (0.07)	1.43 (0.06)	1.71 (0.06)	2.03 (0.07)	2.36 (0.09)	2.58 (0.12)				100	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.6 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)		% UL ³ >UL (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)					
Both															
	1-3	644	1.26 (0.03)	0.72 (0.07)	0.82 (0.06)	1.00 (0.05)	1.24 (0.04)	1.51 (0.05)	1.80 (0.09)	2.00 (0.12)	0.4	<3	30	0.0 (0.0)	
	4-8	956	1.48 (0.03)	0.93 (0.08)	1.03 (0.07)	1.22 (0.05)	1.46 (0.04)	1.72 (0.06)	1.99 (0.09)	2.17 (0.12)	0.5	<3	40	0.0 (0.0)	
Male															
	9-13	589	1.80 (0.05)	1.09 (0.11)	1.22 (0.10)	1.45 (0.08)	1.75 (0.06)	2.11 (0.09)	2.51 (0.15)	2.79 (0.21)	0.8	<3	60	0.0 (0.0)	
	14-18	639	2.10 (0.06)	1.37 (0.18)	1.51 (0.16)	1.76 (0.12)	2.08 (0.08)	2.44 (0.09)	2.81 (0.17)	3.04 (0.23)	1.1	F	80	0.0 (0.0)	
	19-30	481	2.28 (0.09)	1.18 (0.24) ^E	1.38 (0.21)	1.72 (0.16)	2.17 (0.12)	2.72 (0.16)	3.34 (0.29)	3.76 (0.41)	1.1	F	100	0.0 (0.0)	
	31-50	709	2.15 (0.07)	1.32 (0.21)	1.47 (0.19)	1.76 (0.13)	2.11 (0.08)	2.52 (0.12)	2.94 (0.22)	3.22 (0.31)	1.1	F	100	0.0 (0.0)	
	51-70	758	2.11 (0.05)	1.44 (0.18)	1.56 (0.15)	1.79 (0.10)	2.06 (0.07)	2.37 (0.11)	2.66 (0.19)	2.85 (0.25)	1.4	F	100	0.0 (0.0)	
	>70	734	1.78 (0.04)	1.10 (0.13)	1.23 (0.11)	1.46 (0.08)	1.74 (0.05)	2.05 (0.08)	2.37 (0.15)	2.58 (0.20)	1.4	20.6 (6.5) ^E	100	0.0 (0.0)	
	19+	2682	2.14 (0.04)	1.28 (0.10)	1.43 (0.09)	1.72 (0.07)	2.09 (0.05)	2.51 (0.07)	2.94 (0.11)	3.22 (0.16)			100	0.0 (0.0)	
Female															
	9-13	585	1.52 (0.05)	0.87 (0.10)	0.99 (0.09)	1.20 (0.06)	1.46 (0.05)	1.75 (0.08)	2.06 (0.14)	2.27 (0.19)	0.8	F	60	0.0 (0.0)	
	14-18	645	1.56 (0.05)	0.74 (0.07)	0.88 (0.07)	1.13 (0.05)	1.47 (0.05)	1.87 (0.07)	2.33 (0.12)	2.65 (0.17)	1.0	16.4 (3.5) ^E	80	0.0 (0.0)	
	19-30	514	1.52 (0.06)	0.80 (0.12)	0.92 (0.11)	1.14 (0.08)	1.44 (0.06)	1.78 (0.09)	2.13 (0.15)	2.36 (0.20)	1.1	21.8 (6.9) ^E	100	0.0 (0.0)	
	31-50	758	1.59 (0.04)	0.76 (0.05)	0.90 (0.05)	1.16 (0.05)	1.53 (0.05)	1.97 (0.07)	2.42 (0.10)	2.72 (0.13)	1.1	20.8 (3.0)	100	0.0 (0.0)	
	51-70	955	1.64 (0.05)	0.94 (0.11)	1.06 (0.10)	1.28 (0.08)	1.58 (0.06)	1.92 (0.08)	2.28 (0.13)	2.52 (0.18)	1.3	26.3 (6.7) ^E	100	0.0 (0.0)	
	>70	1345	1.54 (0.03)	0.84 (0.06)	0.95 (0.05)	1.17 (0.04)	1.46 (0.04)	1.79 (0.05)	2.13 (0.08)	2.35 (0.10)	1.3	35.7 (3.5)	100	0.0 (0.0)	
	19+	3572	1.58 (0.03)	0.84 (0.04)	0.96 (0.03)	1.20 (0.03)	1.52 (0.03)	1.89 (0.04)	2.28 (0.06)	2.55 (0.08)			100	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.7 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																
	1-3	324	1.20 (0.05)	0.72 (0.04)	0.80 (0.04)	0.96 (0.05)	1.18 (0.06)	1.44 (0.07)	1.71 (0.09)	1.89 (0.10)	0.4	<3		30	0.0 (0.0)	
	4-8	425	1.34 (0.07)	0.87 (0.06)	0.95 (0.06)	1.10 (0.07)	1.29 (0.08)	1.51 (0.08)	1.74 (0.09)	1.88 (0.10)	0.5	0.0 (0.0)		40	0.0 (0.0)	
Male																
	9-13	274	1.64 (0.08)	1.11 (0.15)	1.21 (0.13)	1.40 (0.11)	1.63 (0.10)	1.88 (0.12)	2.12 (0.18)	2.28 (0.23)	0.8	<3		60	0.0 (0.0)	
	14-18	297	2.07 (0.09)	1.00 (0.13)	1.18 (0.12)	1.54 (0.11)	2.00 (0.10)	2.55 (0.12)	3.13 (0.17)	3.51 (0.21)	1.1	F		80	0.0 (0.0)	
	19-30	249	2.24 (0.17)	1.45 (0.29) ^E	1.58 (0.26) ^E	1.82 (0.21)	2.11 (0.17)	2.44 (0.20)	2.77 (0.30)	2.98 (0.39)	1.1	F		100	0.0 (0.0)	
	31-50	309	1.92 (0.13)	1.18 (0.20)	1.30 (0.17)	1.53 (0.14)	1.82 (0.12)	2.16 (0.16)	2.53 (0.27)	2.78 (0.37)	1.1	F		100	0.0 (0.0)	
	51-70	277	1.88 (0.07)	1.12 (0.15)	1.26 (0.13)	1.51 (0.10)	1.83 (0.08)	2.17 (0.11)	2.52 (0.16)	2.74 (0.21)	1.4	F		100	0.0 (0.0)	
	>70	136	1.81 (0.13)	0.95 (0.13)	1.11 (0.13)	1.39 (0.12)	1.75 (0.14)	2.18 (0.18)	2.65 (0.25)	2.98 (0.33)	1.4	25.4 (8.3) ^E		100	0.0 (0.0)	
	19+	971	1.97 (0.06)	1.14 (0.12)	1.27 (0.11)	1.54 (0.09)	1.88 (0.07)	2.25 (0.09)	2.66 (0.14)	2.93 (0.19)				100	0.0 (0.0)	
Female																
	9-13	265	1.40 (0.08)	1.09 (0.19) ^E	1.15 (0.16)	1.25 (0.12)	1.37 (0.09)	1.51 (0.13)	1.63 (0.21)	1.71 (0.28)	0.8	F		60	0.0 (0.0)	
	14-18	290	1.40 (0.07)	0.70 (0.13) ^E	0.82 (0.11)	1.03 (0.09)	1.30 (0.09)	1.60 (0.14)	1.90 (0.20)	2.09 (0.25)	1.0	F		80	0.0 (0.0)	
	19-30	197	1.69 (0.14)	1.38 (0.30) ^E	1.45 (0.27) ^E	1.58 (0.21)	1.72 (0.17)	1.87 (0.20)	2.02 (0.30)	2.10 (0.38) ^E	1.1	F		100	0.0 (0.0)	
	31-50	312	1.61 (0.07)	0.95 (0.18) ^E	1.09 (0.15)	1.33 (0.11)	1.59 (0.08)	1.87 (0.11)	2.18 (0.19)	2.41 (0.26)	1.1	F		100	0.0 (0.0)	
	51-70	312	1.67 (0.08)	1.14 (0.17)	1.23 (0.16)	1.40 (0.13)	1.61 (0.10)	1.84 (0.11)	2.05 (0.16)	2.19 (0.20)	1.3	F		100	0.0 (0.0)	
	>70	239	1.38 (0.06)	0.63 (0.09)	0.75 (0.08)	0.97 (0.07)	1.29 (0.06)	1.68 (0.10)	2.11 (0.16)	2.41 (0.22)	1.3	50.6 (5.2)		100	0.0 (0.0)	
	19+	1060	1.61 (0.05)	0.88 (0.07)	1.00 (0.06)	1.24 (0.05)	1.55 (0.05)	1.91 (0.07)	2.27 (0.11)	2.51 (0.13)				100	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.8 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	129	1.16	(0.05)	0.76 (0.07)	0.85 (0.07)	0.99 (0.07)	1.17 (0.07)	1.35 (0.08)	1.55 (0.10)	1.69 (0.12)	0.4	<3	30	0.0	(0.0)
	4-8	213	1.39	(0.06)	0.90 (0.13)	0.99 (0.12)	1.14 (0.09)	1.35 (0.07)	1.58 (0.09)	1.83 (0.15)	1.99 (0.20)	0.5	<3	40	0.0	(0.0)
Male																
	9-13	122	1.65	(0.10)	1.25 (0.11)	1.33 (0.12)	1.47 (0.13)	1.64 (0.14)	1.83 (0.16)	2.01 (0.19)	2.14 (0.20)	0.8	<3	60	0.0	(0.0)
	14-18	150	2.22	(0.17)	1.18 (0.13)	1.36 (0.14)	1.70 (0.16)	2.14 (0.20)	2.66 (0.26)	3.21 (0.32)	3.57 (0.37)	1.1	F	80	0.0	(0.0)
	19-30	106	2.11	(0.12)	1.09 (0.25) ^E	1.28 (0.23) ^E	1.64 (0.18)	2.08 (0.15)	2.57 (0.20)	3.05 (0.32)	3.36 (0.41)	1.1	F	100	0.0	(0.0)
	31-50	155	1.93	(0.12)	0.97 (0.22) ^E	1.16 (0.20) ^E	1.51 (0.17)	1.93 (0.16)	2.41 (0.18)	2.94 (0.25)	3.31 (0.32)	1.1	F	100	0.0	(0.0)
	51-70	122	1.90	(0.14)	1.00 (0.25) ^E	1.16 (0.22) ^E	1.48 (0.18)	1.89 (0.18)	2.36 (0.28)	2.84 (0.42)	3.15 (0.52)	1.4	F	100	0.0	(0.0)
	>70	88	1.89	(0.11)	1.10 (0.18)	1.22 (0.17)	1.46 (0.15)	1.78 (0.13)	2.15 (0.16)	2.50 (0.21)	2.70 (0.25)	1.4	F	100	0.0	(0.0)
	19+	471	1.96	(0.06)	1.04 (0.10)	1.22 (0.09)	1.53 (0.08)	1.93 (0.08)	2.40 (0.11)	2.89 (0.15)	3.21 (0.18)			100	0.0	(0.0)
Female																
	9-13	103	1.47	(0.13)	0.94 (0.14)	1.03 (0.14)	1.18 (0.13)	1.38 (0.13)	1.60 (0.16)	1.83 (0.21)	1.98 (0.25)	0.8	F	60	0.0	(0.0)
	14-18	142	1.52	(0.09)	1.06 (0.15)	1.15 (0.14)	1.31 (0.12)	1.51 (0.12)	1.73 (0.16)	1.97 (0.22)	2.13 (0.26)	1.0	F	80	0.0	(0.0)
	19-30	111	1.32	(0.08)	0.84 (0.17) ^E	0.93 (0.15)	1.08 (0.12)	1.27 (0.11)	1.46 (0.14)	1.63 (0.19)	1.74 (0.23)	1.1	F	100	0.0	(0.0)
	31-50	146	1.68	(0.11)	0.80 (0.20) ^E	0.94 (0.19) ^E	1.20 (0.16)	1.54 (0.14)	1.93 (0.16)	2.31 (0.22)	2.56 (0.27)	1.1	F	100	0.0	(0.0)
	51-70	184	1.62	(0.09)	1.26 (0.08)	1.34 (0.09)	1.47 (0.10)	1.62 (0.12)	1.78 (0.14)	1.94 (0.16)	2.04 (0.18)	1.3	F	100	0.0	(0.0)
	>70	143	1.60	(0.07)	0.97 (0.15)	1.09 (0.13)	1.31 (0.11)	1.60 (0.10)	1.93 (0.13)	2.29 (0.19)	2.54 (0.25)	1.3	F	100	0.0	(0.0)
	19+	584	1.58	(0.06)	0.92 (0.10)	1.04 (0.10)	1.26 (0.08)	1.53 (0.07)	1.83 (0.08)	2.12 (0.11)	2.31 (0.14)			100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.9 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	169	1.13	(0.05)	0.60 (0.08)	0.69 (0.07)	0.87 (0.06)	1.10 (0.06)	1.38 (0.08)	1.66 (0.12)	1.85 (0.16)	0.4	<3	30	0.0	(0.0)
	4-8	281	1.35	(0.05)	0.81 (0.13)	0.89 (0.11)	1.06 (0.08)	1.27 (0.06)	1.53 (0.08)	1.80 (0.15)	1.98 (0.21)	0.5	<3	40	0.0	(0.0)
Male																
	9-13	183	1.87	(0.12)	1.18 (0.20) ^E	1.30 (0.19)	1.53 (0.16)	1.83 (0.15)	2.19 (0.18)	2.58 (0.27)	2.83 (0.35)	0.8	<3	60	0.0	(0.0)
	14-18	187	2.18	(0.11)	1.33 (0.26) ^E	1.50 (0.23)	1.82 (0.18)	2.21 (0.13)	2.67 (0.28)	3.14 (0.51)	3.46 (0.67) ^E	1.1	F	80	0.0	(0.0)
	19-30	223	2.14	(0.11)	1.17 (0.24) ^E	1.34 (0.21)	1.65 (0.17)	2.03 (0.13)	2.49 (0.32)	2.99 (0.69) ^E	3.34 (1.00) ^E	1.1	F	100	0.0	(0.0)
	31-50	229	2.29	(0.12)	1.23 (0.27) ^E	1.42 (0.25) ^E	1.79 (0.19)	2.28 (0.16)	2.85 (0.24)	3.46 (0.39)	3.88 (0.51)	1.1	F	100	0.0	(0.0)
	51-70	197	1.92	(0.10)	1.09 (0.18) ^E	1.24 (0.16)	1.50 (0.13)	1.83 (0.12)	2.19 (0.16)	2.55 (0.25)	2.78 (0.31)	1.4	F	100	0.0	(0.0)
	>70	72	1.84	(0.12)	1.11 (0.20) ^E	1.24 (0.18)	1.49 (0.16)	1.80 (0.15)	2.15 (0.18)	2.51 (0.23)	2.74 (0.28)	1.4	F	100	0.0	(0.0)
	19+	721	2.13	(0.07)	1.10 (0.08)	1.28 (0.08)	1.62 (0.07)	2.06 (0.08)	2.59 (0.10)	3.16 (0.15)	3.56 (0.19)			100	0.0	(0.0)
Female																
	9-13	165	1.58	(0.11)	0.90 (0.21) ^E	1.03 (0.19) ^E	1.28 (0.15)	1.60 (0.13)	1.96 (0.17)	2.33 (0.26)	2.57 (0.33)	0.8	F	60	0.0	(0.0)
	14-18	206	1.38	(0.07)	0.86 (0.13)	0.96 (0.11)	1.12 (0.09)	1.33 (0.08)	1.57 (0.11)	1.83 (0.18)	2.01 (0.23)	1.0	F	80	0.0	(0.0)
	19-30	191	1.53	(0.12)	0.81 (0.14) ^E	0.92 (0.13)	1.13 (0.12)	1.40 (0.11)	1.78 (0.16)	2.26 (0.29)	2.63 (0.43)	1.1	F	100	0.0	(0.0)
	31-50	258	1.57	(0.08)	1.04 (0.18) ^E	1.16 (0.16)	1.38 (0.14)	1.63 (0.11)	1.88 (0.12)	2.14 (0.18)	2.31 (0.25)	1.1	F	100	0.0	(0.0)
	51-70	249	1.55	(0.08)	1.05 (0.18) ^E	1.15 (0.15)	1.33 (0.11)	1.55 (0.09)	1.79 (0.12)	2.04 (0.18)	2.20 (0.23)	1.3	F	100	0.0	(0.0)
	>70	128	1.66	(0.11)	1.02 (0.12)	1.15 (0.12)	1.39 (0.12)	1.67 (0.13)	1.95 (0.14)	2.22 (0.17)	2.39 (0.20)	1.3	F	100	0.0	(0.0)
	19+	826	1.57	(0.05)	0.94 (0.07)	1.06 (0.07)	1.27 (0.06)	1.55 (0.06)	1.88 (0.07)	2.24 (0.11)	2.50 (0.15)			100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.10 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		UL ³	% >UL	
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)	(<i>SE</i>)		(<i>SE</i>)	
Both																
	1-3	192	1.24	(0.05)	0.83 (0.08)	0.91 (0.07)	1.04 (0.06)	1.19 (0.06)	1.35 (0.08)	1.50 (0.10)	1.60 (0.13)	0.4	<3	30	0.0	(0.0)
	4-8	321	1.58	(0.05)	1.04 (0.11)	1.15 (0.10)	1.34 (0.07)	1.57 (0.06)	1.84 (0.09)	2.10 (0.13)	2.27 (0.16)	0.5	<3	40	0.0	(0.0)
Male																
	9-13	226	1.73	(0.07)	1.00 (0.09)	1.12 (0.08)	1.35 (0.08)	1.64 (0.08)	1.99 (0.12)	2.40 (0.18)	2.70 (0.24)	0.8	<3	60	0.0	(0.0)
	14-18	262	2.61	(0.29)	1.33 (0.33) ^E	1.53 (0.31) ^E	1.90 (0.29)	2.43 (0.29)	3.18 (0.36)	4.04 (0.58)	4.64 (0.78) ^E	1.1	F	80	0.0	(0.0)
	19-30	197	2.45	(0.12)	1.50 (0.16)	1.67 (0.15)	1.98 (0.14)	2.40 (0.14)	2.93 (0.20)	3.53 (0.34)	3.96 (0.46)	1.1	<3	100	0.0	(0.0)
	31-50	282	2.62	(0.18)	1.58 (0.22)	1.76 (0.20)	2.08 (0.18)	2.52 (0.18)	3.06 (0.26)	3.67 (0.41)	4.11 (0.55)	1.1	<3	100	0.0	(0.0)
	51-70	234	2.31	(0.10)	1.29 (0.16)	1.46 (0.15)	1.79 (0.14)	2.21 (0.14)	2.69 (0.16)	3.19 (0.22)	3.51 (0.27)	1.4	F	100	0.0	(0.0)
	>70	119	2.01	(0.15)	0.90 (0.16) ^E	1.07 (0.15)	1.40 (0.16)	1.86 (0.19)	2.44 (0.24)	3.05 (0.33)	3.49 (0.42)	1.4	F	100	0.0	(0.0)
	19+	832	2.44	(0.08)	1.33 (0.07)	1.52 (0.07)	1.86 (0.07)	2.33 (0.08)	2.91 (0.12)	3.57 (0.19)	4.06 (0.27)			100	0.0	(0.0)
Female																
	9-13	226	1.63	(0.10)	1.01 (0.13)	1.13 (0.12)	1.35 (0.11)	1.63 (0.11)	1.96 (0.14)	2.30 (0.20)	2.53 (0.24)	0.8	F	60	0.0	(0.0)
	14-18	242	1.55	(0.09)	0.89 (0.13)	1.00 (0.12)	1.22 (0.11)	1.49 (0.11)	1.82 (0.14)	2.16 (0.19)	2.38 (0.22)	1.0	F	80	0.0	(0.0)
	19-30	208	1.65	(0.10)	1.09 (0.08)	1.19 (0.09)	1.39 (0.10)	1.65 (0.12)	1.96 (0.15)	2.28 (0.19)	2.50 (0.23)	1.1	F	100	0.0	(0.0)
	31-50	263	1.78	(0.08)	0.90 (0.13)	1.04 (0.12)	1.30 (0.11)	1.64 (0.11)	2.00 (0.13)	2.38 (0.18)	2.64 (0.24)	1.1	F	100	0.0	(0.0)
	51-70	322	1.73	(0.11)	1.12 (0.18)	1.24 (0.16)	1.45 (0.13)	1.71 (0.12)	2.01 (0.16)	2.32 (0.25)	2.52 (0.32)	1.3	F	100	0.0	(0.0)
	>70	198	1.66	(0.10)	0.98 (0.17) ^E	1.11 (0.16)	1.34 (0.14)	1.64 (0.14)	1.96 (0.17)	2.25 (0.21)	2.43 (0.24)	1.3	F	100	0.0	(0.0)
	19+	991	1.73	(0.05)	0.97 (0.08)	1.09 (0.08)	1.34 (0.07)	1.68 (0.07)	2.05 (0.09)	2.42 (0.12)	2.67 (0.15)			100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.11 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)		UL ³	% >UL (SE)								
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)					(SE)			(SE)					
Both																								
	1-3	348	1.29	(0.05)	0.91	(0.10)	0.98	(0.09)	1.12	(0.07)	1.28	(0.06)	1.46	(0.07)	1.64	(0.10)	1.76	(0.12)	0.4	0.0	(0.0)	30	0.0	(0.0)
	4-8	554	1.52	(0.06)	0.96	(0.06)	1.06	(0.06)	1.25	(0.06)	1.50	(0.07)	1.81	(0.10)	2.14	(0.15)	2.38	(0.20)	0.5	<3		40	0.0	(0.0)
Male																								
	9-13	409	1.75	(0.07)	1.10	(0.09)	1.21	(0.08)	1.42	(0.08)	1.68	(0.08)	1.97	(0.10)	2.28	(0.13)	2.49	(0.16)	0.8	<3		60	0.0	(0.0)
	14-18	414	2.01	(0.08)	1.29	(0.13)	1.41	(0.12)	1.65	(0.11)	1.98	(0.11)	2.39	(0.15)	2.82	(0.22)	3.11	(0.28)	1.1	F		80	0.0	(0.0)
	19-30	311	2.29	(0.13)	1.38	(0.14)	1.54	(0.14)	1.83	(0.13)	2.20	(0.14)	2.63	(0.17)	3.09	(0.24)	3.40	(0.30)	1.1	<3		100	0.0	(0.0)
	31-50	489	2.10	(0.08)	1.27	(0.13)	1.41	(0.12)	1.68	(0.11)	2.04	(0.10)	2.47	(0.13)	2.91	(0.18)	3.19	(0.22)	1.1	F		100	0.0	(0.0)
	51-70	575	1.98	(0.07)	1.17	(0.12)	1.32	(0.10)	1.58	(0.09)	1.91	(0.08)	2.28	(0.10)	2.64	(0.13)	2.88	(0.17)	1.4	F		100	0.0	(0.0)
	>70	239	1.83	(0.09)	1.06	(0.10)	1.19	(0.10)	1.44	(0.11)	1.76	(0.12)	2.14	(0.14)	2.55	(0.19)	2.82	(0.22)	1.4	22.4	(7.1) ^E	100	0.0	(0.0)
	19+	1614	2.08	(0.05)	1.19	(0.06)	1.34	(0.05)	1.63	(0.05)	2.01	(0.06)	2.46	(0.07)	2.93	(0.10)	3.24	(0.12)				100	0.0	(0.0)
Female																								
	9-13	355	1.52	(0.08)	1.09	(0.13)	1.18	(0.12)	1.33	(0.10)	1.53	(0.10)	1.76	(0.12)	2.00	(0.16)	2.16	(0.19)	0.8	<3		60	0.0	(0.0)
	14-18	410	1.41	(0.08)	0.84	(0.10)	0.94	(0.10)	1.13	(0.09)	1.38	(0.09)	1.67	(0.12)	1.96	(0.17)	2.16	(0.22)	1.0	F		80	0.0	(0.0)
	19-30	384	1.49	(0.06)	1.01	(0.11)	1.11	(0.10)	1.30	(0.08)	1.52	(0.08)	1.77	(0.10)	2.02	(0.13)	2.16	(0.15)	1.1	F		100	0.0	(0.0)
	31-50	585	1.49	(0.05)	0.75	(0.07)	0.89	(0.07)	1.15	(0.06)	1.47	(0.07)	1.81	(0.08)	2.15	(0.10)	2.39	(0.13)	1.1	21.4	(4.4) ^E	100	0.0	(0.0)
	51-70	711	1.55	(0.05)	0.91	(0.07)	1.02	(0.07)	1.23	(0.06)	1.49	(0.06)	1.79	(0.08)	2.09	(0.11)	2.29	(0.13)	1.3	31.5	(5.9) ^E	100	0.0	(0.0)
	>70	342	1.58	(0.06)	0.95	(0.08)	1.08	(0.08)	1.30	(0.07)	1.58	(0.08)	1.88	(0.09)	2.18	(0.12)	2.37	(0.14)	1.3	24.8	(6.1) ^E	100	0.0	(0.0)
	19+	2022	1.52	(0.03)	0.86	(0.04)	0.99	(0.04)	1.22	(0.04)	1.50	(0.04)	1.82	(0.04)	2.12	(0.06)	2.33	(0.07)				100	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.12 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR (SE)		UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	622	1.15	(0.03)	0.70 (0.05)	0.78 (0.04)	0.94 (0.04)	1.14 (0.04)	1.37 (0.05)	1.62 (0.07)	1.79 (0.09)	0.4	<3	30	0.0 (0.0)	
	4-8	919	1.36	(0.03)	0.86 (0.08)	0.94 (0.07)	1.10 (0.05)	1.30 (0.04)	1.53 (0.06)	1.76 (0.10)	1.92 (0.13)	0.5	<3	40	0.0 (0.0)	
Male																
	9-13	579	1.78	(0.08)	1.31 (0.17)	1.40 (0.15)	1.57 (0.12)	1.77 (0.10)	2.01 (0.12)	2.24 (0.17)	2.39 (0.22)	0.8	<3	60	0.0 (0.0)	
	14-18	634	2.17	(0.08)	1.20 (0.10)	1.38 (0.09)	1.72 (0.09)	2.15 (0.09)	2.65 (0.12)	3.19 (0.17)	3.55 (0.22)	1.1	F	80	0.0 (0.0)	
	19-30	578	2.15	(0.08)	1.12 (0.15)	1.30 (0.14)	1.63 (0.12)	2.04 (0.10)	2.54 (0.12)	3.06 (0.19)	3.42 (0.25)	1.1	F	100	0.0 (0.0)	
	31-50	693	2.16	(0.09)	1.17 (0.15)	1.34 (0.14)	1.69 (0.12)	2.14 (0.11)	2.62 (0.13)	3.19 (0.21)	3.59 (0.28)	1.1	F	100	0.0 (0.0)	
	51-70	596	1.91	(0.07)	1.03 (0.10)	1.19 (0.09)	1.48 (0.08)	1.84 (0.08)	2.26 (0.10)	2.68 (0.14)	2.96 (0.17)	1.4	20.4 (4.9) ^E	100	0.0 (0.0)	
	>70	296	1.84	(0.07)	1.09 (0.13)	1.22 (0.12)	1.46 (0.10)	1.78 (0.09)	2.13 (0.11)	2.49 (0.15)	2.72 (0.19)	1.4	F	100	0.0 (0.0)	
	19+	2163	2.07	(0.04)	1.11 (0.06)	1.28 (0.05)	1.59 (0.05)	2.00 (0.05)	2.49 (0.07)	3.01 (0.09)	3.36 (0.12)			100	0.0 (0.0)	
Female																
	9-13	533	1.52	(0.07)	1.02 (0.13)	1.12 (0.12)	1.30 (0.10)	1.51 (0.08)	1.76 (0.11)	2.00 (0.16)	2.15 (0.20)	0.8	F	60	0.0 (0.0)	
	14-18	638	1.41	(0.05)	0.84 (0.07)	0.95 (0.06)	1.13 (0.06)	1.35 (0.06)	1.62 (0.07)	1.89 (0.10)	2.08 (0.13)	1.0	13.7 (4.3) ^E	80	0.0 (0.0)	
	19-30	499	1.53	(0.08)	0.95 (0.12)	1.05 (0.11)	1.22 (0.09)	1.46 (0.08)	1.75 (0.11)	2.07 (0.18)	2.30 (0.24)	1.1	F	100	0.0 (0.0)	
	31-50	716	1.60	(0.05)	0.93 (0.12)	1.05 (0.11)	1.29 (0.09)	1.61 (0.07)	1.91 (0.08)	2.24 (0.12)	2.47 (0.16)	1.1	F	100	0.0 (0.0)	
	51-70	745	1.59	(0.05)	1.24 (0.14)	1.31 (0.12)	1.44 (0.09)	1.59 (0.06)	1.75 (0.08)	1.90 (0.13)	2.00 (0.17)	1.3	F	100	0.0 (0.0)	
	>70	510	1.57	(0.06)	0.81 (0.06)	0.95 (0.06)	1.21 (0.06)	1.55 (0.07)	1.92 (0.08)	2.31 (0.10)	2.57 (0.13)	1.3	31.2 (5.1)	100	0.0 (0.0)	
	19+	2470	1.58	(0.03)	0.93 (0.05)	1.04 (0.05)	1.26 (0.04)	1.55 (0.04)	1.88 (0.05)	2.23 (0.07)	2.47 (0.09)			100	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 24.13 Vitamin B₆ (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR (SE)	UL ³	% >UL (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)				
Both														
	1-3	2117	1.26 (0.02)	0.75 (0.03)	0.85 (0.03)	1.01 (0.03)	1.23 (0.03)	1.49 (0.03)	1.77 (0.05)	1.95 (0.06)	0.4	<3	30	0.0 (0.0)
	4-8	3235	1.49 (0.02)	0.94 (0.03)	1.04 (0.03)	1.22 (0.03)	1.46 (0.03)	1.73 (0.03)	2.01 (0.05)	2.19 (0.06)	0.5	<3	40	0.0 (0.0)
Male														
	9-13	2080	1.83 (0.04)	1.16 (0.06)	1.28 (0.05)	1.50 (0.05)	1.78 (0.05)	2.11 (0.06)	2.45 (0.08)	2.69 (0.10)	0.8	<3	60	0.0 (0.0)
	14-18	2288	2.22 (0.05)	1.32 (0.06)	1.48 (0.06)	1.78 (0.05)	2.19 (0.06)	2.69 (0.07)	3.23 (0.11)	3.60 (0.15)	1.1	<3	80	0.0 (0.0)
	19-30	1804	2.29 (0.06)	1.31 (0.07)	1.48 (0.07)	1.79 (0.06)	2.22 (0.07)	2.74 (0.09)	3.31 (0.13)	3.70 (0.17)	1.1	F	100	0.0 (0.0)
	31-50	2596	2.23 (0.05)	1.28 (0.06)	1.46 (0.06)	1.78 (0.05)	2.17 (0.05)	2.65 (0.07)	3.18 (0.11)	3.54 (0.15)	1.1	F	100	0.0 (0.0)
	51-70	2550	2.08 (0.04)	1.23 (0.06)	1.38 (0.05)	1.66 (0.05)	2.03 (0.05)	2.46 (0.05)	2.89 (0.08)	3.16 (0.09)	1.4	10.9 (2.2) ^E	100	0.0 (0.0)
	>70	1520	1.83 (0.04)	1.02 (0.05)	1.16 (0.05)	1.43 (0.05)	1.78 (0.06)	2.19 (0.07)	2.61 (0.08)	2.89 (0.10)	1.4	23.1 (3.4)	100	0.0 (0.0)
	19+	8470	2.17 (0.03)	1.23 (0.03)	1.39 (0.03)	1.71 (0.03)	2.11 (0.03)	2.59 (0.04)	3.11 (0.06)	3.47 (0.08)			100	0.0 (0.0)
Female														
	9-13	1980	1.54 (0.03)	0.93 (0.04)	1.04 (0.04)	1.24 (0.03)	1.49 (0.04)	1.79 (0.05)	2.09 (0.07)	2.29 (0.09)	0.8	F	60	0.0 (0.0)
	14-18	2256	1.51 (0.03)	0.86 (0.03)	0.98 (0.03)	1.19 (0.03)	1.47 (0.03)	1.79 (0.04)	2.13 (0.06)	2.37 (0.07)	1.0	11.1 (1.7)	80	0.0 (0.0)
	19-30	1854	1.59 (0.04)	1.00 (0.06)	1.11 (0.06)	1.31 (0.05)	1.56 (0.05)	1.85 (0.06)	2.14 (0.08)	2.32 (0.10)	1.1	9.6 (3.2) ^E	100	0.0 (0.0)
	31-50	2686	1.65 (0.03)	0.85 (0.03)	0.98 (0.03)	1.25 (0.04)	1.60 (0.04)	2.00 (0.05)	2.40 (0.07)	2.68 (0.08)	1.1	15.9 (1.9)	100	0.0 (0.0)
	51-70	3200	1.66 (0.03)	1.05 (0.05)	1.16 (0.05)	1.37 (0.04)	1.63 (0.04)	1.93 (0.04)	2.24 (0.07)	2.44 (0.09)	1.3	19.4 (3.4) ^E	100	0.0 (0.0)
	>70	2610	1.55 (0.03)	0.87 (0.04)	0.99 (0.04)	1.21 (0.04)	1.50 (0.04)	1.85 (0.04)	2.20 (0.05)	2.44 (0.06)	1.3	32.5 (3.1)	100	0.0 (0.0)
	19+	10350	1.63 (0.02)	0.92 (0.02)	1.05 (0.02)	1.28 (0.02)	1.59 (0.02)	1.94 (0.03)	2.30 (0.04)	2.55 (0.05)			100	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

25. Vitamin B₁₂ (µg/d): Usual intakes from food

Table 25.1 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	3.5 (0.4)	1.9 (0.4) ^E	2.2 (0.4) ^E	2.8 (0.4)	3.5 (0.4)	4.4 (0.5)	5.3 (0.8)	6.0 (1.1) ^E	0.7	<3	
	4-8	127	3.3 (0.2)	2.3 (0.2)	2.4 (0.2)	2.8 (0.3)	3.2 (0.3)	3.6 (0.3)	4.0 (0.3)	4.3 (0.4)	1.0	0.0 (0.0)	
Male	9-13	111	4.8 (0.6)	2.7 (0.6) ^E	3.0 (0.6) ^E	3.7 (0.6)	4.6 (0.6)	5.7 (0.9)	7.0 (1.2) ^E	7.8 (1.6) ^E	1.5	<3	
	14-18	107	4.7 (0.4)	2.7 (0.3)	3.1 (0.3)	3.8 (0.4)	4.7 (0.5)	5.9 (0.6)	7.0 (0.7)	7.8 (0.8)	2.0	<3	
	19-30	77	3.8 (0.3)	2.2 (0.6) ^E	2.4 (0.6) ^E	2.9 (0.5)	3.6 (0.4)	4.3 (0.5)	5.1 (0.8)	5.6 (1.2) ^E	2.0	F	
	31-50	145	6.5 (2.0) ^E	3.3 (0.4)	3.6 (0.5)	4.0 (0.6)	4.6 (0.7)	5.2 (0.9) ^E	6.0 (1.3) ^E	6.5 (1.6) ^E	2.0	<3	
	51-70	182	4.2 (0.4)	2.3 (0.3)	2.6 (0.3)	3.1 (0.3)	3.8 (0.4)	4.7 (0.6)	5.6 (0.7)	6.3 (0.8)	2.0	F	
	>70	63	6.2 (1.9) ^E	F	F	F	F	F	F	F	2.0	F	
	19+	467	5.2 (0.8)	2.5 (0.4) ^E	2.8 (0.4)	3.3 (0.4)	4.2 (0.4)	5.3 (0.5)	6.7 (1.0)	7.8 (1.5) ^E	2.0	F	
	Female	9-13	96	2.9 (0.2)	F	1.3 (0.4) ^E	1.9 (0.3) ^E	2.7 (0.3)	3.7 (0.4)	4.8 (0.6)	5.6 (0.7)	1.5	F
14-18		105	3.1 (0.4)	1.5 (0.2)	1.7 (0.3)	2.2 (0.3)	2.8 (0.4)	3.6 (0.6)	4.7 (0.8) ^E	5.5 (1.0) ^E	2.0	F	
19-30		91	4.1 (1.2) ^E	1.8 (0.3) ^E	2.0 (0.4) ^E	2.6 (0.5) ^E	3.4 (0.8) ^E	4.7 (1.4) ^E	F	F	2.0	F	
31-50		167	3.0 (0.4)	1.2 (0.3) ^E	1.5 (0.3) ^E	2.0 (0.4) ^E	2.7 (0.4)	3.7 (0.6)	4.9 (0.9) ^E	5.8 (1.1) ^E	2.0	F	
51-70		198	F	1.8 (0.3) ^E	2.1 (0.4) ^E	2.9 (0.9) ^E	F	F	F	F	2.0	F	
>70		74	3.0 (0.3)	1.7 (0.2)	1.9 (0.2)	2.2 (0.3)	2.6 (0.3)	3.1 (0.4)	3.6 (0.5)	4.0 (0.6)	2.0	F	
19+		530	4.6 (1.0) ^E	1.8 (0.4) ^E	2.0 (0.4) ^E	2.5 (0.4) ^E	3.4 (0.5)	5.0 (0.9) ^E	7.6 (2.1) ^E	F	2.0	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.2 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	58	3.6 (0.4)	2.2 (0.4) ^E	2.4 (0.4)	2.9 (0.4)	3.5 (0.4)	4.2 (0.6)	5.0 (0.8)	5.5 (1.0) ^E	0.7	<3
	4-8	110	3.9 (0.4)	2.5 (0.4) ^E	2.7 (0.4)	3.2 (0.4)	3.7 (0.3)	4.3 (0.4)	4.9 (0.5)	5.2 (0.6)	1.0	<3
Male	9-13	95	5.1 (1.3) ^E	2.3 (0.4) ^E	2.6 (0.5) ^E	3.3 (0.7) ^E	4.6 (1.1) ^E	F	F	F	1.5	<3
	14-18	87	5.5 (0.6)	2.5 (0.7) ^E	3.0 (0.7) ^E	3.9 (0.7) ^E	5.3 (0.8)	7.2 (1.1)	9.5 (1.7) ^E	11.2 (2.3) ^E	2.0	F
	19-30	70	8.4 (2.4) ^E	4.0 (0.6)	4.4 (0.7)	5.5 (1.1) ^E	7.7 (2.2) ^E	F	F	F	2.0	<3
	31-50	109	8.1 (2.6) ^E	2.2 (0.5) ^E	2.7 (0.6) ^E	3.8 (0.9) ^E	5.8 (1.7) ^E	F	F	F	2.0	F
	51-70	128	4.6 (0.5)	2.5 (0.5) ^E	2.8 (0.5) ^E	3.3 (0.5)	4.0 (0.4)	4.9 (0.5)	6.0 (0.9)	6.7 (1.2) ^E	2.0	F
	>70	65	3.7 (0.4)	1.5 (0.4) ^E	1.7 (0.4) ^E	2.3 (0.4) ^E	3.1 (0.5)	4.3 (0.7)	5.7 (1.1) ^E	6.8 (1.6) ^E	2.0	F
	19+	372	6.7 (1.1) ^E	2.4 (0.5) ^E	2.8 (0.5) ^E	3.7 (0.6) ^E	5.1 (0.9) ^E	7.7 (1.4) ^E	12.0 (2.9) ^E	16.2 (4.7) ^E	2.0	F
	Female	9-13	75	3.5 (0.4)	2.3 (0.5) ^E	2.5 (0.5) ^E	2.8 (0.5) ^E	3.3 (0.5) ^E	3.8 (0.7) ^E	4.3 (0.8) ^E	4.6 (1.0) ^E	1.5
14-18	81	3.5 (0.5)	F	F	2.0 (0.6) ^E	2.9 (0.7) ^E	4.2 (0.9) ^E	5.8 (1.4) ^E	7.1 (2.1) ^E	2.0	F	
19-30	101	5.7 (1.5) ^E	2.3 (0.7) ^E	2.8 (0.8) ^E	3.9 (0.9) ^E	5.4 (1.3) ^E	7.5 (2.0) ^E	10.1 (3.0) ^E	12.1 (3.9) ^E	2.0	F	
31-50	116	3.6 (0.4)	1.4 (0.3) ^E	1.7 (0.3) ^E	2.3 (0.4)	3.3 (0.5)	4.8 (0.7)	6.5 (1.3) ^E	8.0 (1.9) ^E	2.0	F	
51-70	146	3.7 (0.3)	1.4 (0.2) ^E	1.7 (0.3)	2.4 (0.3)	3.3 (0.3)	4.6 (0.4)	6.3 (0.7)	7.7 (0.9)	2.0	F	
>70	94	F	F	F	F	3.3 (1.1) ^E	5.1 (1.6) ^E	F	F	2.0	F	
19+	457	4.2 (0.4)	1.6 (0.2)	1.9 (0.2)	2.6 (0.2)	3.6 (0.4)	5.3 (0.6)	7.5 (1.0)	9.3 (1.4)	2.0	11.8 (3.6) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.3 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both												
	1-3	112	3.0 (0.2)	2.1 (0.4) ^E	2.2 (0.4) ^E	2.5 (0.3)	2.8 (0.3)	3.2 (0.3)	3.6 (0.5)	3.8 (0.6)	0.7	<3
	4-8	177	3.9 (0.2)	2.2 (0.5) ^E	2.5 (0.4)	3.0 (0.3)	3.7 (0.3)	4.5 (0.4)	5.3 (0.7)	5.9 (0.8)	1.0	<3
Male												
	9-13	111	4.5 (0.5)	1.8 (0.6) ^E	2.2 (0.5) ^E	2.9 (0.5)	3.9 (0.5)	5.2 (0.6)	7.0 (1.1)	8.5 (1.6) ^E	1.5	F
	14-18	113	5.3 (0.7)	2.6 (0.8) ^E	3.0 (0.8) ^E	3.7 (0.8) ^E	4.8 (1.0) ^E	6.2 (1.3) ^E	7.8 (1.7) ^E	9.0 (2.0) ^E	2.0	F
	19-30	91	5.5 (0.7)	F	3.1 (0.9) ^E	3.9 (0.8) ^E	5.0 (0.8)	6.3 (1.0)	7.7 (1.6) ^E	8.6 (2.1) ^E	2.0	F
	31-50	101	4.9 (0.5)	3.3 (0.8) ^E	3.6 (0.7) ^E	4.2 (0.7)	5.0 (0.7)	5.9 (1.1) ^E	7.0 (1.7) ^E	7.7 (2.4) ^E	2.0	F
	51-70	134	5.0 (0.6)	2.8 (0.6) ^E	3.2 (0.6) ^E	3.8 (0.6)	4.7 (0.6)	5.9 (0.7)	7.3 (1.2)	8.3 (1.7) ^E	2.0	F
	>70	56	4.7 (0.5)	2.2 (0.7) ^E	2.6 (0.7) ^E	3.4 (0.7) ^E	4.5 (0.7)	6.1 (1.0) ^E	8.2 (1.9) ^E	9.8 (2.8) ^E	2.0	F
	19+	382	5.0 (0.3)	2.4 (0.3)	2.8 (0.3)	3.5 (0.3)	4.6 (0.4)	6.1 (0.5)	8.0 (0.9)	9.4 (1.2)	2.0	F
Female												
	9-13	105	3.2 (0.4)	1.7 (0.4) ^E	1.9 (0.3) ^E	2.2 (0.3)	2.8 (0.3)	3.4 (0.4)	4.1 (0.6)	4.6 (0.8) ^E	1.5	F
	14-18	120	3.2 (0.5)	1.5 (0.5) ^E	1.7 (0.5) ^E	2.2 (0.4) ^E	2.9 (0.5) ^E	3.9 (0.8) ^E	F	F	2.0	F
	19-30	91	3.4 (0.3)	2.3 (0.5) ^E	2.5 (0.5) ^E	2.9 (0.5) ^E	3.4 (0.5)	4.0 (0.5)	4.6 (0.7)	4.9 (0.9) ^E	2.0	F
	31-50	159	4.2 (0.3)	1.9 (0.3)	2.3 (0.3)	3.1 (0.4)	4.2 (0.5)	5.5 (0.6)	6.9 (0.7)	7.8 (0.9)	2.0	F
	51-70	174	4.3 (0.8) ^E	1.9 (0.6) ^E	2.2 (0.6) ^E	2.8 (0.6) ^E	3.7 (0.7) ^E	5.0 (1.0) ^E	6.8 (1.8) ^E	F	2.0	F
	>70	80	5.0 (1.6) ^E	F	F	F	F	F	F	F	2.0	F
	19+	504	4.2 (0.4)	2.0 (0.3)	2.3 (0.3)	3.0 (0.3)	3.9 (0.3)	5.3 (0.6)	7.0 (1.1)	8.4 (1.7) ^E	2.0	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.4 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	99	F	2.9 (0.6) ^E	3.2 (0.7) ^E	4.0 (1.1) ^E	5.5 (1.8) ^E	F	F	F	0.7	0.0 (0.0)	
	4-8	140	F	2.7 (0.2)	2.8 (0.3)	3.1 (0.4)	3.7 (0.7) ^E	4.7 (1.3) ^E	F	F	1.0	0.0 (0.0)	
Male	9-13	92	5.0 (0.6)	2.5 (0.7) ^E	2.9 (0.6) ^E	3.4 (0.6) ^E	4.2 (0.6)	5.1 (0.7)	6.1 (1.0)	6.8 (1.2) ^E	1.5	<3	
	14-18	107	5.4 (0.6)	3.5 (0.7) ^E	3.8 (0.7) ^E	4.4 (0.6)	5.1 (0.7)	6.0 (0.9)	6.8 (1.2) ^E	7.4 (1.5) ^E	2.0	<3	
	19-30	73	5.6 (0.6)	3.7 (0.9) ^E	4.1 (0.8) ^E	4.8 (0.8)	5.7 (0.8)	6.8 (1.2) ^E	7.9 (1.8) ^E	8.7 (2.4) ^E	2.0	F	
	31-50	134	5.5 (1.1) ^E	3.4 (0.5)	3.7 (0.6)	4.3 (0.7) ^E	5.0 (1.0) ^E	6.0 (1.3) ^E	7.1 (1.8) ^E	7.9 (2.2) ^E	2.0	<3	
	51-70	131	6.2 (0.9)	3.9 (1.1) ^E	4.3 (1.0) ^E	4.9 (1.0) ^E	5.8 (1.0) ^E	6.9 (1.2) ^E	8.0 (1.9) ^E	8.9 (2.7) ^E	2.0	F	
	>70	55	4.1 (0.5)	F	2.3 (0.7) ^E	2.9 (0.6) ^E	3.7 (0.7) ^E	4.8 (0.9) ^E	6.2 (1.7) ^E	F	2.0	F	
	19+	393	5.6 (0.6)	3.1 (0.9) ^E	3.5 (0.8) ^E	4.2 (0.8) ^E	5.3 (0.8)	6.7 (1.0)	8.4 (1.5) ^E	9.6 (2.0) ^E	2.0	<3	
Female	9-13	79	3.9 (0.4)	F	2.1 (0.6) ^E	2.7 (0.5) ^E	3.6 (0.5)	4.7 (0.7)	6.2 (1.5) ^E	F	1.5	F	
	14-18	104	4.1 (0.9) ^E	1.3 (0.4) ^E	1.6 (0.4) ^E	2.2 (0.5) ^E	3.1 (0.6) ^E	4.5 (1.0) ^E	6.3 (1.7) ^E	7.8 (2.4) ^E	2.0	F	
	19-30	101	3.5 (0.6) ^E	F	2.1 (0.6) ^E	2.7 (0.6) ^E	3.6 (0.8) ^E	5.0 (1.3) ^E	F	F	2.0	F	
	31-50	143	F	F	F	F	F	F	F	F	2.0	F	
	51-70	193	4.5 (0.8) ^E	1.6 (0.2)	1.8 (0.2)	2.3 (0.3)	3.3 (0.6) ^E	5.3 (1.2) ^E	8.7 (2.7) ^E	F	2.0	F	
	>70	94	4.1 (0.9) ^E	2.4 (0.7) ^E	2.6 (0.7) ^E	3.0 (0.7) ^E	3.6 (0.8) ^E	4.3 (1.2) ^E	F	F	2.0	F	
	19+	531	4.6 (0.8)	1.8 (0.5) ^E	2.0 (0.5) ^E	2.6 (0.6) ^E	3.6 (0.8) ^E	5.4 (1.1) ^E	8.3 (1.9) ^E	11.0 (2.9) ^E	2.0	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.5 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	311	3.6 (0.2)	2.0 (0.3)	2.3 (0.3)	2.8 (0.3)	3.5 (0.2)	4.3 (0.3)	5.1 (0.3)	5.7 (0.4)	0.7	<3	
	4-8	485	4.0 (0.2)	2.2 (0.4) ^E	2.5 (0.4)	3.0 (0.3)	3.8 (0.3)	4.8 (0.4)	5.9 (0.7)	6.5 (0.9)	1.0	<3	
Male	9-13	277	4.6 (0.3)	3.1 (0.5)	3.4 (0.5)	3.9 (0.4)	4.7 (0.4)	5.5 (0.5)	6.4 (0.7)	7.0 (0.8)	1.5	<3	
	14-18	339	5.9 (0.5)	2.8 (0.6) ^E	3.4 (0.5)	4.4 (0.5)	5.9 (0.6)	7.7 (0.8)	9.9 (1.2)	11.4 (1.6)	2.0	F	
	19-30	237	5.4 (0.8)	2.2 (0.7) ^E	2.6 (0.7) ^E	3.4 (0.7) ^E	4.9 (0.8) ^E	7.1 (1.5) ^E	10.1 (3.0) ^E	F	2.0	F	
	31-50	423	5.1 (0.6)	2.4 (0.7) ^E	2.7 (0.6) ^E	3.4 (0.6) ^E	4.4 (0.5)	5.8 (0.6)	7.6 (1.1)	8.9 (1.6) ^E	2.0	F	
	51-70	387	5.3 (0.4)	2.1 (0.5) ^E	2.4 (0.5) ^E	3.1 (0.5)	4.3 (0.5)	6.2 (0.6)	9.3 (1.5)	12.3 (2.8) ^E	2.0	F	
	>70	132	4.9 (1.0) ^E	1.6 (0.5) ^E	1.9 (0.5) ^E	2.6 (0.6) ^E	3.9 (0.8) ^E	6.0 (1.5) ^E	9.3 (3.1) ^E	F	2.0	F	
	19+	1179	5.2 (0.3)	2.2 (0.3)	2.5 (0.2)	3.2 (0.3)	4.3 (0.3)	6.0 (0.4)	8.6 (0.8)	10.8 (1.3)	2.0	F	
	Female	9-13	281	3.6 (0.2)	2.3 (0.4) ^E	2.5 (0.4)	2.9 (0.3)	3.4 (0.3)	3.9 (0.4)	4.5 (0.5)	4.8 (0.7)	1.5	<3
14-18	321	3.4 (0.2)	2.0 (0.3)	2.2 (0.3)	2.6 (0.3)	3.3 (0.2)	4.0 (0.3)	4.9 (0.5)	5.5 (0.7)	2.0	F		
19-30	249	3.5 (0.4)	2.3 (0.4) ^E	2.5 (0.4)	2.8 (0.3)	3.2 (0.3)	3.7 (0.4)	4.2 (0.6)	4.5 (0.8) ^E	2.0	F		
31-50	364	3.8 (0.3)	1.8 (0.4) ^E	2.1 (0.4) ^E	2.6 (0.4)	3.4 (0.4)	4.7 (0.5)	6.3 (1.0)	7.6 (1.5) ^E	2.0	F		
51-70	467	4.4 (0.5)	2.0 (0.5) ^E	2.3 (0.4) ^E	2.9 (0.4)	3.8 (0.3)	4.9 (0.5)	6.3 (0.9)	7.4 (1.3) ^E	2.0	F		
>70	215	3.4 (0.6) ^E	1.7 (0.2)	1.8 (0.2)	2.2 (0.2)	2.8 (0.3)	3.8 (0.6)	5.5 (1.5) ^E	F	2.0	F		
19+	1295	3.9 (0.2)	2.0 (0.2)	2.3 (0.2)	2.7 (0.2)	3.4 (0.2)	4.5 (0.3)	5.9 (0.5)	7.1 (0.8)	2.0	F		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.6 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	644	3.2	(0.1)	1.6 (0.2)	1.9 (0.2)	2.4 (0.2)	3.1 (0.1)	3.9 (0.1)	4.7 (0.3)	5.3 (0.4)	0.7	<3
	4-8	956	3.5	(0.1)	1.8 (0.2)	2.0 (0.2)	2.6 (0.2)	3.3 (0.1)	4.2 (0.2)	5.2 (0.3)	5.9 (0.4)	1.0	<3
Male													
	9-13	589	4.3	(0.3)	2.4 (0.1)	2.7 (0.1)	3.2 (0.2)	3.9 (0.2)	4.8 (0.3)	5.9 (0.5)	6.6 (0.7)	1.5	<3
	14-18	639	5.3	(0.3)	2.3 (0.3)	2.8 (0.3)	3.6 (0.3)	4.9 (0.3)	6.7 (0.4)	8.8 (0.7)	10.3 (1.1)	2.0	F
	19-30	481	5.4	(0.6)	F	2.4 (0.7) ^E	3.2 (0.6) ^E	4.4 (0.6)	7.0 (1.0)	11.2 (3.0) ^E	F	2.0	F
	31-50	709	5.3	(0.4)	2.1 (0.6) ^E	2.5 (0.6) ^E	3.3 (0.5)	4.6 (0.5)	6.9 (0.7)	10.3 (1.7) ^E	13.3 (3.0) ^E	2.0	F
	51-70	758	4.5	(0.3)	2.1 (0.6) ^E	2.5 (0.6) ^E	3.3 (0.6) ^E	4.5 (0.6)	6.0 (0.8)	7.6 (1.1)	8.8 (1.5) ^E	2.0	F
	>70	734	4.2	(0.4)	1.8 (0.6) ^E	2.1 (0.6) ^E	2.6 (0.5) ^E	3.4 (0.4)	4.8 (0.5)	6.8 (1.2) ^E	8.5 (2.1) ^E	2.0	F
	19+	2682	5.0	(0.3)	2.0 (0.2)	2.4 (0.2)	3.1 (0.2)	4.3 (0.3)	6.5 (0.5)	9.5 (1.0)	12.0 (1.6)	2.0	F
Female													
	9-13	585	3.3	(0.1)	1.6 (0.3)	1.8 (0.2)	2.3 (0.2)	3.0 (0.1)	3.8 (0.2)	4.8 (0.4)	5.4 (0.5)	1.5	F
	14-18	645	3.3	(0.2)	1.2 (0.3) ^E	1.5 (0.2)	2.2 (0.2)	3.0 (0.2)	4.1 (0.2)	5.4 (0.4)	6.3 (0.6)	2.0	20.6 (5.7) ^E
	19-30	514	3.3	(0.5)	1.4 (0.3) ^E	1.6 (0.3) ^E	2.1 (0.3)	2.8 (0.3)	3.9 (0.5)	5.4 (1.2) ^E	6.7 (1.9) ^E	2.0	F
	31-50	758	3.6	(0.3)	1.9 (0.4) ^E	2.1 (0.3)	2.5 (0.3)	3.1 (0.2)	4.0 (0.3)	5.2 (0.8)	6.0 (1.3) ^E	2.0	F
	51-70	955	3.7	(0.3)	1.6 (0.4) ^E	1.9 (0.3) ^E	2.4 (0.3)	3.0 (0.3)	4.2 (0.3)	5.9 (0.7)	7.3 (1.2) ^E	2.0	F
	>70	1345	3.3	(0.3)	1.2 (0.1)	1.4 (0.1)	1.9 (0.1)	2.6 (0.2)	4.0 (0.4)	6.1 (1.0) ^E	8.1 (1.8) ^E	2.0	29.6 (4.4)
	19+	3572	3.5	(0.2)	1.5 (0.1)	1.7 (0.1)	2.2 (0.1)	2.9 (0.1)	4.1 (0.2)	5.8 (0.5)	7.3 (0.8)	2.0	16.9 (3.6) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.7 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both	1-3	324	3.1	(0.2)	1.4 (0.2)	1.7 (0.2)	2.3 (0.2)	3.1 (0.2)	4.0 (0.3)	4.9 (0.5)	5.6 (0.6)	0.7	<3
	4-8	425	3.3	(0.1)	2.1 (0.3) ^E	2.3 (0.3)	2.7 (0.2)	3.2 (0.2)	3.7 (0.2)	4.3 (0.4)	4.7 (0.5)	1.0	<3
Male	9-13	274	4.1	(0.2)	2.0 (0.5) ^E	2.4 (0.4) ^E	3.0 (0.3)	3.9 (0.2)	5.0 (0.4)	6.1 (0.7)	6.9 (0.9)	1.5	F
	14-18	297	5.5	(0.3)	2.3 (0.5) ^E	2.8 (0.5) ^E	3.8 (0.5)	5.2 (0.4)	7.2 (0.5)	9.5 (0.8)	11.2 (1.2)	2.0	F
	19-30	249	5.7	(0.7)	2.7 (0.3)	3.2 (0.3)	4.0 (0.4)	5.0 (0.5)	6.2 (0.9)	7.6 (1.6) ^E	8.6 (2.3) ^E	2.0	<3
	31-50	309	4.5	(0.4)	2.4 (0.2)	2.7 (0.2)	3.4 (0.3)	4.3 (0.4)	5.2 (0.5)	6.2 (0.6)	6.8 (0.8)	2.0	F
	51-70	277	5.3	(0.5)	2.5 (0.7) ^E	2.9 (0.7) ^E	3.8 (0.6)	5.2 (0.6)	7.1 (0.9)	9.2 (1.7) ^E	10.7 (2.7) ^E	2.0	F
	>70	136	4.0	(0.4)	1.3 (0.4) ^E	1.6 (0.4) ^E	2.4 (0.4) ^E	3.6 (0.4)	5.2 (0.6)	7.1 (0.9)	8.5 (1.2)	2.0	F
	19+	971	5.0	(0.3)	2.3 (0.4) ^E	2.7 (0.4)	3.4 (0.3)	4.5 (0.3)	5.9 (0.4)	7.5 (0.6)	8.7 (0.9)	2.0	F
	Female	9-13	265	3.7	(0.2)	1.6 (0.4) ^E	1.9 (0.3) ^E	2.5 (0.3)	3.2 (0.2)	4.4 (0.3)	6.0 (0.8)	7.3 (1.4) ^E	1.5
14-18	290	3.2	(0.2)	2.2 (0.4) ^E	2.4 (0.4)	2.7 (0.3)	3.1 (0.2)	3.6 (0.4)	4.0 (0.6)	4.3 (0.7) ^E	2.0	F	
19-30	197	3.3	(0.3)	1.6 (0.2)	1.9 (0.2)	2.4 (0.2)	3.1 (0.3)	3.9 (0.4)	4.9 (0.6)	5.5 (0.8)	2.0	F	
31-50	312	3.5	(0.3)	1.4 (0.4) ^E	1.7 (0.4) ^E	2.2 (0.3)	2.9 (0.3)	4.1 (0.4)	5.8 (0.9)	7.3 (1.4) ^E	2.0	F	
51-70	312	4.6	(0.8) ^E	1.9 (0.5) ^E	2.1 (0.5) ^E	2.6 (0.5) ^E	3.5 (0.5)	5.0 (0.9) ^E	7.0 (1.9) ^E	F	2.0	F	
>70	239	2.8	(0.2)	1.5 (0.4) ^E	1.8 (0.3) ^E	2.2 (0.3)	2.7 (0.2)	3.4 (0.3)	4.1 (0.6)	4.6 (0.8) ^E	2.0	F	
19+	1060	3.6	(0.3)	1.5 (0.2)	1.8 (0.2)	2.3 (0.2)	3.0 (0.2)	4.0 (0.2)	5.6 (0.5)	7.1 (0.9)	2.0	15.8 (4.7) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.8 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	129	3.5 (0.3)	1.8 (0.4) ^E	2.2 (0.4) ^E	2.7 (0.4)	3.5 (0.4)	4.5 (0.5)	5.4 (0.6)	5.9 (0.7)	0.7	<3	
	4-8	213	3.2 (0.2)	1.9 (0.3)	2.1 (0.3)	2.5 (0.2)	3.1 (0.2)	3.7 (0.2)	4.5 (0.3)	5.0 (0.5)	1.0	<3	
Male													
	9-13	122	4.1 (0.4)	2.6 (0.5) ^E	2.9 (0.5) ^E	3.4 (0.5)	4.0 (0.5)	4.8 (0.7)	5.5 (0.9)	6.0 (1.1) ^E	1.5	<3	
	14-18	150	5.2 (0.4)	2.4 (0.5) ^E	2.9 (0.5)	3.7 (0.4)	4.8 (0.5)	6.1 (0.5)	7.5 (0.7)	8.4 (0.9)	2.0	F	
	19-30	106	5.8 (0.7)	F	F	3.5 (0.9) ^E	5.1 (1.0) ^E	7.5 (1.5) ^E	11.1 (3.0) ^E	F	2.0	F	
	31-50	155	5.6 (1.1) ^E	1.9 (0.5) ^E	2.4 (0.5) ^E	3.2 (0.5) ^E	4.5 (0.6)	6.6 (1.0)	9.6 (2.2) ^E	12.3 (3.9) ^E	2.0	F	
	51-70	122	7.4 (2.2) ^E	2.4 (0.7) ^E	2.9 (0.7) ^E	3.8 (0.8) ^E	5.2 (1.2) ^E	8.0 (2.3) ^E	F	F	2.0	F	
	>70	88	5.3 (0.8)	F	2.4 (0.7) ^E	3.3 (0.7) ^E	4.6 (0.8)	6.6 (1.2) ^E	9.2 (2.2) ^E	11.3 (3.1) ^E	2.0	F	
	19+	471	6.1 (0.8)	2.0 (0.3)	2.5 (0.3)	3.4 (0.3)	4.7 (0.4)	7.1 (0.7)	11.2 (1.8)	15.3 (3.2) ^E	2.0	F	
Female													
	9-13	103	3.2 (0.3)	1.9 (0.3)	2.1 (0.3)	2.5 (0.3)	3.1 (0.4)	3.7 (0.4)	4.4 (0.5)	4.9 (0.6)	1.5	F	
	14-18	142	3.5 (0.3)	1.8 (0.4) ^E	2.2 (0.4) ^E	2.7 (0.4)	3.5 (0.4)	4.4 (0.6)	5.3 (0.8)	6.0 (1.0)	2.0	F	
	19-30	111	3.1 (0.4)	F	1.5 (0.4) ^E	2.0 (0.4) ^E	2.8 (0.5) ^E	3.7 (0.6) ^E	4.7 (0.9) ^E	5.4 (1.1) ^E	2.0	F	
	31-50	146	3.9 (0.4)	F	1.6 (0.4) ^E	2.2 (0.4) ^E	3.0 (0.5)	4.4 (0.6)	5.8 (0.9)	6.6 (1.0)	2.0	F	
	51-70	184	3.4 (0.2)	2.0 (0.2)	2.3 (0.2)	2.7 (0.3)	3.3 (0.3)	4.1 (0.4)	4.9 (0.7)	5.5 (0.9) ^E	2.0	F	
	>70	143	4.4 (0.7) ^E	1.5 (0.4) ^E	1.8 (0.4) ^E	2.4 (0.4) ^E	3.5 (0.5)	5.1 (0.9) ^E	7.4 (1.8) ^E	9.4 (3.0) ^E	2.0	F	
	19+	584	3.6 (0.2)	1.6 (0.2)	1.9 (0.2)	2.4 (0.2)	3.1 (0.2)	4.0 (0.3)	5.5 (0.6)	7.0 (1.0)	2.0	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.9 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	169	3.2 (0.2)	1.3 (0.3) ^E	1.7 (0.2)	2.3 (0.2)	3.0 (0.2)	3.9 (0.3)	5.1 (0.5)	6.0 (0.7)	0.7	<3	
	4-8	281	3.4 (0.2)	2.1 (0.1)	2.3 (0.1)	2.7 (0.2)	3.2 (0.2)	3.8 (0.2)	4.5 (0.3)	5.0 (0.4)	1.0	0.0	(0.0)
Male													
	9-13	183	6.1 (1.3) ^E	2.8 (0.8) ^E	3.2 (0.8) ^E	4.1 (0.7) ^E	5.4 (0.8)	7.2 (1.4) ^E	9.7 (2.7) ^E	F	1.5	<3	
	14-18	187	5.3 (0.3)	3.0 (0.7) ^E	3.5 (0.7) ^E	4.3 (0.5)	5.4 (0.4)	6.6 (0.6)	8.0 (1.1)	8.9 (1.4)	2.0	<3	
	19-30	223	4.9 (0.4)	2.6 (0.3)	2.9 (0.3)	3.6 (0.3)	4.6 (0.4)	5.7 (0.5)	6.8 (0.6)	7.6 (0.7)	2.0	F	
	31-50	229	5.0 (0.5)	2.5 (0.7) ^E	2.8 (0.6) ^E	3.6 (0.5)	4.6 (0.5)	5.9 (0.7)	7.3 (1.2)	8.3 (1.6) ^E	2.0	F	
	51-70	197	4.9 (0.5)	2.5 (0.7) ^E	2.9 (0.7) ^E	3.6 (0.6) ^E	4.7 (0.6)	6.1 (0.8)	7.8 (1.5) ^E	9.0 (2.1) ^E	2.0	F	
	>70	72	4.4 (0.8) ^E	2.0 (0.3) ^E	2.3 (0.4)	2.8 (0.4)	3.5 (0.5)	4.6 (0.8) ^E	6.1 (1.4) ^E	7.3 (2.2) ^E	2.0	F	
	19+	721	4.9 (0.3)	2.9 (0.5) ^E	3.2 (0.5)	3.8 (0.4)	4.7 (0.3)	5.8 (0.4)	6.9 (0.7)	7.7 (1.0)	2.0	<3	
Female													
	9-13	165	3.1 (0.2)	1.8 (0.2)	2.0 (0.2)	2.4 (0.2)	3.0 (0.2)	3.7 (0.3)	4.3 (0.3)	4.8 (0.4)	1.5	F	
	14-18	206	3.6 (0.5)	1.7 (0.6) ^E	2.0 (0.5) ^E	2.5 (0.5) ^E	3.3 (0.4)	4.2 (0.5)	5.2 (0.9)	6.0 (1.1) ^E	2.0	F	
	19-30	191	3.3 (0.3)	2.3 (0.6) ^E	2.5 (0.5) ^E	2.9 (0.4)	3.4 (0.4)	3.9 (0.5)	4.4 (0.9) ^E	4.8 (1.2) ^E	2.0	F	
	31-50	258	3.4 (0.3)	2.2 (0.6) ^E	2.4 (0.6) ^E	2.9 (0.5) ^E	3.5 (0.5)	4.1 (0.6)	4.8 (0.8) ^E	5.3 (1.1) ^E	2.0	F	
	51-70	249	3.7 (0.3)	1.7 (0.4) ^E	2.0 (0.4) ^E	2.5 (0.4)	3.3 (0.4)	4.5 (0.5)	6.0 (0.9)	7.3 (1.6) ^E	2.0	F	
	>70	128	4.2 (0.6)	1.9 (0.5) ^E	2.3 (0.5) ^E	3.0 (0.5) ^E	4.0 (0.6)	5.2 (0.8)	6.5 (1.0)	7.5 (1.3) ^E	2.0	F	
	19+	826	3.5 (0.2)	2.1 (0.4) ^E	2.3 (0.4)	2.8 (0.3)	3.5 (0.3)	4.4 (0.3)	5.5 (0.5)	6.2 (0.7)	2.0	F	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.10 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	192	3.3 (0.2)	1.8 (0.3)	2.1 (0.3)	2.6 (0.2)	3.2 (0.2)	4.0 (0.3)	4.8 (0.5)	5.3 (0.6)	0.7	<3	
	4-8	321	3.6 (0.3)	1.9 (0.2)	2.2 (0.2)	2.7 (0.2)	3.2 (0.2)	4.2 (0.4)	5.9 (1.0) ^E	7.4 (1.8) ^E	1.0	<3	
Male													
	9-13	226	4.4 (0.3)	2.4 (0.4)	2.7 (0.3)	3.3 (0.3)	4.1 (0.4)	5.3 (0.5)	6.7 (0.8)	7.9 (1.2)	1.5	<3	
	14-18	262	5.8 (0.6)	2.3 (0.4) ^E	2.9 (0.4)	3.9 (0.4)	5.4 (0.5)	7.7 (0.8)	10.8 (1.5)	13.4 (2.4) ^E	2.0	F	
	19-30	197	5.5 (0.5)	2.3 (0.5) ^E	2.7 (0.5) ^E	3.6 (0.5)	5.1 (0.6)	7.1 (0.9)	9.2 (1.5)	10.8 (2.0) ^E	2.0	F	
	31-50	282	5.7 (0.5)	2.4 (0.8) ^E	2.8 (0.8) ^E	3.8 (0.7) ^E	5.3 (0.6)	7.2 (0.7)	9.4 (1.2)	11.1 (1.7)	2.0	F	
	51-70	234	4.7 (0.3)	2.6 (0.2)	2.9 (0.3)	3.6 (0.3)	4.4 (0.3)	5.4 (0.4)	6.4 (0.5)	7.1 (0.6)	2.0	<3	
	>70	119	3.9 (0.5)	1.7 (0.2)	2.0 (0.3)	2.6 (0.3)	3.4 (0.4)	4.6 (0.6)	5.9 (0.9)	6.8 (1.1)	2.0	F	
	19+	832	5.2 (0.3)	2.6 (0.4)	3.1 (0.4)	3.9 (0.4)	5.1 (0.3)	6.5 (0.4)	8.0 (0.6)	8.9 (0.8)	2.0	F	
Female													
	9-13	226	4.3 (0.6)	2.1 (0.2)	2.4 (0.3)	3.0 (0.4)	4.0 (0.6)	5.3 (0.9) ^E	6.8 (1.5) ^E	8.0 (2.2) ^E	1.5	<3	
	14-18	242	2.9 (0.2)	1.3 (0.3) ^E	1.6 (0.3)	2.0 (0.2)	2.6 (0.2)	3.4 (0.3)	4.2 (0.4)	4.7 (0.5)	2.0	F	
	19-30	208	3.9 (0.7) ^E	F	2.1 (0.6) ^E	2.8 (0.6) ^E	3.7 (0.7) ^E	4.9 (1.0) ^E	6.3 (1.7) ^E	7.3 (2.4) ^E	2.0	F	
	31-50	263	5.2 (1.6) ^E	F	2.0 (0.6) ^E	2.5 (0.7) ^E	3.4 (1.0) ^E	5.0 (1.5) ^E	F	F	2.0	F	
	51-70	322	3.5 (0.3)	1.3 (0.3) ^E	1.5 (0.3) ^E	2.1 (0.3)	3.1 (0.3)	4.4 (0.5)	5.9 (0.9)	7.0 (1.2) ^E	2.0	F	
	>70	198	3.7 (0.4)	1.6 (0.3) ^E	1.9 (0.3) ^E	2.5 (0.4)	3.2 (0.4)	4.3 (0.7)	6.4 (1.4) ^E	8.5 (2.6) ^E	2.0	F	
	19+	991	4.3 (0.7)	1.6 (0.2)	1.9 (0.2)	2.5 (0.2)	3.4 (0.4)	5.1 (0.7)	7.5 (1.6) ^E	9.7 (2.5) ^E	2.0	13.0 (4.1) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.11 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		(SE)	
Both														
	1-3	348	4.4	(1.1) ^E	2.2 (0.5) ^E	2.5 (0.6) ^E	3.0 (0.6) ^E	3.9 (0.9) ^E	5.4 (1.4) ^E	F	F	0.7	<3	
	4-8	554	4.4	(0.8) ^E	2.6 (0.3)	2.8 (0.3)	3.2 (0.2)	3.6 (0.2)	4.3 (0.3)	5.0 (0.6)	5.6 (0.9) ^E	1.0	0.0 (0.0)	
Male														
	9-13	409	4.8	(0.3)	2.2 (0.3)	2.5 (0.3)	3.2 (0.3)	4.2 (0.3)	5.5 (0.4)	7.1 (0.6)	8.4 (0.9)	1.5	<3	
	14-18	414	5.2	(0.3)	2.7 (0.4)	3.1 (0.4)	3.9 (0.4)	4.9 (0.4)	6.2 (0.5)	7.7 (0.8)	8.8 (0.9)	2.0	<3	
	19-30	311	5.3	(0.4)	2.6 (0.5) ^E	3.0 (0.5)	3.8 (0.4)	4.9 (0.4)	6.2 (0.6)	7.7 (1.0)	8.7 (1.4)	2.0	F	
	31-50	489	5.6	(0.6)	3.3 (0.6) ^E	3.6 (0.6) ^E	4.1 (0.6)	4.9 (0.6)	6.0 (0.7)	7.3 (1.2) ^E	8.2 (1.7) ^E	2.0	<3	
	51-70	575	5.2	(0.4)	2.9 (0.5) ^E	3.2 (0.5)	3.9 (0.4)	4.7 (0.4)	5.8 (0.5)	7.1 (0.8)	8.0 (1.1)	2.0	<3	
	>70	239	4.8	(0.5)	1.9 (0.4) ^E	2.2 (0.4) ^E	2.9 (0.5)	4.0 (0.7)	5.7 (1.0) ^E	8.3 (1.8) ^E	10.8 (2.7) ^E	2.0	F	
	19+	1614	5.4	(0.3)	2.6 (0.2)	2.9 (0.2)	3.6 (0.2)	4.6 (0.3)	6.2 (0.4)	8.2 (0.6)	9.8 (0.9)	2.0	<3	
Female														
	9-13	355	3.4	(0.2)	1.6 (0.2)	1.9 (0.2)	2.3 (0.2)	3.1 (0.2)	4.1 (0.3)	5.3 (0.5)	6.2 (0.7)	1.5	F	
	14-18	410	3.5	(0.4)	1.4 (0.2)	1.7 (0.2)	2.2 (0.2)	2.9 (0.3)	4.1 (0.4)	5.9 (1.0) ^E	7.7 (1.8) ^E	2.0	F	
	19-30	384	3.7	(0.4)	2.1 (0.3)	2.3 (0.3)	2.8 (0.3)	3.5 (0.4)	4.6 (0.6)	6.1 (1.1) ^E	7.3 (1.6) ^E	2.0	F	
	31-50	585	4.2	(0.6)	1.5 (0.2)	1.8 (0.2)	2.4 (0.2)	3.4 (0.3)	5.1 (0.7)	7.8 (1.6) ^E	10.4 (2.8) ^E	2.0	14.0 (4.1) ^E	
	51-70	711	5.1	(0.8)	1.6 (0.3) ^E	1.9 (0.4) ^E	2.6 (0.5) ^E	4.0 (0.7) ^E	6.6 (1.4) ^E	10.9 (2.9) ^E	15.3 (4.8) ^E	2.0	F	
	>70	342	4.3	(0.7) ^E	1.7 (0.3) ^E	2.0 (0.3) ^E	2.5 (0.4)	3.4 (0.5)	5.0 (1.0) ^E	7.6 (2.3) ^E	F	2.0	F	
	19+	2022	4.4	(0.4)	1.7 (0.1)	2.0 (0.1)	2.6 (0.2)	3.6 (0.3)	5.5 (0.5)	8.7 (1.2)	11.9 (2.0) ^E	2.0	10.1 (2.9) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.12 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both													
	1-3	622	3.2	(0.1)	1.5 (0.1)	1.8 (0.1)	2.4 (0.1)	3.2 (0.2)	4.0 (0.2)	4.9 (0.3)	5.6 (0.4)	0.7	<3
	4-8	919	3.3	(0.1)	2.1 (0.2)	2.3 (0.2)	2.7 (0.2)	3.2 (0.1)	3.8 (0.2)	4.5 (0.3)	5.0 (0.4)	1.0	<3
Male													
	9-13	579	5.3	(0.8)	2.5 (0.4) ^E	2.9 (0.4)	3.7 (0.4)	4.5 (0.5)	6.0 (0.9)	8.2 (1.9) ^E	10.4 (3.1) ^E	1.5	<3
	14-18	634	5.3	(0.2)	2.6 (0.3)	3.1 (0.3)	4.0 (0.3)	5.2 (0.3)	6.6 (0.4)	8.3 (0.6)	9.4 (0.7)	2.0	F
	19-30	578	5.2	(0.3)	2.6 (0.5) ^E	3.0 (0.5)	3.8 (0.4)	4.8 (0.3)	6.1 (0.5)	7.5 (0.8)	8.4 (1.2)	2.0	F
	31-50	693	5.0	(0.4)	2.4 (0.5) ^E	2.8 (0.4)	3.5 (0.4)	4.5 (0.4)	5.9 (0.5)	7.5 (0.9)	8.7 (1.3)	2.0	F
	51-70	596	5.4	(0.6)	2.3 (0.5) ^E	2.8 (0.5) ^E	3.6 (0.5)	4.8 (0.5)	6.8 (0.7)	9.6 (1.4)	12.0 (2.3) ^E	2.0	F
	>70	296	4.5	(0.4)	2.1 (0.2)	2.4 (0.2)	3.1 (0.3)	4.0 (0.4)	5.3 (0.6)	7.0 (1.0)	8.4 (1.4)	2.0	F
	19+	2163	5.1	(0.2)	2.3 (0.2)	2.7 (0.2)	3.4 (0.2)	4.6 (0.2)	6.1 (0.3)	8.0 (0.5)	9.7 (0.8)	2.0	F
Female													
	9-13	533	3.2	(0.1)	1.8 (0.3)	2.0 (0.3)	2.5 (0.2)	3.1 (0.2)	3.9 (0.2)	4.7 (0.5)	5.3 (0.7)	1.5	F
	14-18	638	3.5	(0.3)	1.7 (0.4) ^E	2.0 (0.3) ^E	2.5 (0.3)	3.3 (0.3)	4.3 (0.4)	5.3 (0.6)	6.0 (0.7)	2.0	F
	19-30	499	3.2	(0.2)	1.6 (0.3) ^E	1.9 (0.3)	2.4 (0.3)	3.1 (0.3)	3.9 (0.3)	4.9 (0.5)	5.5 (0.7)	2.0	F
	31-50	716	3.5	(0.2)	1.7 (0.3) ^E	2.0 (0.3)	2.5 (0.3)	3.2 (0.4)	4.3 (0.4)	5.5 (0.5)	6.4 (0.7)	2.0	F
	51-70	745	3.9	(0.3)	2.0 (0.3) ^E	2.2 (0.3)	2.6 (0.3)	3.3 (0.3)	4.4 (0.3)	6.1 (0.8)	7.5 (1.3) ^E	2.0	F
	>70	510	3.9	(0.3)	1.6 (0.2)	1.9 (0.2)	2.6 (0.3)	3.5 (0.3)	4.7 (0.5)	6.3 (0.7)	7.6 (1.1)	2.0	F
	19+	2470	3.6	(0.1)	1.7 (0.1)	2.0 (0.1)	2.5 (0.1)	3.2 (0.2)	4.2 (0.2)	5.5 (0.3)	6.6 (0.5)	2.0	9.9 (3.0) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 25.13 Vitamin B₁₂ (µg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both													
	1-3	2117	3.4 (0.1)	1.7 (0.1)	2.0 (0.1)	2.6 (0.1)	3.2 (0.1)	4.1 (0.1)	5.2 (0.3)	6.0 (0.5)	0.7	<3	
	4-8	3235	3.6 (0.1)	2.1 (0.1)	2.3 (0.1)	2.8 (0.1)	3.4 (0.1)	4.3 (0.1)	5.3 (0.3)	6.0 (0.4)	1.0	<3	
Male													
	9-13	2080	4.6 (0.2)	2.6 (0.2)	2.9 (0.2)	3.5 (0.1)	4.2 (0.1)	5.1 (0.2)	6.3 (0.4)	7.4 (0.7)	1.5	<3	
	14-18	2288	5.5 (0.2)	2.5 (0.1)	3.0 (0.1)	3.9 (0.2)	5.2 (0.2)	7.0 (0.3)	9.1 (0.5)	10.6 (0.7)	2.0	1.7 (0.5) ^E	
	19-30	1804	5.4 (0.3)	2.4 (0.2)	2.9 (0.2)	3.6 (0.2)	4.6 (0.2)	6.5 (0.4)	8.8 (0.9)	10.7 (1.4)	2.0	F	
	31-50	2596	5.3 (0.2)	2.4 (0.3)	2.8 (0.3)	3.6 (0.2)	4.7 (0.3)	6.5 (0.4)	8.9 (0.7)	10.8 (1.1)	2.0	F	
	51-70	2550	4.9 (0.2)	2.3 (0.2)	2.6 (0.2)	3.3 (0.2)	4.3 (0.2)	5.9 (0.3)	8.0 (0.6)	9.7 (0.8)	2.0	F	
	>70	1520	4.4 (0.3)	1.8 (0.2)	2.1 (0.2)	2.7 (0.2)	3.7 (0.3)	5.3 (0.5)	7.9 (0.9)	10.2 (1.5)	2.0	F	
	19+	8470	5.1 (0.1)	2.3 (0.1)	2.6 (0.1)	3.4 (0.1)	4.4 (0.1)	6.2 (0.2)	8.6 (0.4)	10.6 (0.6)	2.0	2.7 (0.8) ^E	
Female													
	9-13	1980	3.5 (0.1)	1.8 (0.1)	2.1 (0.1)	2.5 (0.1)	3.2 (0.1)	4.1 (0.1)	5.1 (0.2)	5.8 (0.3)	1.5	F	
	14-18	2256	3.3 (0.1)	1.5 (0.1)	1.8 (0.1)	2.3 (0.1)	3.1 (0.1)	4.1 (0.2)	5.2 (0.2)	6.0 (0.3)	2.0	15.8 (2.7) ^E	
	19-30	1854	3.4 (0.2)	1.8 (0.2)	2.0 (0.2)	2.4 (0.2)	3.1 (0.2)	4.1 (0.4)	5.4 (0.7)	6.4 (1.1)	2.0	F	
	31-50	2686	3.9 (0.3)	1.6 (0.1)	1.8 (0.1)	2.4 (0.1)	3.2 (0.2)	4.6 (0.3)	6.8 (0.7)	8.6 (1.2)	2.0	13.7 (3.2) ^E	
	51-70	3200	4.0 (0.2)	1.8 (0.2)	2.1 (0.2)	2.6 (0.1)	3.3 (0.2)	4.6 (0.2)	6.4 (0.5)	7.9 (0.8)	2.0	F	
	>70	2610	3.5 (0.2)	1.6 (0.1)	1.8 (0.1)	2.3 (0.1)	3.0 (0.2)	4.2 (0.3)	6.0 (0.5)	7.7 (0.8)	2.0	15.3 (3.6) ^E	
	19+	10350	3.8 (0.1)	1.7 (0.1)	2.0 (0.1)	2.5 (0.1)	3.2 (0.1)	4.4 (0.2)	6.2 (0.4)	7.9 (0.6)	2.0	11.1 (1.9) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

26. Vitamin C (mg/d): Usual intakes from food (by smoking status)

Table 26.1¹ Vitamin C (mg/d): Usual intakes from food, by sex, region and smoking status,² household population aged 19 and older, 2004³

Age-Sex Group	Region	Smoking Status	Percentiles (and SE) of usual intake										EAR ⁴	% <EAR	(SE)	UL ⁵	% >UL	(SE)
			n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Males 19+	Atlantic region	Non-Smoker	1140	119	(6)	40 (4)	51 (4)	75 (5)	112 (7)	157 (10)	211 (14)	251 (18)	75	24.7	(3.8)	2000	0.0	(0.0)
		Smoker	474	92	(7)	31 (6) ^E	39 (6)	57 (7)	84 (9)	121 (12)	163 (17)	190 (20)	110	68.7	(7.1)	2000	0.0	(0.0)
	Quebec	Non-Smoker	800	156	(7)	55 (7)	72 (7)	107 (7)	154 (8)	211 (11)	278 (15)	328 (20)	75	11.0	(2.4) ^E	2000	0.0	(0.0)
		Smoker	378	135	(11)	70 (20) ^E	82 (20) ^E	105 (18) ^E	138 (17)	180 (18)	228 (27)	261 (36)	110	^F		2000	0.0	(0.0)
	Ontario	Non-Smoker	1990	135	(5)	49 (7)	61 (7)	87 (6)	124 (5)	173 (7)	226 (13)	264 (18)	75	17.6	(3.8) ^E	2000	0.0	(0.0)
		Smoker	690	106	(7)	^F	49 (15) ^E	68 (12) ^E	96 (8)	133 (11)	175 (23)	204 (33)	110	60.9	(7.7)	2000	0.0	(0.0)
	Prairie region	Non-Smoker	1484	143	(8)	37 (5)	49 (5)	76 (7)	122 (8)	191 (12)	277 (23)	344 (36)	75	24.6	(3.8)	2000	<3	
		Smoker	679	92	(7)	^F	44 (13) ^E	61 (12) ^E	84 (10)	114 (11)	148 (18)	171 (23)	110	72.2	(9.6)	2000	0.0	(0.0)
	British Columbia	Non-Smoker	611	151	(8)	48 (8) ^E	60 (9)	87 (10)	133 (10)	192 (13)	248 (19)	287 (26)	75	17.9	(5.0) ^E	2000	0.0	(0.0)
		Smoker	219	115	(10)	39 (13) ^E	49 (13) ^E	70 (14) ^E	107 (15)	153 (18)	207 (26)	248 (36)	110	52.1	(10.1) ^E	2000	0.0	(0.0)
Females 19+	Atlantic region	Non-Smoker	1530	107	(4)	32 (3)	42 (3)	64 (4)	96 (5)	139 (7)	189 (10)	225 (12)	60	22.2	(3.1)	2000	0.0	(0.0)
		Smoker	491	84	(6)	30 (7) ^E	38 (8) ^E	55 (8)	79 (9)	109 (11)	142 (14)	166 (17)	95	65.0	(8.2)	2000	0.0	(0.0)
	Quebec	Non-Smoker	926	138	(7)	48 (5)	61 (6)	87 (6)	126 (8)	176 (10)	232 (14)	273 (18)	60	9.7	(2.6) ^E	2000	0.0	(0.0)
		Smoker	368	115	(10)	44 (10) ^E	53 (10) ^E	75 (11)	109 (12)	156 (17)	208 (25)	245 (34)	95	40.0	(9.2) ^E	2000	0.0	(0.0)
	Ontario	Non-Smoker	2867	128	(3)	41 (4)	53 (4)	78 (4)	116 (4)	164 (5)	217 (9)	255 (12)	60	13.8	(2.1)	2000	0.0	(0.0)
		Smoker	705	92	(6)	21 (4) ^E	29 (5) ^E	45 (5)	74 (6)	117 (9)	171 (15)	210 (21)	95	64.1	(4.3)	2000	0.0	(0.0)
	Prairie region	Non-Smoker	1848	113	(4)	37 (4)	47 (4)	69 (4)	101 (5)	143 (7)	193 (10)	229 (12)	60	18.1	(2.7)	2000	0.0	(0.0)
		Smoker	621	102	(8)	48 (13) ^E	57 (13) ^E	74 (12) ^E	99 (12)	133 (14)	170 (20)	195 (26)	95	46.0	(12.2) ^E	2000	0.0	(0.0)
	British Columbia	Non-Smoker	799	125	(5)	47 (7)	58 (7)	84 (8)	123 (8)	156 (9)	196 (12)	222 (15)	60	11.2	(3.4) ^E	2000	0.0	(0.0)
		Smoker	192	100	(14)	^F	51 (16) ^E	69 (17) ^E	96 (18) ^E	133 (24) ^E	178 (39) ^E	212 (55) ^E	95	^F		2000	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Some domains were too small to produce reliable estimates. Only the domains with a large enough sample are included.

² Smokers are defined as those who smoke daily or occasionally.

³ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

⁴ EAR is the Estimated Average Requirement. Note that the EAR for smokers is increased by 35 mg/day. For additional detail, see footnote 9 in Appendix A.

⁵ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 26.2¹ Vitamin C (mg/d): Usual intakes from food, by DRI age–sex group and smoking status,² household population, Canada excluding territories, 2004³

Sex	Age (years)	Smoking Status	n	Mean	(SE)	Percentiles (and SE) of usual intake						EAR ⁴	% <EAR	(SE)	UL ⁵	% >UL	(SE)
						5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)						
Male	14-18	Non-Smoker	1917	165	(6)	57 (5)	72 (6)	104 (6)	153 (8)	217 (11)	288 (16)	63	6.9	(1.7) ^E	1800	0.0	(0.0)
		Smoker	368	153	(11)	46 (13) ^E	59 (13) ^E	89 (14)	135 (15)	198 (20)	270 (32)	98	29.7	(7.7) ^E	1800	<3	
	19-30	Non-Smoker	1136	171	(9)	52 (7)	66 (8)	100 (9)	153 (10)	226 (13)	307 (20)	75	13.6	(3.4) ^E	2000	0.0	(0.0)
		Smoker	667	131	(11)	67 (19) ^E	77 (18) ^E	97 (16)	124 (14)	160 (16)	198 (25)	110	^F		2000	0.0	(0.0)
	31-50	Non-Smoker	1647	138	(5)	43 (5)	55 (5)	82 (6)	126 (6)	183 (8)	244 (13)	75	20.7	(3.1)	2000	0.0	(0.0)
		Smoker	949	104	(6)	41 (9) ^E	51 (9) ^E	70 (8)	98 (8)	135 (10)	176 (15)	110	59.2	(6.7)	2000	0.0	(0.0)
	51-70	Non-Smoker	1871	140	(5)	48 (4)	61 (5)	89 (5)	129 (6)	181 (9)	240 (13)	75	16.8	(2.7)	2000	0.0	(0.0)
		Smoker	677	105	(9)	25 (6) ^E	35 (7) ^E	56 (8)	92 (9)	143 (13)	206 (22)	110	60.4	(6.0)	2000	0.0	(0.0)
	>70	Non-Smoker	1371	111	(4)	33 (4)	44 (4)	67 (4)	102 (5)	147 (6)	197 (9)	75	30.5	(3.0)	2000	0.0	(0.0)
		Smoker	147	115	(18)	^F	39 (11) ^E	61 (13) ^E	95 (18) ^E	147 (24)	205 (32)	110	58.5	(11.9) ^E	2000	0.0	(0.0)
Female	14-18	Non-Smoker	6025	142	(3)	44 (2)	56 (3)	86 (3)	129 (4)	188 (5)	256 (8)	75	19.3	(1.5)	2000	0.0	(0.0)
		Smoker	2440	111	(4)	41 (5)	51 (5)	71 (5)	103 (5)	145 (7)	194 (10)	110	55.3	(4.0)	2000	0.0	(0.0)
	19-30	Non-Smoker	1876	149	(5)	57 (5)	71 (5)	101 (5)	142 (6)	192 (7)	246 (11)	56	4.7	(1.4) ^E	1800	0.0	(0.0)
		Smoker	378	136	(12)	45 (11) ^E	57 (11) ^E	84 (13)	123 (15)	174 (29) ^E	233 (37)	91	29.5	(8.8) ^E	1800	0.0	(0.0)
	31-50	Non-Smoker	1228	142	(6)	52 (6)	65 (6)	94 (7)	136 (7)	188 (9)	242 (13)	60	7.9	(2.3) ^E	2000	0.0	(0.0)
		Smoker	626	109	(9)	43 (12) ^E	51 (11) ^E	68 (10)	93 (10)	126 (14)	164 (23)	95	51.6	(10.6) ^E	2000	0.0	(0.0)
	51-70	Non-Smoker	1816	124	(4)	36 (3)	47 (3)	71 (4)	108 (4)	157 (6)	214 (9)	60	17.7	(2.2)	2000	0.0	(0.0)
		Smoker	869	98	(7)	35 (6)	43 (6)	62 (7)	92 (8)	133 (10)	181 (14)	95	52.1	(6.2)	2000	0.0	(0.0)
	>70	Non-Smoker	2551	126	(4)	46 (3)	58 (4)	82 (4)	117 (4)	160 (6)	206 (8)	60	11.2	(1.9) ^E	2000	0.0	(0.0)
		Smoker	648	102	(6)	28 (6) ^E	38 (6) ^E	59 (7)	90 (8)	133 (11)	186 (17)	95	53.4	(5.6)	2000	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Some domains were too small to produce reliable estimates. Only the domains with a large enough sample are included.

² Smokers are defined as those who smoke daily or occasionally.

³ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

⁴ EAR is the Estimated Average Requirement. Note that the EAR for smokers is increased by 35 mg/day. For additional detail, see footnote 9 in Appendix A.

⁵ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

27. Vitamin D (µg/d): Usual intakes from food

Table 27.1 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							% AI ² >AI (SE)			% UL ³ >UL (SE)		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	79	6.1	(0.4)	2.9 (0.7) ^E	3.5 (0.7) ^E	4.7 (0.6)	6.3 (0.6)	8.0 (0.7)	9.5 (1.0)	10.6 (1.2)	5	70.7 (9.7)	50	0.0 (0.0)		
	4-8	127	6.1	(0.4)	3.5 (0.3)	4.0 (0.4)	4.9 (0.4)	6.0 (0.5)	7.3 (0.6)	8.7 (0.8)	9.7 (1.1)	5	72.4 (9.3)	50	0.0 (0.0)		
Male																	
	9-13	111	7.9	(0.9)	F	4.2 (1.1) ^E	5.7 (1.0) ^E	7.7 (1.0)	9.7 (1.3)	12.0 (2.0)	13.7 (2.5) ^E	5	82.8 (10.9)	50	<3		
	14-18	107	6.2	(0.7)	F	F	3.7 (0.9) ^E	5.8 (0.9)	8.7 (1.4)	11.8 (2.4) ^E	13.7 (3.2) ^E	5	59.2 (12.4) ^E	50	<3		
	19-30	77	4.5	(0.8) ^E	1.4 (0.4) ^E	1.8 (0.4) ^E	2.5 (0.5) ^E	3.7 (0.7) ^E	5.4 (1.1) ^E	7.6 (1.6) ^E	9.1 (2.0) ^E	5	F	50	<3		
	31-50	145	7.7	(2.0) ^E	4.1 (1.2) ^E	4.5 (1.2) ^E	5.2 (1.2) ^E	6.1 (1.2) ^E	7.2 (1.5) ^E	8.4 (2.2) ^E	9.2 (2.9) ^E	5	F	50	0.0 (0.0)		
	51-70	182	6.3	(1.1) ^E	2.5 (0.3)	2.9 (0.4)	3.9 (0.5)	5.2 (0.7)	7.0 (1.2) ^E	9.2 (2.1) ^E	10.8 (3.0) ^E	10	F	50	<3		
	>70	63	9.1	(1.5) ^E	3.6 (0.9) ^E	4.3 (0.9) ^E	5.6 (1.0) ^E	7.7 (1.3) ^E	11.2 (1.9) ^E	16.8 (3.5) ^E	22.1 (5.4) ^E	15	F	50	<3		
	19+	467	6.7	(0.9)	2.7 (0.5) ^E	3.2 (0.5) ^E	4.1 (0.6)	5.6 (0.6)	7.6 (0.8)	10.1 (1.3)	12.0 (1.7)			50	0.0 (0.0)		
Female																	
	9-13	96	5.6	(0.6)	F	2.4 (0.7) ^E	3.7 (0.7) ^E	5.3 (0.8)	7.2 (1.1)	9.2 (1.6) ^E	10.6 (2.0) ^E	5	54.7 (14.0) ^E	50	0.0 (0.0)		
	14-18	105	4.9	(1.0) ^E	F	F	F	4.1 (1.0) ^E	6.1 (1.4) ^E	9.0 (2.4) ^E	11.4 (3.4) ^E	5	F	50	<3		
	19-30	91	4.4	(1.0) ^E	F	F	2.5 (0.7) ^E	3.6 (0.9) ^E	5.3 (1.3) ^E	7.5 (2.2) ^E	9.3 (3.0) ^E	5	F	50	<3		
	31-50	167	3.8	(0.3)	1.6 (0.2)	2.0 (0.2)	2.8 (0.3)	3.6 (0.4)	4.7 (0.6)	6.1 (0.8)	7.1 (1.0)	5	F	50	0.0 (0.0)		
	51-70	198	4.9	(0.5)	F	2.7 (0.8) ^E	3.4 (0.7) ^E	4.6 (0.6)	6.0 (0.8)	7.8 (1.4) ^E	9.2 (2.0) ^E	10	F	50	<3		
	>70	74	5.0	(0.5)	2.6 (0.7) ^E	3.0 (0.6) ^E	3.7 (0.6) ^E	4.5 (0.6)	5.5 (0.8)	6.5 (1.0)	7.1 (1.2) ^E	15	<3	50	0.0 (0.0)		
	19+	530	4.4	(0.3)	2.2 (0.3)	2.6 (0.3)	3.3 (0.3)	4.3 (0.3)	5.7 (0.4)	7.2 (0.7)	8.3 (1.0)			50	0.0 (0.0)		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.2 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							% AI ² >AI (SE)			% UL ³ >UL (SE)		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	58	7.9	(1.1)	3.2 (1.0) ^E	4.1 (1.0) ^E	5.7 (1.0) ^E	7.9 (1.1)	10.4 (1.5)	12.8 (2.0)	14.5 (2.4)	5	82.3	(8.3)	50	0.0	(0.0)
	4-8	110	7.7	(1.0)	5.4 (1.2) ^E	5.8 (1.1) ^E	6.6 (1.1)	7.5 (1.1)	8.6 (1.3)	9.5 (1.5)	10.1 (1.7) ^E	5	97.4	(11.6)	50	0.0	(0.0)
Male																	
	9-13	95	6.1	(0.4)	4.1 (0.8) ^E	4.5 (0.7)	5.2 (0.6)	6.0 (0.6)	6.9 (0.7)	7.7 (1.0)	8.2 (1.2)	5	80.2	(14.0) ^E	50	0.0	(0.0)
	14-18	87	8.4	(1.8) ^E	F	3.8 (1.1) ^E	5.2 (1.3) ^E	7.3 (1.6) ^E	10.4 (2.7) ^E	14.4 (4.2) ^E	17.8 (5.6) ^E	5	77.5	(12.8)	50	<3	
	19-30	70	12.1	(2.2) ^E	F	6.2 (1.9) ^E	8.4 (2.0) ^E	11.8 (2.6) ^E	16.3 (3.8) ^E	21.3 (5.3) ^E	24.5 (6.3) ^E	5	95.1	(5.7)	50	<3	
	31-50	109	6.2	(0.8)	F	F	3.7 (0.8) ^E	5.9 (0.9)	8.7 (1.2)	11.9 (1.8)	14.1 (2.3)	5	60.1	(11.3) ^E	50	<3	
	51-70	128	6.8	(0.7)	2.9 (0.7) ^E	3.4 (0.7) ^E	4.3 (0.6)	5.7 (0.7)	8.0 (1.0)	11.3 (2.0) ^E	14.3 (3.4) ^E	10	F		50	<3	
	>70	65	6.1	(0.7)	2.1 (0.6) ^E	2.6 (0.6) ^E	3.7 (0.7) ^E	5.2 (0.8)	7.3 (1.1)	10.4 (1.9) ^E	13.0 (2.7) ^E	15	F		50	<3	
	19+	372	7.6	(0.6)	2.5 (0.4) ^E	3.2 (0.5)	4.7 (0.5)	6.8 (0.6)	9.9 (0.9)	14.0 (1.5)	17.2 (2.1)				50	<3	
Female																	
	9-13	75	7.3	(1.3) ^E	F	F	6.0 (1.8) ^E	7.5 (1.9) ^E	9.2 (2.2) ^E	10.9 (2.7) ^E	12.0 (3.0) ^E	5	88.8	(23.1) ^E	50	0.0	(0.0)
	14-18	81	6.6	(1.0)	F	F	3.6 (0.9) ^E	5.2 (1.0) ^E	7.3 (1.4) ^E	9.6 (2.0) ^E	11.2 (2.6) ^E	5	52.7	(15.9) ^E	50	<3	
	19-30	101	5.5	(0.5)	F	2.3 (0.8) ^E	3.7 (0.8) ^E	5.6 (0.8)	8.0 (1.0)	10.5 (1.4)	12.0 (1.8)	5	57.8	(11.3) ^E	50	0.0	(0.0)
	31-50	116	5.2	(0.8)	F	2.1 (0.6) ^E	3.2 (0.6) ^E	4.8 (0.8)	7.0 (1.1)	9.5 (1.7) ^E	11.2 (2.1) ^E	5	47.4	(12.0) ^E	50	<3	
	51-70	146	5.7	(0.7)	2.1 (0.4) ^E	2.5 (0.4) ^E	3.5 (0.5)	5.0 (0.6)	7.0 (0.8)	9.9 (1.4)	12.5 (2.1) ^E	10	F		50	<3	
	>70	94	4.7	(0.5)	1.9 (0.4) ^E	2.2 (0.4) ^E	2.9 (0.5)	4.0 (0.6)	5.5 (0.7)	7.5 (1.1)	8.9 (1.4)	15	<3		50	0.0	(0.0)
	19+	457	5.3	(0.4)	1.7 (0.2)	2.2 (0.2)	3.3 (0.3)	4.9 (0.4)	7.0 (0.6)	9.6 (0.9)	11.5 (1.2)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.3 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							% AI ² >AI (SE)			% UL ³ >UL (SE)		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	112	6.2	(0.5)	F	3.8 (1.1) ^E	4.9 (0.8)	6.2 (0.6)	7.5 (0.7)	8.7 (1.0)	9.5 (1.4)	5	73.0 (14.4) ^E	50	0.0	(0.0)	
	4-8	177	7.3	(0.6)	2.8 (0.6) ^E	3.6 (0.5)	5.1 (0.5)	7.0 (0.6)	9.4 (0.8)	12.1 (1.2)	14.1 (1.5)	5	76.0 (6.1)	50	0.0	(0.0)	
Male																	
	9-13	111	7.1	(0.5)	2.9 (0.8) ^E	3.5 (0.7) ^E	4.7 (0.6)	6.3 (0.5)	8.4 (0.7)	10.7 (1.1)	12.2 (1.6)	5	70.0 (9.5)	50	0.0	(0.0)	
	14-18	113	10.5	(3.4) ^E	F	F	F	8.3 (2.6) ^E	11.2 (3.4) ^E	F	F	5	87.9 (9.5)	50	<3		
	19-30	91	7.8	(0.9)	F	3.3 (1.0) ^E	4.9 (1.0) ^E	7.1 (1.0)	9.8 (1.4)	12.8 (1.9)	14.8 (2.4)	5	73.6 (11.5)	50	<3		
	31-50	101	5.8	(0.6)	3.1 (0.8) ^E	3.5 (0.8) ^E	4.3 (0.7)	5.3 (0.7)	6.6 (0.8)	7.8 (1.2)	8.7 (1.6) ^E	5	58.1 (17.1) ^E	50	0.0	(0.0)	
	51-70	134	6.7	(0.7)	3.0 (0.9) ^E	3.5 (0.8) ^E	4.6 (0.8)	6.3 (0.7)	8.3 (0.9)	10.6 (1.4)	12.4 (2.1) ^E	10	F	50	<3		
	>70	56	8.0	(1.2)	F	F	5.2 (1.4) ^E	7.4 (1.5) ^E	10.4 (2.2) ^E	14.0 (3.6) ^E	16.6 (5.1) ^E	15	F	50	<3		
	19+	382	6.7	(0.4)	2.7 (0.4)	3.3 (0.4)	4.4 (0.4)	6.1 (0.4)	8.4 (0.6)	11.0 (0.9)	12.9 (1.2)			50	0.0	(0.0)	
Female																	
	9-13	105	5.5	(0.7)	F	2.5 (0.7) ^E	3.5 (0.6) ^E	4.8 (0.7)	6.4 (0.8)	8.2 (1.2)	9.3 (1.5)	5	46.9 (13.8) ^E	50	0.0	(0.0)	
	14-18	120	5.0	(0.5)	F	F	3.2 (0.6) ^E	4.5 (0.6)	6.1 (0.9)	7.7 (1.4) ^E	8.8 (1.8) ^E	5	41.5 (13.6) ^E	50	0.0	(0.0)	
	19-30	91	5.4	(0.6)	2.8 (0.9) ^E	3.3 (0.9) ^E	4.2 (0.9) ^E	5.4 (0.9)	6.8 (1.0)	8.2 (1.3)	9.3 (1.6) ^E	5	58.9 (19.2) ^E	50	0.0	(0.0)	
	31-50	159	5.8	(0.5)	2.2 (0.4) ^E	2.8 (0.4)	4.1 (0.5)	5.7 (0.6)	7.6 (0.8)	9.9 (1.1)	11.6 (1.3)	5	60.7 (9.7)	50	0.0	(0.0)	
	51-70	174	5.7	(0.8)	2.0 (0.6) ^E	2.4 (0.6) ^E	3.3 (0.6) ^E	4.7 (0.7)	6.6 (1.0)	8.9 (1.5) ^E	10.5 (2.0) ^E	10	F	50	<3		
	>70	80	6.5	(1.0)	2.4 (0.7) ^E	2.8 (0.8) ^E	3.7 (0.9) ^E	5.3 (1.2) ^E	7.6 (1.7) ^E	10.5 (2.5) ^E	12.7 (3.3) ^E	15	F	50	<3		
	19+	504	5.8	(0.3)	2.1 (0.3)	2.6 (0.3)	3.8 (0.4)	5.3 (0.4)	7.4 (0.6)	10.2 (0.9)	12.4 (1.3)			50	0.0	(0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.4 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake										% AI ² >AI (SE)			% UL ³ >UL (SE)		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			(SE)			(SE)
Both																	
	1-3	99	8.1	(0.8)	4.5 (1.2) ^E	5.2 (1.1) ^E	6.3 (0.9)	7.7 (0.9)	9.6 (1.2)	11.6 (1.7)	13.0 (2.2) ^E	5	91.6	(7.8)	50	0.0	(0.0)
	4-8	140	5.9	(0.4)	3.7 (0.7) ^E	4.1 (0.6)	4.8 (0.5)	5.6 (0.4)	6.6 (0.5)	7.6 (0.8)	8.3 (1.0)	5	68.9	(12.9) ^E	50	0.0	(0.0)
Male																	
	9-13	92	7.0	(0.6)	3.5 (0.8) ^E	4.1 (0.8) ^E	5.2 (0.7)	6.5 (0.8)	8.1 (1.0)	10.0 (1.6)	11.3 (2.0) ^E	5	77.9	(12.2)	50	<3	
	14-18	107	9.8	(1.0)	5.6 (1.2) ^E	6.3 (1.2) ^E	7.6 (1.2)	9.6 (1.2)	12.1 (1.5)	15.4 (2.2)	17.9 (2.9)	5	97.3	(4.2)	50	<3	
	19-30	73	7.4	(0.7)	^F	4.2 (1.2) ^E	5.6 (1.1) ^E	7.4 (1.1)	9.7 (1.4)	12.4 (2.4) ^E	14.5 (3.3) ^E	5	81.6	(11.7)	50	<3	
	31-50	134	6.5	(0.7)	3.8 (1.0) ^E	4.3 (0.9) ^E	5.1 (0.8)	6.1 (0.8)	7.4 (0.9)	8.7 (1.4)	9.6 (1.8) ^E	5	77.0	(16.3) ^E	50	0.0	(0.0)
	51-70	131	7.2	(0.8)	3.6 (0.8) ^E	4.1 (0.8) ^E	5.0 (0.8)	6.4 (0.8)	8.1 (1.1)	10.1 (1.8) ^E	11.5 (2.4) ^E	10	^F		50	<3	
	>70	55	6.3	(0.6)	^F	3.5 (1.1) ^E	4.8 (1.0) ^E	6.3 (0.9)	8.0 (0.9)	9.7 (1.1)	10.7 (1.3)	15	<3		50	0.0	(0.0)
	19+	393	6.9	(0.4)	3.4 (0.5)	4.0 (0.5)	5.0 (0.5)	6.5 (0.5)	8.4 (0.7)	10.5 (1.0)	12.1 (1.3)				50	0.0	(0.0)
Female																	
	9-13	79	6.9	(0.7)	3.5 (0.8) ^E	4.0 (0.8) ^E	5.0 (0.8)	6.4 (1.0)	8.1 (1.2)	9.9 (1.5)	11.1 (1.7)	5	74.8	(13.1) ^E	50	0.0	(0.0)
	14-18	104	5.7	(1.1) ^E	^F	^F	5.2 (1.4) ^E	6.9 (1.6) ^E	9.3 (2.3) ^E	11.9 (3.3) ^E	13.5 (4.2) ^E	5	77.4	(20.9) ^E	50	<3	
	19-30	101	5.7	(1.1) ^E	^F	^F	3.0 (0.9) ^E	4.9 (1.1) ^E	7.9 (1.7) ^E	11.8 (2.9) ^E	15.0 (4.0) ^E	5	49.1	(14.0) ^E	50	<3	
	31-50	143	5.4	(0.8)	1.9 (0.3)	2.3 (0.3)	3.3 (0.4)	4.8 (0.7)	6.7 (1.4) ^E	9.3 (3.0) ^E	^F	5	46.4	(12.1) ^E	50	<3	
	51-70	193	4.9	(0.4)	2.4 (0.5) ^E	2.7 (0.4)	3.4 (0.4)	4.4 (0.4)	5.6 (0.6)	7.3 (1.0)	8.6 (1.5) ^E	10	^F		50	0.0	(0.0)
	>70	94	5.1	(0.6)	2.1 (0.5) ^E	2.5 (0.6) ^E	3.3 (0.7) ^E	4.4 (0.9) ^E	5.9 (1.1) ^E	7.7 (1.3) ^E	9.0 (1.5) ^E	15	<3		50	0.0	(0.0)
	19+	531	5.3	(0.4)	2.2 (0.3)	2.6 (0.3)	3.5 (0.3)	4.8 (0.4)	6.5 (0.6)	8.8 (1.1)	10.5 (1.5)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.5 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI (SE)	UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)					95th (SE)
Both															
	1-3	311	6.4	(0.3)	2.5 (0.3)	3.2 (0.3)	4.6 (0.4)	6.2 (0.4)	8.2 (0.5)	10.3 (0.6)	11.7 (0.8)	5	68.7 (5.6)	50	0.0 (0.0)
	4-8	485	6.2	(0.4)	2.6 (0.5) ^E	3.2 (0.5)	4.3 (0.4)	5.8 (0.4)	7.5 (0.6)	9.4 (1.0)	10.7 (1.3)	5	64.0 (7.1)	50	0.0 (0.0)
Male															
	9-13	277	7.2	(0.4)	3.6 (0.6)	4.2 (0.6)	5.4 (0.5)	7.0 (0.6)	9.0 (0.7)	11.1 (1.1)	12.7 (1.4)	5	80.4 (7.4)	50	0.0 (0.0)
	14-18	339	7.7	(0.5)	2.8 (0.7) ^E	3.6 (0.6) ^E	5.2 (0.6)	7.4 (0.6)	10.4 (0.8)	13.9 (1.2)	16.4 (1.7)	5	77.0 (6.2)	50	<3
	19-30	237	5.2	(0.4)	^F	2.4 (0.6) ^E	3.2 (0.5)	4.4 (0.5)	5.9 (0.6)	7.6 (1.1)	8.9 (1.5) ^E	5	39.3 (11.0) ^E	50	0.0 (0.0)
	31-50	423	6.2	(0.7)	2.3 (0.3)	2.8 (0.4)	3.9 (0.4)	5.5 (0.6)	7.8 (1.0)	10.5 (2.0) ^E	12.5 (3.3) ^E	5	57.7 (7.9)	50	<3
	51-70	387	6.9	(0.7)	2.0 (0.5) ^E	2.6 (0.5) ^E	3.8 (0.5)	5.8 (0.6)	8.8 (1.0)	13.2 (2.1)	16.9 (3.4) ^E	10	19.4 (5.6) ^E	50	<3
	>70	132	6.9	(1.4) ^E	^F	^F	3.5 (1.0) ^E	5.1 (1.2) ^E	7.8 (1.9) ^E	12.0 (3.4) ^E	^F	15	^F	50	<3
	19+	1179	6.3	(0.4)	2.2 (0.3)	2.7 (0.3)	3.7 (0.3)	5.3 (0.4)	7.8 (0.6)	11.1 (1.0)	13.8 (1.4)			50	<3
Female															
	9-13	281	6.0	(0.5)	2.6 (0.6) ^E	3.1 (0.6) ^E	4.2 (0.5)	5.5 (0.5)	7.1 (0.7)	9.0 (1.0)	10.2 (1.2)	5	59.4 (10.6) ^E	50	0.0 (0.0)
	14-18	321	5.0	(0.4)	2.3 (0.6) ^E	2.7 (0.5) ^E	3.6 (0.5)	4.7 (0.5)	6.3 (0.7)	8.1 (1.1)	9.4 (1.6)	5	44.8 (11.5) ^E	50	<3
	19-30	249	5.0	(0.5)	2.0 (0.6) ^E	2.5 (0.6) ^E	3.3 (0.5)	4.6 (0.6)	6.3 (0.8)	8.3 (1.2)	9.8 (1.7) ^E	5	42.8 (12.0) ^E	50	0.0 (0.0)
	31-50	364	6.5	(1.2) ^E	2.4 (0.6) ^E	2.8 (0.7) ^E	3.8 (0.9) ^E	5.7 (1.1) ^E	8.6 (1.8) ^E	12.6 (3.1) ^E	16.1 (4.7) ^E	5	58.5 (13.2) ^E	50	<3
	51-70	467	6.2	(0.9)	2.0 (0.6) ^E	2.4 (0.6) ^E	3.4 (0.6) ^E	5.2 (0.8)	8.1 (1.1)	12.2 (2.3) ^E	15.7 (3.5) ^E	10	^F	50	<3
	>70	215	5.6	(0.9)	2.6 (0.6) ^E	3.0 (0.6) ^E	3.9 (0.6)	5.2 (0.8)	6.9 (1.0)	8.9 (1.6) ^E	10.4 (2.2) ^E	15	^F	50	<3
	19+	1295	6.0	(0.5)	2.3 (0.3)	2.8 (0.3)	3.7 (0.4)	5.3 (0.7)	7.8 (0.9)	11.2 (1.4)	14.0 (2.0)			50	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.6 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									AI ²	% >AI (<i>SE</i>)	UL ³	% >UL (<i>SE</i>)										
		n	Mean	(<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)					95th (<i>SE</i>)									
Both																								
	1-3	644	6.5	(0.2)	1.7	(0.4) ^E	2.6	(0.3)	4.2	(0.3)	6.2	(0.3)	8.6	(0.3)	11.2	(0.5)	13.1	(0.8)	5	65.0	(3.4)	50	0.0	(0.0)
	4-8	956	5.9	(0.2)	2.7	(0.4)	3.2	(0.4)	4.2	(0.3)	5.6	(0.2)	7.4	(0.3)	9.4	(0.6)	10.7	(0.8)	5	60.1	(5.1)	50	0.0	(0.0)
Male																								
	9-13	589	6.5	(0.3)	2.9	(0.5) ^E	3.5	(0.5)	4.6	(0.4)	6.2	(0.3)	8.0	(0.4)	9.9	(0.7)	11.1	(0.9)	5	69.4	(6.2)	50	0.0	(0.0)
	14-18	639	7.2	(0.4)	2.8	(0.5) ^E	3.5	(0.5)	4.9	(0.4)	6.8	(0.4)	9.2	(0.6)	12.1	(1.0)	14.1	(1.4)	5	73.2	(5.9)	50	0.0	(0.0)
	19-30	481	5.8	(0.5)	3.3	(1.1) ^E	3.8	(1.0) ^E	4.7	(0.8) ^E	5.8	(0.7)	7.1	(0.8)	8.5	(1.3)	9.5	(2.0) ^E	5	67.7	(16.0) ^E	50	<3	
	31-50	709	5.3	(0.4)	1.8	(0.5) ^E	2.2	(0.5) ^E	3.2	(0.4)	4.6	(0.4)	6.6	(0.5)	9.1	(1.0)	11.1	(1.5)	5	43.6	(6.2)	50	<3	
	51-70	758	6.5	(0.8)	1.8	(0.4) ^E	2.3	(0.4)	3.3	(0.4)	4.8	(0.5)	7.6	(0.8)	11.7	(1.9)	15.4	(3.1) ^E	10	^F		50	<3	
	>70	734	5.5	(0.4)	2.0	(0.5) ^E	2.5	(0.5) ^E	3.5	(0.4)	4.9	(0.4)	6.8	(0.5)	9.1	(0.9)	10.9	(1.3)	15	^F		50	<3	
	19+	2682	5.7	(0.3)	2.1	(0.3)	2.5	(0.3)	3.5	(0.3)	5.0	(0.3)	7.3	(0.5)	10.4	(0.9)	12.9	(1.4)				50	<3	
Female																								
	9-13	585	5.4	(0.2)	2.1	(0.5) ^E	2.6	(0.4)	3.6	(0.3)	4.9	(0.3)	6.6	(0.4)	8.4	(0.7)	9.6	(1.0)	5	48.8	(5.3)	50	0.0	(0.0)
	14-18	645	5.1	(0.3)	1.5	(0.5) ^E	2.0	(0.5) ^E	2.9	(0.4)	4.4	(0.3)	6.5	(0.5)	9.4	(1.2)	11.7	(2.1) ^E	5	40.9	(5.5)	50	<3	
	19-30	514	4.0	(0.2)	1.7	(0.5) ^E	2.1	(0.4) ^E	2.8	(0.3)	3.8	(0.3)	5.0	(0.3)	6.3	(0.6)	7.3	(0.9)	5	25.0	(6.6) ^E	50	0.0	(0.0)
	31-50	758	4.7	(0.2)	1.7	(0.4) ^E	2.2	(0.3)	3.0	(0.3)	4.2	(0.3)	5.9	(0.4)	8.0	(0.7)	9.6	(1.0)	5	36.2	(4.8)	50	0.0	(0.0)
	51-70	955	4.4	(0.2)	1.6	(0.4) ^E	2.0	(0.3) ^E	2.7	(0.3)	3.8	(0.3)	5.5	(0.3)	7.6	(0.6)	9.3	(1.0)	10	^F		50	0.0	(0.0)
	>70	1345	4.8	(0.2)	1.7	(0.2)	2.1	(0.2)	2.9	(0.2)	4.2	(0.2)	6.1	(0.3)	8.4	(0.6)	10.2	(0.8)	15	<3		50	0.0	(0.0)
	19+	3572	4.5	(0.1)	1.6	(0.2)	2.0	(0.2)	2.8	(0.1)	4.0	(0.1)	5.6	(0.2)	7.8	(0.3)	9.4	(0.5)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.7 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI		UL ³	% >UL										
					(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)		(SE)	(SE)		(SE)	(SE)									
Both																										
	1-3	324	6.3	(0.4)	2.0	(0.4) ^E	2.7	(0.4)	4.1	(0.4)	6.1	(0.5)	8.5	(0.6)	11.0	(0.8)	12.7	(1.0)	5	63.4	(6.7)	50	0.0	(0.0)		
	4-8	425	5.9	(0.2)	2.7	(0.6) ^E	3.2	(0.5)	4.2	(0.4)	5.5	(0.3)	7.0	(0.4)	8.6	(0.7)	9.7	(1.1)	5	59.8	(7.5)	50	0.0	(0.0)		
Male																										
	9-13	274	7.0	(0.4)	3.4	(1.0) ^E	4.0	(0.9) ^E	5.2	(0.7)	6.7	(0.5)	8.4	(0.5)	10.2	(0.9)	11.3	(1.2)	5	77.6	(10.4)	50	0.0	(0.0)		
	14-18	297	8.4	(0.7)	F		2.8	(0.8) ^E	4.8	(0.7)	7.7	(0.8)	11.5	(1.2)	16.0	(2.0)	19.3	(2.7)	5	73.3	(6.8)	50	<3			
	19-30	249	7.5	(1.2)	F		3.6	(1.1) ^E	4.8	(0.9) ^E	6.5	(0.8)	8.6	(1.1)	11.0	(1.9) ^E	12.8	(2.7) ^E	5	71.9	(13.0) ^E	50	0.0	(0.0)		
	31-50	309	5.1	(0.4)	F		2.5	(0.8) ^E	3.4	(0.7) ^E	4.8	(0.5)	6.6	(0.7)	8.3	(1.1)	9.4	(1.4)	5	46.3	(11.5) ^E	50	0.0	(0.0)		
	51-70	277	5.7	(0.4)			2.7	(0.6) ^E	3.1	(0.6) ^E	4.1	(0.5)	5.4	(0.4)	7.1	(0.6)	9.0	(1.1)	10.4	(1.6)	10	F		50	0.0	(0.0)
	>70	136	5.5	(0.4)			2.1	(0.5) ^E	2.5	(0.5) ^E	3.5	(0.5)	5.0	(0.4)	6.9	(0.6)	9.2	(1.0)	10.8	(1.4)	15	<3		50	0.0	(0.0)
	19+	971	5.8	(0.4)			2.1	(0.4) ^E	2.6	(0.4)	3.7	(0.4)	5.2	(0.3)	7.2	(0.4)	9.5	(0.7)	11.2	(1.0)				50	0.0	(0.0)
Female																										
	9-13	265	6.9	(0.6)			2.4	(0.6) ^E	3.0	(0.6) ^E	4.2	(0.5)	6.0	(0.4)	8.4	(0.7)	11.3	(1.7)	13.8	(2.8) ^E	5	63.3	(7.9)	50	<3	
	14-18	290	5.2	(0.4)			F		2.8	(0.9) ^E	3.8	(0.8) ^E	5.3	(0.8)	7.1	(0.8)	9.0	(1.1)	10.3	(1.4)	5	54.9	(14.4) ^E	50	0.0	(0.0)
	19-30	197	5.4	(0.7)			F		F		3.0	(0.6) ^E	4.7	(0.7)	7.0	(1.1)	9.9	(2.0) ^E	12.3	(3.0) ^E	5	45.7	(10.5) ^E	50	<3	
	31-50	312	4.4	(0.3)			1.6	(0.4) ^E	2.0	(0.4) ^E	2.8	(0.4)	3.9	(0.3)	5.3	(0.5)	6.7	(0.7)	7.7	(0.9)	5	29.2	(7.8) ^E	50	0.0	(0.0)
	51-70	312	5.4	(0.7)			1.8	(0.5) ^E	2.3	(0.5) ^E	3.1	(0.5)	4.4	(0.5)	6.1	(0.7)	8.3	(1.3)	9.9	(1.8) ^E	10	F		50	0.0	(0.0)
	>70	239	5.0	(0.5)			1.8	(0.4) ^E	2.3	(0.4) ^E	3.2	(0.4)	4.7	(0.4)	6.6	(0.7)	9.1	(1.2)	11.0	(1.8) ^E	15	F		50	<3	
	19+	1060	5.0	(0.3)			1.5	(0.2)	1.9	(0.2)	2.9	(0.2)	4.2	(0.2)	6.0	(0.3)	8.5	(0.6)	10.5	(0.9)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.8 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI (SE)		UL ³	% >UL (SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	129	7.8	(0.8)	3.1 (0.7) ^E	3.9 (0.7) ^E	5.5 (0.7)	7.7 (0.8)	10.3 (1.1)	13.0 (1.5)	14.8 (1.8)	5	80.5 (7.2)	50	0.0 (0.0)	
	4-8	213	6.9	(0.7)	2.6 (0.4) ^E	3.2 (0.4)	4.3 (0.4)	5.7 (0.4)	7.6 (0.5)	10.2 (1.3)	12.7 (2.5) ^E	5	62.3 (6.5)	50	<3	
Male																
	9-13	122	6.6	(0.5)	2.8 (0.7) ^E	3.5 (0.7) ^E	4.7 (0.7)	6.3 (0.7)	8.1 (0.9)	10.0 (1.1)	11.2 (1.3)	5	70.2 (10.8)	50	0.0 (0.0)	
	14-18	150	7.3	(0.5)	2.0 (0.5) ^E	2.7 (0.5) ^E	4.2 (0.5)	6.2 (0.6)	8.8 (0.8)	11.8 (1.2)	13.9 (1.6)	5	64.6 (7.0)	50	0.0 (0.0)	
	19-30	106	6.1	(0.6)	F	3.2 (0.9) ^E	4.3 (0.8) ^E	5.6 (0.9)	7.3 (1.1)	9.0 (1.5) ^E	10.1 (1.9) ^E	5	61.6 (16.7) ^E	50	0.0 (0.0)	
	31-50	155	6.4	(0.6)	F	3.0 (0.9) ^E	4.2 (0.8) ^E	5.9 (0.8)	8.0 (1.0)	10.4 (1.5)	12.2 (2.0)	5	63.0 (13.5) ^E	50	<3	
	51-70	122	7.2	(0.7)	3.8 (1.2) ^E	4.5 (1.1) ^E	5.8 (1.0) ^E	7.4 (1.0)	9.4 (1.3)	11.4 (1.9)	12.8 (2.4) ^E	10	F	50	0.0 (0.0)	
	>70	88	6.9	(0.7)	2.8 (0.6) ^E	3.3 (0.7) ^E	4.5 (0.7)	6.3 (0.8)	8.6 (1.0)	11.3 (1.6)	13.3 (2.0)	15	F	50	0.0 (0.0)	
	19+	471	6.6	(0.3)	2.6 (0.4) ^E	3.2 (0.4)	4.4 (0.4)	6.3 (0.4)	8.6 (0.6)	11.1 (0.9)	12.8 (1.2)			50	0.0 (0.0)	
Female																
	9-13	103	5.6	(0.4)	3.2 (0.9) ^E	3.7 (0.8) ^E	4.5 (0.7)	5.6 (0.6)	6.8 (0.8)	8.2 (1.2)	9.1 (1.6) ^E	5	64.0 (15.9) ^E	50	0.0 (0.0)	
	14-18	142	5.0	(0.5)	2.0 (0.6) ^E	2.4 (0.6) ^E	3.3 (0.7) ^E	4.5 (0.9) ^E	6.4 (1.2) ^E	8.5 (1.6) ^E	9.8 (2.0) ^E	5	F	50	0.0 (0.0)	
	19-30	111	5.4	(0.9)	2.2 (0.4) ^E	2.8 (0.5) ^E	3.7 (0.7) ^E	5.0 (1.0) ^E	6.7 (1.4) ^E	8.5 (1.8) ^E	9.8 (2.1) ^E	5	F	50	0.0 (0.0)	
	31-50	146	4.9	(0.6)	F	F	2.5 (0.5) ^E	3.7 (0.6)	5.5 (0.8)	7.5 (1.3) ^E	8.9 (1.7) ^E	5	30.4 (10.0) ^E	50	0.0 (0.0)	
	51-70	184	5.2	(0.6)	F	2.6 (0.8) ^E	3.4 (0.8) ^E	4.8 (0.6)	7.0 (0.9)	9.5 (1.8) ^E	11.2 (2.5) ^E	10	F	50	0.0 (0.0)	
	>70	143	5.7	(0.5)	2.3 (0.5) ^E	2.9 (0.5) ^E	4.0 (0.6)	5.6 (0.7)	7.7 (0.9)	10.0 (1.4)	11.6 (1.8)	15	F	50	0.0 (0.0)	
	19+	584	5.2	(0.3)	1.8 (0.3)	2.3 (0.3)	3.2 (0.3)	4.6 (0.3)	6.3 (0.5)	8.4 (0.7)	9.9 (0.9)			50	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.9 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake													
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	AI ²	% >AI	(SE)	UL ³	% >UL	(SE)	
Both																		
	1-3	169	6.2	(0.4)	F	F	3.5 (0.5)	5.8 (0.5)	8.5 (0.7)	11.9 (1.3)	14.4 (1.9)	5	59.7	(5.9)	50	<3		
	4-8	281	5.8	(0.3)	2.5 (0.4)	3.1 (0.4)	4.2 (0.3)	5.5 (0.3)	7.1 (0.4)	8.7 (0.5)	9.9 (0.7)	5	60.4	(5.5)	50	0.0	(0.0)	
Male																		
	9-13	183	8.3	(0.8)	3.8 (1.0) ^E	4.5 (1.0) ^E	5.9 (0.8)	7.9 (0.8)	10.6 (1.2)	14.1 (2.3)	16.8 (3.4) ^E	5	85.1	(8.0)	50	<3		
	14-18	187	7.5	(0.5)	4.2 (1.3) ^E	4.9 (1.1) ^E	6.2 (0.9)	7.9 (0.7)	9.8 (1.2)	11.6 (1.8)	12.8 (2.3) ^E	5	89.2	(8.8)	50	0.0	(0.0)	
	19-30	223	5.6	(0.4)	3.2 (0.8) ^E	3.6 (0.8) ^E	4.4 (0.6)	5.3 (0.5)	6.3 (0.6)	7.4 (1.1)	8.1 (1.4) ^E	5	58.3	(14.4) ^E	50	0.0	(0.0)	
	31-50	229	5.8	(0.6)	2.5 (0.8) ^E	3.0 (0.8) ^E	3.9 (0.7) ^E	5.2 (0.8)	7.0 (1.1)	9.3 (2.0) ^E	10.9 (3.0) ^E	5	54.0	(13.5) ^E	50	<3		
	51-70	197	6.2	(0.8)	F	F	4.0 (1.0) ^E	5.5 (0.9) ^E	7.5 (1.3) ^E	10.0 (2.3) ^E	11.9 (3.4) ^E	10	F		50	<3		
	>70	72	7.0	(0.8)	3.5 (1.0) ^E	4.0 (1.0) ^E	5.0 (0.9) ^E	6.3 (0.8)	8.0 (1.0)	9.9 (1.5)	11.2 (1.9) ^E	15	<3		50	0.0	(0.0)	
	19+	721	5.9	(0.3)	2.2 (0.6) ^E	2.7 (0.5) ^E	3.8 (0.5)	5.3 (0.4)	7.4 (0.5)	10.0 (1.0)	12.0 (1.4)				50	0.0	(0.0)	
Female																		
	9-13	165	4.6	(0.3)	F	2.5 (0.6) ^E	3.3 (0.5)	4.5 (0.4)	5.8 (0.6)	7.3 (1.0)	8.4 (1.4)	5	38.9	(10.5) ^E	50	0.0	(0.0)	
	14-18	206	4.5	(0.4)	F	2.3 (0.6) ^E	3.1 (0.5) ^E	4.2 (0.4)	5.6 (0.6)	7.0 (1.0)	8.0 (1.4) ^E	5	34.0	(11.1) ^E	50	0.0	(0.0)	
	19-30	191	3.9	(0.4)	2.3 (0.7) ^E	2.6 (0.7) ^E	3.1 (0.6) ^E	3.8 (0.6)	4.5 (0.6)	5.3 (0.9) ^E	5.8 (1.2) ^E	5	F		50	0.0	(0.0)	
	31-50	258	5.2	(0.5)	F	2.6 (0.8) ^E	3.6 (0.7) ^E	5.1 (0.7)	6.9 (0.9)	9.0 (1.2)	10.4 (1.6)	5	51.2	(13.4) ^E	50	0.0	(0.0)	
	51-70	249	4.4	(0.4)	1.9 (0.6) ^E	2.3 (0.6) ^E	3.1 (0.5) ^E	4.2 (0.5)	5.6 (0.6)	7.2 (0.9)	8.3 (1.2)	10	F		50	0.0	(0.0)	
	>70	128	7.5	(2.0) ^E	F	F	F	6.5 (1.9) ^E	9.3 (2.5) ^E	12.5 (3.6) ^E	14.8 (4.8) ^E	15	F		50	<3		
	19+	826	5.0	(0.3)	2.3 (0.4) ^E	2.7 (0.4)	3.6 (0.4)	4.7 (0.4)	6.2 (0.4)	7.9 (0.6)	9.0 (0.9)				50	0.0	(0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.10 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)					Percentiles (<i>and SE</i>) of usual intake						% AI ² >AI (SE)			% UL ³ >UL (SE)		
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	192	6.4	(0.4)	2.3 (0.5) ^E	3.0 (0.5) ^E	4.3 (0.5)	6.1 (0.5)	8.3 (0.7)	10.5 (1.0)	11.9 (1.2)	5	66.1	(8.0)	50	0.0 (0.0)	
	4-8	321	5.6	(0.3)	2.6 (0.4) ^E	3.2 (0.4)	4.2 (0.4)	5.5 (0.3)	7.0 (0.4)	8.5 (0.6)	9.6 (0.8)	5	60.5	(6.9)	50	0.0 (0.0)	
Male																	
	9-13	226	7.2	(0.4)	3.3 (0.5)	4.0 (0.5)	5.3 (0.5)	7.0 (0.5)	9.1 (0.7)	11.6 (1.0)	13.5 (1.3)	5	78.8	(5.9)	50	0.0 (0.0)	
	14-18	262	7.9	(0.6)	3.5 (0.7) ^E	4.2 (0.7) ^E	5.7 (0.7)	7.7 (0.7)	10.3 (1.0)	13.3 (1.6)	15.5 (2.1)	5	82.7	(7.0)	50	0.0 (0.0)	
	19-30	197	6.6	(0.5)	2.6 (0.8) ^E	3.2 (0.7) ^E	4.4 (0.7)	6.3 (0.7)	8.6 (0.9)	11.0 (1.4)	12.6 (1.8)	5	67.0	(10.5)	50	0.0 (0.0)	
	31-50	282	6.4	(0.6)	F	2.9 (0.8) ^E	4.0 (0.7) ^E	5.7 (0.6)	8.0 (0.8)	11.0 (1.7)	13.4 (2.5) ^E	5	59.9	(11.7) ^E	50	<3	
	51-70	234	10.3	(1.9) ^E	F	5.9 (1.9) ^E	7.6 (1.9) ^E	9.7 (2.0) ^E	12.4 (2.4) ^E	15.6 (3.6) ^E	18.0 (4.9) ^E	10	F		50	<3	
	>70	119	6.8	(1.0)	F	F	4.0 (1.0) ^E	5.8 (1.1) ^E	8.5 (1.5) ^E	12.5 (2.9) ^E	15.9 (4.5) ^E	15	F		50	<3	
	19+	832	7.5	(0.6)	3.1 (0.4)	3.6 (0.4)	4.8 (0.5)	7.0 (0.7)	9.8 (1.0)	13.2 (1.4)	15.9 (2.0)				50	<3	
Female																	
	9-13	226	6.4	(0.6)	2.0 (0.4) ^E	2.6 (0.4)	3.8 (0.5)	5.6 (0.6)	8.2 (0.9)	11.5 (1.4)	14.1 (1.9)	5	58.3	(7.5)	50	<3	
	14-18	242	5.0	(0.4)	1.5 (0.4) ^E	1.9 (0.4) ^E	2.9 (0.5)	4.5 (0.5)	6.6 (0.6)	8.8 (0.9)	10.2 (1.1)	5	42.7	(7.7) ^E	50	0.0 (0.0)	
	19-30	208	5.8	(0.9)	2.9 (0.9) ^E	3.3 (0.9) ^E	4.3 (0.9) ^E	5.5 (0.9)	7.0 (1.0)	8.6 (1.4) ^E	9.7 (1.8) ^E	5	60.1	(18.1) ^E	50	0.0 (0.0)	
	31-50	263	4.7	(0.4)	2.1 (0.6) ^E	2.4 (0.6) ^E	3.1 (0.5) ^E	4.1 (0.5)	5.7 (0.6)	7.7 (1.1)	9.3 (1.7) ^E	5	33.8	(9.8) ^E	50	<3	
	51-70	322	4.4	(0.5)	F	1.9 (0.5) ^E	2.8 (0.5) ^E	4.2 (0.6)	6.3 (1.0)	8.9 (1.7) ^E	10.9 (2.4) ^E	10	F		50	<3	
	>70	198	4.9	(0.5)	1.8 (0.5) ^E	2.2 (0.5) ^E	3.0 (0.5)	4.3 (0.6)	6.1 (0.8)	8.4 (1.4)	10.2 (1.9) ^E	15	F		50	0.0 (0.0)	
	19+	991	4.8	(0.3)	1.8 (0.2)	2.2 (0.2)	3.0 (0.3)	4.3 (0.3)	6.4 (0.4)	9.1 (0.9)	11.4 (1.4)				50	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.11 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	Percentiles (<i>and SE</i>) of usual intake									% AI ² >AI (SE)			% UL ³ >UL (SE)										
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			(SE)			(SE)							
Both																								
	1-3	348	6.9	(0.4)	3.2	(0.6) ^E	4.0	(0.5)	5.4	(0.4)	6.9	(0.4)	8.6	(0.5)	10.4	(0.8)	11.8	(1.0)	5	80.0	(5.6)	50	0.0	(0.0)
	4-8	554	6.6	(0.3)	3.5	(0.3)	4.1	(0.3)	5.1	(0.3)	6.4	(0.3)	8.0	(0.4)	9.7	(0.5)	10.9	(0.7)	5	76.8	(4.4)	50	0.0	(0.0)
Male																								
	9-13	409	7.2	(0.3)	3.3	(0.4)	3.9	(0.4)	5.1	(0.4)	6.7	(0.4)	8.7	(0.6)	11.0	(0.8)	12.7	(1.1)	5	75.9	(5.5)	50	0.0	(0.0)
	14-18	414	9.2	(1.4)	3.3	(0.6) ^E	4.1	(0.6)	5.6	(0.7)	7.7	(0.9)	11.0	(1.4)	16.0	(2.7) ^E	20.4	(4.3) ^E	5	81.9	(6.0)	50	<3	
	19-30	311	7.2	(0.5)	2.6	(0.6) ^E	3.3	(0.6) ^E	4.7	(0.6)	6.7	(0.6)	9.2	(0.8)	12.2	(1.2)	14.3	(1.6)	5	71.0	(7.3)	50	0.0	(0.0)
	31-50	489	6.5	(0.5)	3.1	(0.4)	3.6	(0.4)	4.6	(0.4)	6.0	(0.5)	7.7	(0.6)	9.6	(1.0)	10.9	(1.3)	5	67.9	(8.5)	50	0.0	(0.0)
	51-70	575	6.8	(0.4)	3.4	(0.6) ^E	3.9	(0.6)	4.9	(0.5)	6.3	(0.5)	8.0	(0.5)	10.1	(0.9)	11.5	(1.3)	10	^F		50	0.0	(0.0)
	>70	239	7.5	(0.6)	2.8	(0.5) ^E	3.5	(0.6)	4.9	(0.6)	7.0	(0.7)	9.9	(1.0)	13.7	(1.6)	17.0	(2.2)	15	^F		50	<3	
	19+	1614	6.8	(0.3)	2.8	(0.2)	3.3	(0.2)	4.4	(0.2)	6.2	(0.3)	8.5	(0.4)	11.2	(0.6)	13.2	(0.8)				50	0.0	(0.0)
Female																								
	9-13	355	6.1	(0.4)	2.6	(0.4)	3.1	(0.4)	4.2	(0.4)	5.6	(0.5)	7.5	(0.6)	9.6	(0.9)	10.9	(1.0)	5	61.5	(8.4)	50	0.0	(0.0)
	14-18	410	5.3	(0.4)	1.5	(0.3) ^E	2.1	(0.3)	3.2	(0.3)	4.7	(0.4)	6.9	(0.7)	9.8	(1.5)	12.3	(2.3) ^E	5	45.3	(6.4)	50	<3	
	19-30	384	5.3	(0.5)	1.9	(0.3)	2.3	(0.3)	3.3	(0.4)	4.8	(0.5)	7.0	(0.7)	9.6	(1.1)	11.5	(1.4)	5	47.7	(7.7)	50	0.0	(0.0)
	31-50	585	5.2	(0.3)	2.0	(0.3)	2.5	(0.3)	3.6	(0.3)	5.1	(0.4)	7.0	(0.6)	9.4	(1.1)	11.3	(1.6)	5	51.0	(6.8)	50	<3	
	51-70	711	5.3	(0.4)	2.0	(0.2)	2.5	(0.3)	3.4	(0.3)	4.7	(0.4)	6.6	(0.6)	8.7	(0.8)	10.4	(1.1)	10	^F		50	<3	
	>70	342	5.6	(0.5)	2.6	(0.5) ^E	3.1	(0.5) ^E	3.9	(0.6)	5.0	(0.6)	6.6	(0.8)	8.6	(1.2)	10.0	(1.5)	15	<3		50	0.0	(0.0)
	19+	2022	5.3	(0.2)	2.0	(0.1)	2.5	(0.2)	3.5	(0.2)	4.9	(0.2)	6.8	(0.3)	9.3	(0.5)	11.2	(0.8)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.12 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						AI ²	% >AI		UL ³	% >UL									
					(SE)	(SE)	(SE)	(SE)	(SE)	(SE)		(SE)	(SE)		(SE)	(SE)								
Both																								
	1-3	622	6.5	(0.3)	1.5	(0.4) ^E	2.4	(0.4)	4.0	(0.3)	6.2	(0.4)	8.8	(0.4)	11.7	(0.7)	13.8	(1.0)	5	63.5	(4.0)	50	0.0	(0.0)
	4-8	919	6.0	(0.2)	2.3	(0.2)	2.9	(0.2)	4.1	(0.2)	5.5	(0.2)	7.2	(0.2)	9.2	(0.4)	10.8	(0.7)	5	58.6	(3.6)	50	<3	
Male																								
	9-13	579	7.7	(0.5)	3.4	(0.4)	4.0	(0.4)	5.4	(0.4)	7.3	(0.5)	9.7	(0.7)	12.6	(1.3)	14.9	(1.8)	5	79.8	(5.0)	50	<3	
	14-18	634	7.7	(0.3)	2.3	(0.4) ^E	3.2	(0.4)	4.9	(0.5)	7.6	(0.5)	10.0	(0.6)	13.5	(0.9)	15.5	(1.1)	5	74.1	(4.3)	50	0.0	(0.0)
	19-30	578	6.0	(0.4)	2.2	(0.5) ^E	2.7	(0.5) ^E	3.8	(0.4)	5.4	(0.4)	7.3	(0.5)	9.5	(0.9)	11.0	(1.2)	5	55.7	(6.5)	50	0.0	(0.0)
	31-50	693	5.7	(0.4)	2.1	(0.5) ^E	2.6	(0.5) ^E	3.6	(0.4)	5.1	(0.4)	7.3	(0.6)	10.0	(1.1)	12.0	(1.6)	5	51.9	(7.4)	50	0.0	(0.0)
	51-70	596	6.3	(0.5)	2.3	(0.3)	2.8	(0.4)	3.9	(0.4)	5.6	(0.6)	8.1	(0.8)	11.2	(1.2)	13.7	(1.5)	10	14.3	(4.1) ^E	50	<3	
	>70	296	6.6	(0.5)	2.8	(0.4)	3.3	(0.4)	4.4	(0.4)	5.9	(0.5)	7.9	(0.6)	10.2	(0.9)	11.9	(1.1)	15	^F		50	0.0	(0.0)
	19+	2163	6.0	(0.2)	2.1	(0.2)	2.7	(0.2)	3.7	(0.2)	5.3	(0.3)	7.6	(0.3)	10.5	(0.6)	12.7	(0.8)				50	0.0	(0.0)
Female																								
	9-13	533	5.3	(0.3)	2.4	(0.4) ^E	2.9	(0.4)	3.8	(0.3)	4.9	(0.3)	6.6	(0.5)	8.7	(1.3)	10.2	(2.4) ^E	5	48.8	(5.9)	50	<3	
	14-18	638	4.7	(0.2)	1.6	(0.2)	2.0	(0.3)	3.0	(0.3)	4.4	(0.3)	6.2	(0.4)	8.3	(0.6)	9.6	(0.8)	5	40.4	(5.2)	50	0.0	(0.0)
	19-30	499	4.5	(0.3)	1.7	(0.3) ^E	2.1	(0.3) ^E	2.9	(0.4)	4.0	(0.4)	5.5	(0.6)	7.2	(0.9)	8.4	(1.2)	5	32.5	(8.4) ^E	50	0.0	(0.0)
	31-50	716	5.0	(0.3)	1.9	(0.5) ^E	2.4	(0.5) ^E	3.2	(0.5)	4.5	(0.5)	6.3	(0.5)	8.4	(0.7)	9.9	(0.9)	5	41.6	(8.2) ^E	50	0.0	(0.0)
	51-70	745	4.8	(0.3)	1.8	(0.4) ^E	2.3	(0.3)	3.2	(0.3)	4.3	(0.3)	6.1	(0.4)	8.0	(0.7)	9.4	(0.9)	10	^F		50	0.0	(0.0)
	>70	510	6.4	(1.0)	2.1	(0.4) ^E	2.5	(0.5) ^E	3.8	(0.7) ^E	6.0	(1.1) ^E	8.5	(1.5) ^E	11.8	(2.0) ^E	14.3	(2.6) ^E	15	^F		50	<3	
	19+	2470	5.0	(0.2)	1.8	(0.2)	2.2	(0.2)	3.1	(0.2)	4.5	(0.2)	6.4	(0.3)	8.7	(0.4)	10.4	(0.6)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 27.13 Vitamin D (µg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)				Percentiles (<i>and SE</i>) of usual intake							AI ²	% >AI	(SE)	UL ³	% >UL	(SE)
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																	
	1-3	2117	6.5	(0.1)	2.1 (0.2)	2.9 (0.2)	4.4 (0.2)	6.3 (0.2)	8.5 (0.2)	10.9 (0.3)	12.5 (0.4)	5	67.0	(2.5)	50	0.0	(0.0)
	4-8	3235	6.0	(0.1)	2.5 (0.1)	3.1 (0.1)	4.2 (0.1)	5.6 (0.1)	7.4 (0.2)	9.4 (0.3)	10.8 (0.4)	5	60.7	(2.5)	50	0.0	(0.0)
Male																	
	9-13	2080	7.0	(0.2)	3.1 (0.2)	3.7 (0.2)	4.9 (0.2)	6.6 (0.2)	8.8 (0.3)	11.1 (0.4)	12.8 (0.5)	5	73.9	(2.8)	50	0.0	(0.0)
	14-18	2288	7.6	(0.2)	2.7 (0.2)	3.5 (0.2)	5.0 (0.2)	7.2 (0.3)	10.0 (0.3)	13.6 (0.6)	16.2 (0.8)	5	75.1	(2.4)	50	0.0	(0.0)
	19-30	1804	5.9	(0.2)	2.2 (0.2)	2.7 (0.2)	3.7 (0.2)	5.2 (0.3)	7.2 (0.4)	9.7 (0.7)	11.6 (1.0)	5	53.0	(4.2)	50	0.0	(0.0)
	31-50	2596	5.8	(0.2)	2.3 (0.2)	2.8 (0.2)	3.7 (0.2)	5.1 (0.2)	7.2 (0.3)	9.9 (0.6)	11.9 (0.9)	5	52.0	(4.0)	50	0.0	(0.0)
	51-70	2550	7.1	(0.5)	2.3 (0.2)	2.8 (0.2)	4.0 (0.3)	5.9 (0.4)	9.0 (0.7)	13.5 (1.2)	17.3 (1.8)	10	20.4	(3.6) ^E	50	<3	
	>70	1520	6.3	(0.4)	2.2 (0.2)	2.7 (0.2)	3.7 (0.2)	5.3 (0.4)	7.7 (0.5)	11.0 (0.8)	13.8 (1.1)	15	3.8	(1.1) ^E	50	<3	
	19+	8470	6.2	(0.2)	2.2 (0.1)	2.7 (0.1)	3.7 (0.1)	5.3 (0.2)	7.8 (0.3)	11.1 (0.5)	13.8 (0.7)				50	0.0	(0.0)
Female																	
	9-13	1980	5.7	(0.2)	2.2 (0.2)	2.8 (0.2)	3.8 (0.2)	5.2 (0.2)	7.0 (0.2)	9.2 (0.4)	10.7 (0.5)	5	53.0	(3.1)	50	0.0	(0.0)
	14-18	2256	5.0	(0.2)	1.5 (0.1)	2.0 (0.2)	3.0 (0.2)	4.4 (0.2)	6.4 (0.3)	8.9 (0.4)	10.7 (0.7)	5	40.9	(2.9)	50	0.0	(0.0)
	19-30	1854	4.7	(0.2)	1.7 (0.2)	2.2 (0.2)	3.0 (0.2)	4.2 (0.2)	5.8 (0.3)	7.8 (0.4)	9.3 (0.6)	5	36.0	(3.5)	50	0.0	(0.0)
	31-50	2686	5.2	(0.3)	1.9 (0.2)	2.3 (0.2)	3.2 (0.2)	4.5 (0.3)	6.6 (0.5)	9.6 (1.0)	12.0 (1.5)	5	42.0	(4.4)	50	<3	
	51-70	3200	5.0	(0.3)	1.7 (0.2)	2.1 (0.2)	3.0 (0.2)	4.4 (0.3)	6.7 (0.4)	9.8 (0.9)	12.3 (1.3)	10	9.3	(2.5) ^E	50	<3	
	>70	2610	5.3	(0.3)	2.0 (0.2)	2.5 (0.2)	3.4 (0.2)	4.7 (0.3)	6.8 (0.4)	9.4 (0.6)	11.5 (0.9)	15	^F		50	0.0	(0.0)
	19+	10350	5.0	(0.1)	1.8 (0.1)	2.2 (0.1)	3.1 (0.1)	4.4 (0.1)	6.4 (0.2)	9.2 (0.4)	11.5 (0.7)				50	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

28. Zinc (mg/d): Usual intakes from food

Table 28.1 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	79	7.4	(0.6)	5.1 (0.8)	5.5 (0.8)	6.3 (0.7)	7.2 (0.7)	8.2 (0.8)	9.1 (1.0)	9.7 (1.1)	2.5	<3	7	F	
	4-8	127	8.9	(0.5)	7.0 (0.9)	7.4 (0.9)	8.0 (0.8)	8.8 (0.7)	9.6 (0.8)	10.3 (1.1)	10.8 (1.4)	4.0	<3	12	F	
Male																
	9-13	111	12.7	(1.2)	9.2 (1.4)	9.8 (1.4)	11.1 (1.3)	12.7 (1.3)	14.4 (1.8)	16.2 (2.4)	17.4 (2.9) ^E	7.0	F	23	F	
	14-18	107	12.9	(0.9)	9.8 (1.7) ^E	10.5 (1.6)	11.8 (1.4)	13.5 (1.2)	15.4 (1.4)	17.4 (2.1)	18.8 (2.8)	8.5	F	34	<3	
	19-30	77	13.0	(0.9)	8.1 (1.1)	9.0 (1.2)	10.8 (1.2)	13.1 (1.3)	15.5 (1.4)	17.8 (1.6)	19.3 (1.9)	9.4	F	40	<3	
	31-50	145	13.1	(1.1)	7.9 (1.4) ^E	8.7 (1.3)	10.2 (1.1)	12.1 (1.2)	14.3 (1.5)	16.7 (2.1)	18.4 (2.7)	9.4	F	40	<3	
	51-70	182	10.5	(0.7)	6.4 (0.5)	7.0 (0.6)	8.2 (0.7)	9.8 (0.8)	11.8 (0.9)	13.9 (1.2)	15.3 (1.3)	9.4	43.0 (11.8) ^E	40	0.0 (0.0)	
	>70	63	11.8	(1.2)	9.1 (1.4)	9.8 (1.5)	11.0 (1.7)	12.5 (2.0)	14.3 (2.4) ^E	16.1 (2.8) ^E	17.3 (3.0) ^E	9.4	F	40	0.0 (0.0)	
	19+	467	12.2	(0.5)	8.3 (0.9)	9.0 (0.9)	10.3 (0.7)	11.9 (0.6)	13.8 (0.8)	15.7 (1.2)	16.9 (1.5)	9.4	F	40	0.0 (0.0)	
Female																
	9-13	96	8.9	(0.5)	5.4 (0.9)	6.0 (0.8)	7.2 (0.7)	8.7 (0.7)	10.5 (0.8)	12.6 (1.1)	14.0 (1.4)	7.0	F	23	<3	
	14-18	105	9.3	(0.8)	5.9 (1.1) ^E	6.4 (1.0)	7.5 (0.9)	8.9 (0.9)	10.7 (1.2)	12.7 (1.6)	14.1 (2.1)	7.3	F	34	<3	
	19-30	91	8.6	(0.6)	6.0 (0.9)	6.4 (0.8)	7.3 (0.8)	8.4 (0.8)	9.5 (0.9)	10.7 (1.2)	11.5 (1.5)	6.8	F	40	0.0 (0.0)	
	31-50	167	9.4	(0.8)	5.5 (1.0) ^E	6.2 (1.0)	7.6 (0.9)	9.4 (1.0)	11.5 (1.2)	13.8 (1.6)	15.3 (2.0)	6.8	F	40	<3	
	51-70	198	10.1	(1.3)	6.3 (0.6)	7.1 (0.7)	8.2 (0.8)	9.7 (1.1)	11.8 (1.6)	14.0 (2.1)	15.7 (2.6) ^E	6.8	F	40	0.0 (0.0)	
	>70	74	9.0	(0.8)	5.0 (0.7)	5.6 (0.8)	6.8 (0.8)	8.4 (1.0)	10.5 (1.2)	12.7 (1.6)	14.3 (1.8)	6.8	F	40	0.0 (0.0)	
	19+	530	9.4	(0.5)	6.6 (0.9)	7.2 (0.8)	8.2 (0.7)	9.5 (0.6)	11.1 (0.7)	12.6 (1.0)	13.7 (1.3)	6.8	F	40	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.2 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)				
Both																
	1-3	58	7.8	(0.6)	6.8 (0.8)	7.0 (0.7)	7.4 (0.7)	7.8 (0.7)	8.3 (0.8)	8.7 (1.1)	9.0 (1.3)	2.5	0.0 (0.0)	7	90.6	(24.9) ^E
	4-8	110	8.5	(0.4)	7.2 (0.8)	7.5 (0.7)	7.9 (0.5)	8.4 (0.4)	8.9 (0.5)	9.3 (0.7)	9.6 (0.9)	4.0	<3	12	<3	
Male																
	9-13	95	11.8	(0.9)	9.6 (1.6)	10.1 (1.5)	11.0 (1.4)	12.0 (1.3)	13.2 (1.4)	14.3 (1.5)	15.0 (1.7)	7.0	F	23	<3	
	14-18	87	14.3	(1.0)	7.8 (1.7) ^E	9.1 (1.5) ^E	11.4 (1.2)	14.2 (1.2)	17.8 (1.7)	22.1 (2.8)	25.4 (3.8)	8.5	F	34	<3	
	19-30	70	15.0	(1.0)	9.0 (1.9) ^E	10.4 (1.7) ^E	12.8 (1.5)	15.7 (1.3)	18.8 (1.4)	21.7 (1.7)	23.6 (2.1)	9.4	F	40	<3	
	31-50	109	12.3	(0.8)	7.5 (1.2)	8.2 (1.2)	9.5 (1.1)	11.4 (1.0)	13.7 (1.3)	16.2 (1.9)	17.8 (2.4)	9.4	F	40	<3	
	51-70	128	11.4	(0.7)	6.9 (1.2) ^E	7.6 (1.1)	9.0 (0.8)	10.8 (0.7)	12.9 (1.0)	15.2 (1.5)	16.8 (2.0)	9.4	F	40	0.0 (0.0)	
	>70	65	9.4	(0.6)	6.0 (1.0)	6.6 (0.9)	7.5 (0.8)	8.8 (0.8)	10.3 (1.0)	11.8 (1.4)	12.8 (1.7)	9.4	61.3 (18.5) ^E	40	0.0 (0.0)	
	19+	372	12.3	(0.4)	7.2 (0.6)	8.0 (0.6)	9.6 (0.6)	11.8 (0.6)	14.5 (0.8)	17.2 (1.1)	19.0 (1.3)	9.4	22.7 (6.0) ^E	40	0.0 (0.0)	
Female																
	9-13	75	8.9	(0.6)	5.6 (0.8)	6.1 (0.8)	7.2 (0.9)	8.6 (1.0)	10.1 (1.1)	11.6 (1.3)	12.7 (1.4)	7.0	F	23	<3	
	14-18	81	8.6	(0.7)	4.5 (1.2) ^E	5.3 (1.1) ^E	6.7 (0.9)	8.4 (0.9)	10.0 (1.0)	11.5 (1.2)	12.5 (1.3)	7.3	F	34	0.0 (0.0)	
	19-30	101	9.7	(0.7)	6.4 (1.0)	7.2 (1.0)	8.6 (0.9)	10.3 (0.9)	12.1 (1.1)	14.0 (1.5)	15.2 (1.8)	6.8	F	40	0.0 (0.0)	
	31-50	116	9.6	(0.8)	5.8 (0.9)	6.5 (0.8)	7.9 (0.8)	9.6 (0.9)	11.5 (1.3)	13.3 (1.7)	14.5 (1.9)	6.8	F	40	0.0 (0.0)	
	51-70	146	9.9	(0.6)	5.3 (0.7)	6.1 (0.7)	7.6 (0.7)	9.7 (0.7)	12.1 (0.9)	14.5 (1.3)	16.1 (1.6)	6.8	F	40	0.0 (0.0)	
	>70	94	7.7	(0.6)	4.5 (0.3)	5.0 (0.3)	6.0 (0.4)	7.4 (0.6)	9.1 (0.8)	11.1 (1.2)	12.6 (1.5)	6.8	39.7 (9.5) ^E	40	0.0 (0.0)	
	19+	457	9.5	(0.4)	5.5 (0.4)	6.2 (0.4)	7.7 (0.4)	9.6 (0.5)	11.6 (0.7)	13.6 (0.9)	15.0 (1.1)	6.8	15.4 (3.9) ^E	40	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.3 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	%	(SE)	UL ³	%	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)			<EAR	
Both																
	1-3	112	6.9	(0.4)	5.4 (0.4)	5.7 (0.4)	6.2 (0.4)	6.9 (0.4)	7.6 (0.5)	8.4 (0.5)	8.9 (0.6)	2.5	0.0	(0.0)	7	F
	4-8	177	9.2	(0.5)	5.8 (0.7)	6.4 (0.7)	7.4 (0.7)	8.7 (0.7)	10.2 (0.7)	11.8 (0.8)	12.8 (0.9)	4.0	<3		12	F
Male																
	9-13	111	11.4	(0.7)	6.6 (0.6)	7.2 (0.6)	8.5 (0.7)	10.5 (0.7)	13.0 (1.0)	15.7 (1.4)	17.7 (2.0)	7.0	F		23	<3
	14-18	113	12.9	(1.3)	7.7 (1.8) ^E	8.5 (1.8) ^E	10.2 (1.8) ^E	12.3 (1.9)	15.0 (2.4)	17.7 (3.0) ^E	19.6 (3.5) ^E	8.5	F		34	<3
	19-30	91	14.4	(1.1)	8.2 (1.9) ^E	9.3 (1.7) ^E	11.5 (1.5)	14.1 (1.5)	17.1 (1.8)	20.0 (2.5)	21.9 (3.1)	9.4	F		40	<3
	31-50	101	14.2	(1.2)	8.9 (1.6) ^E	9.9 (1.4)	11.6 (1.3)	14.0 (1.5)	16.9 (2.2)	20.3 (3.4)	22.8 (4.4) ^E	9.4	F		40	<3
	51-70	134	13.0	(1.0)	8.2 (1.6) ^E	9.0 (1.4)	10.5 (1.2)	12.4 (1.1)	14.6 (1.3)	16.9 (1.9)	18.4 (2.5)	9.4	F		40	<3
	>70	56	10.0	(0.8)	7.6 (1.1)	8.1 (1.1)	9.0 (1.1)	10.1 (1.1)	11.3 (1.1)	12.5 (1.1)	13.2 (1.2)	9.4	F		40	0.0 (0.0)
	19+	382	13.5	(0.6)	7.7 (0.7)	8.7 (0.7)	10.6 (0.7)	13.1 (0.7)	16.1 (1.1)	19.5 (1.6)	21.8 (2.1)	9.4	14.8	(4.8) ^E	40	<3
Female																
	9-13	105	9.0	(0.9)	5.7 (0.9)	6.2 (0.9)	7.1 (0.8)	8.4 (0.8)	10.0 (1.0)	11.8 (1.5)	13.1 (1.9)	7.0	F		23	<3
	14-18	120	8.3	(0.9)	3.9 (1.3) ^E	4.7 (1.2) ^E	6.2 (1.0)	8.0 (0.9)	10.2 (1.2)	12.8 (1.6)	14.7 (2.1)	7.3	F		34	<3
	19-30	91	9.7	(0.6)	6.1 (0.9)	6.9 (0.7)	8.1 (0.6)	9.5 (0.8)	11.7 (1.4)	13.9 (2.1)	15.1 (2.5)	6.8	F		40	0.0 (0.0)
	31-50	159	10.9	(0.7)	5.2 (0.8)	6.2 (0.8)	8.3 (0.8)	10.9 (0.9)	13.7 (1.1)	16.5 (1.4)	18.4 (1.7)	6.8	F		40	<3
	51-70	174	9.1	(0.5)	6.7 (1.0)	7.2 (0.9)	8.0 (0.7)	9.0 (0.6)	10.1 (0.7)	11.2 (0.9)	11.9 (1.2)	6.8	F		40	0.0 (0.0)
	>70	80	8.6	(0.7)	5.3 (1.1) ^E	6.0 (1.1) ^E	7.2 (1.0)	8.7 (1.0)	10.3 (1.2)	12.0 (1.6)	13.1 (1.9)	6.8	F		40	0.0 (0.0)
	19+	504	9.8	(0.3)	5.5 (0.4)	6.3 (0.4)	7.8 (0.4)	9.8 (0.5)	12.0 (0.6)	14.2 (0.8)	15.6 (0.9)	6.8	14.7	(4.2) ^E	40	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.4 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	99	8.2	(0.5)	5.8 (0.8)	6.3 (0.8)	7.2 (0.6)	8.4 (0.6)	9.7 (0.8)	11.0 (1.3)	11.9 (1.6)	2.5	<3	7	79.2	(12.4)
	4-8	140	9.7	(0.7)	6.5 (0.9)	7.1 (0.9)	8.2 (0.8)	9.7 (0.7)	11.3 (0.9)	13.1 (1.3)	14.4 (1.7)	4.0	<3	12	F	
Male																
	9-13	92	12.0	(1.2)	8.4 (1.5) ^E	9.0 (1.4)	10.1 (1.3)	11.4 (1.3)	12.9 (1.5)	14.4 (1.9)	15.3 (2.2)	7.0	F	23	<3	
	14-18	107	14.6	(1.0)	8.7 (1.2)	9.8 (1.2)	11.9 (1.2)	14.8 (1.2)	18.5 (2.0)	22.5 (2.9)	25.4 (3.6)	8.5	F	34	F	
	19-30	73	16.9	(1.8)	10.5 (2.3) ^E	11.7 (2.2) ^E	14.0 (2.1)	17.0 (2.3)	20.6 (3.0)	24.5 (4.2) ^E	27.2 (5.2) ^E	9.4	F	40	<3	
	31-50	134	13.0	(1.0)	8.9 (1.5) ^E	9.6 (1.4)	10.8 (1.2)	12.4 (1.1)	14.3 (1.5)	16.3 (2.6)	17.7 (3.6) ^E	9.4	F	40	<3	
	51-70	131	12.0	(1.0)	7.1 (1.4) ^E	7.9 (1.3)	9.3 (1.1)	11.1 (1.0)	13.3 (1.2)	15.6 (1.7)	17.3 (2.2)	9.4	F	40	<3	
	>70	55	11.6	(1.3)	5.9 (1.3) ^E	6.7 (1.2) ^E	8.2 (1.2)	10.2 (1.3)	12.6 (1.7)	15.2 (2.4)	17.0 (3.0) ^E	9.4	F	40	<3	
	19+	393	13.4	(0.6)	7.6 (0.9)	8.5 (0.9)	10.2 (0.8)	12.7 (0.7)	15.7 (0.9)	19.0 (1.6)	21.4 (2.1)	9.4	F	40	<3	
Female																
	9-13	79	9.9	(0.9)	5.9 (1.0) ^E	6.6 (1.0)	8.1 (1.0)	10.1 (1.1)	12.3 (1.4)	14.6 (2.0)	16.3 (2.5)	7.0	F	23	<3	
	14-18	104	8.8	(0.5)	6.6 (0.6)	7.0 (0.6)	7.9 (0.6)	8.8 (0.7)	9.9 (0.9)	11.1 (1.1)	12.0 (1.2)	7.3	F	34	0.0	(0.0)
	19-30	101	10.3	(1.9) ^E	6.8 (1.8) ^E	7.5 (1.8) ^E	8.7 (1.8) ^E	10.7 (2.1) ^E	13.4 (3.1) ^E	F	F	6.8	F	40	F	
	31-50	143	9.1	(0.7)	6.1 (1.3) ^E	6.6 (1.2) ^E	7.4 (1.0)	8.5 (0.8)	9.7 (1.1)	11.0 (1.8) ^E	11.8 (2.5) ^E	6.8	F	40	<3	
	51-70	193	8.3	(0.4)	6.3 (0.9)	6.6 (0.9)	7.2 (0.7)	8.0 (0.6)	8.7 (0.6)	9.5 (0.9)	9.9 (1.1)	6.8	F	40	0.0	(0.0)
	>70	94	8.3	(0.6)	4.7 (0.9) ^E	5.3 (0.9) ^E	6.4 (0.8)	7.9 (0.9)	9.6 (1.0)	11.4 (1.3)	12.6 (1.6)	6.8	F	40	0.0	(0.0)
	19+	531	9.0	(0.5)	5.1 (0.3)	5.7 (0.3)	6.8 (0.3)	8.3 (0.4)	10.4 (0.6)	12.8 (1.2)	14.6 (2.0)	6.8	24.5	(5.3) ^E	40	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.5 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)								
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)												
Both																								
	1-3	311	8.1	(0.3)	5.3	(0.6)	5.9	(0.6)	6.9	(0.5)	8.2	(0.5)	9.5	(0.5)	10.9	(0.7)	11.7	(0.8)	2.5	<3		7	73.1	(9.6)
	4-8	485	10.3	(0.4)	7.6	(0.9)	8.1	(0.8)	9.1	(0.6)	10.3	(0.5)	11.6	(0.6)	12.9	(1.0)	13.7	(1.2)	4.0	<3		12		F
Male																								
	9-13	277	13.1	(0.6)	9.8	(1.3)	10.6	(1.1)	12.0	(0.9)	13.8	(0.9)	15.7	(1.1)	17.6	(1.5)	18.8	(1.8)	7.0	<3		23	<3	
	14-18	339	16.2	(1.0)	9.4	(1.2)	10.7	(1.1)	13.2	(1.1)	16.5	(1.2)	20.8	(1.7)	25.8	(2.6)	29.4	(3.3)	8.5	F		34		F
	19-30	237	13.8	(0.7)	9.3	(0.7)	10.2	(0.7)	11.9	(0.8)	14.1	(0.9)	16.4	(1.1)	18.7	(1.3)	20.2	(1.5)	9.4	F		40	0.0	(0.0)
	31-50	423	13.9	(0.7)	8.7	(1.1)	9.7	(1.0)	11.5	(0.8)	13.7	(0.8)	16.3	(1.0)	19.1	(1.5)	21.0	(2.0)	9.4	F		40	<3	
	51-70	387	12.0	(0.5)	7.3	(0.9)	8.1	(0.8)	9.6	(0.7)	11.6	(0.6)	13.9	(0.8)	16.4	(1.3)	18.2	(1.8)	9.4	F		40	<3	
	>70	132	10.0	(0.7)	5.5	(1.2) ^E	6.3	(1.2) ^E	7.8	(1.1)	9.8	(1.1)	12.1	(1.3)	14.7	(1.6)	16.6	(1.9)	9.4	45.5	(14.9) ^E	40	<3	
	19+	1179	13.0	(0.4)	8.0	(0.5)	8.9	(0.5)	10.6	(0.4)	12.8	(0.5)	15.4	(0.6)	18.1	(0.8)	20.0	(1.0)	9.4	13.6	(3.4) ^E	40	0.0	(0.0)
Female																								
	9-13	281	10.3	(0.5)	6.2	(0.8)	6.9	(0.7)	8.1	(0.6)	9.7	(0.6)	11.5	(0.7)	13.5	(1.0)	14.9	(1.3)	7.0	F		23	<3	
	14-18	321	9.8	(0.4)	6.4	(0.5)	7.1	(0.5)	8.3	(0.5)	9.8	(0.5)	11.6	(0.6)	13.5	(0.8)	14.8	(1.0)	7.3	F		34	0.0	(0.0)
	19-30	249	9.9	(0.5)	6.4	(0.5)	7.0	(0.5)	8.2	(0.5)	9.7	(0.6)	11.3	(0.7)	12.9	(0.9)	13.9	(1.0)	6.8	F		40	0.0	(0.0)
	31-50	364	10.4	(0.5)	6.3	(0.6)	6.9	(0.6)	8.1	(0.6)	9.9	(0.6)	12.4	(0.8)	15.1	(1.4)	17.2	(2.0)	6.8	F		40	<3	
	51-70	467	10.2	(0.4)	7.4	(0.8)	8.0	(0.7)	9.1	(0.5)	10.4	(0.5)	11.9	(0.6)	13.5	(1.0)	14.5	(1.3)	6.8	F		40	0.0	(0.0)
	>70	215	8.0	(0.3)	6.5	(0.7)	6.9	(0.6)	7.5	(0.6)	8.3	(0.5)	9.1	(0.6)	9.9	(0.9)	10.4	(1.1)	6.8	F		40	0.0	(0.0)
	19+	1295	10.0	(0.2)	6.6	(0.4)	7.3	(0.4)	8.4	(0.3)	10.0	(0.3)	11.7	(0.4)	13.6	(0.6)	14.9	(0.8)	6.8	F		40	0.0	(0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.6 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)	UL ³	% >UL (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)				
Both														
	1-3	644	7.2 (0.2)	4.5 (0.4)	5.0 (0.3)	6.0 (0.2)	7.2 (0.2)	8.5 (0.3)	9.9 (0.5)	10.8 (0.7)	2.5	<3	7	53.7 (4.6)
	4-8	956	9.1 (0.4)	6.1 (0.6)	6.7 (0.5)	7.6 (0.4)	8.7 (0.2)	10.2 (0.3)	11.8 (0.7)	12.9 (1.0)	4.0	<3	12	F
Male														
	9-13	589	11.4 (0.3)	8.0 (1.1)	8.7 (0.9)	9.8 (0.7)	11.2 (0.4)	12.9 (0.6)	14.5 (1.1)	15.6 (1.5)	7.0	F	23	<3
	14-18	639	14.1 (0.4)	7.8 (0.8)	8.9 (0.7)	10.9 (0.6)	13.7 (0.5)	17.3 (0.7)	21.2 (1.2)	24.0 (1.6)	8.5	F	34	<3
	19-30	481	14.0 (0.6)	9.8 (1.5)	10.6 (1.3)	12.0 (1.0)	13.7 (0.7)	15.6 (1.0)	17.6 (1.7)	18.8 (2.3)	9.4	F	40	0.0 (0.0)
	31-50	709	13.7 (0.5)	11.6 (2.1) ^E	12.1 (1.9)	13.0 (1.4)	14.1 (0.7)	15.2 (0.9)	16.4 (2.0)	17.1 (2.9) ^E	9.4	F	40	<3
	51-70	758	12.0 (0.4)	6.6 (0.8)	7.4 (0.7)	9.1 (0.6)	11.4 (0.5)	14.0 (0.6)	17.1 (1.1)	19.5 (1.6)	9.4	28.1 (6.7) ^E	40	<3
	>70	734	10.2 (0.3)	6.9 (1.0)	7.4 (0.8)	8.5 (0.6)	9.9 (0.4)	11.4 (0.5)	13.1 (1.0)	14.1 (1.3)	9.4	41.4 (11.1) ^E	40	0.0 (0.0)
	19+	2682	13.0 (0.3)	7.9 (0.7)	8.8 (0.6)	10.5 (0.5)	12.7 (0.3)	15.3 (0.4)	18.2 (0.8)	20.2 (1.1)	9.4	14.5 (4.5) ^E	40	<3
Female														
	9-13	585	9.2 (0.2)	5.5 (0.5)	6.2 (0.4)	7.4 (0.3)	8.9 (0.3)	10.6 (0.4)	12.5 (0.7)	13.8 (0.9)	7.0	19.9 (4.9) ^E	23	<3
	14-18	645	9.8 (0.3)	6.0 (0.7)	6.7 (0.6)	8.0 (0.5)	9.7 (0.3)	11.8 (0.5)	14.0 (0.9)	15.4 (1.2)	7.3	F	34	0.0 (0.0)
	19-30	514	8.9 (0.3)	4.7 (0.5)	5.4 (0.5)	6.6 (0.4)	8.3 (0.4)	10.4 (0.5)	12.7 (0.8)	14.4 (1.1)	6.8	27.3 (6.2) ^E	40	0.0 (0.0)
	31-50	758	9.3 (0.3)	6.0 (0.7)	6.7 (0.6)	7.8 (0.4)	9.1 (0.3)	10.8 (0.4)	12.5 (0.8)	13.6 (1.1)	6.8	F	40	0.0 (0.0)
	51-70	955	9.8 (0.3)	5.5 (0.8)	6.2 (0.7)	7.6 (0.5)	9.3 (0.4)	11.5 (0.5)	13.9 (1.0)	15.5 (1.4)	6.8	F	40	0.0 (0.0)
	>70	1345	8.5 (0.2)	4.5 (0.3)	5.1 (0.3)	6.4 (0.2)	8.0 (0.2)	10.1 (0.3)	12.4 (0.5)	14.2 (0.7)	6.8	31.6 (3.7)	40	0.0 (0.0)
	19+	3572	9.3 (0.2)	5.1 (0.2)	5.8 (0.2)	7.1 (0.2)	8.8 (0.2)	11.0 (0.2)	13.3 (0.4)	15.0 (0.5)	6.8	21.0 (2.5)	40	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.7 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR		(SE)	UL ³	% >UL		(SE)						
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)												
Both																								
	1-3	324	6.9	(0.2)	4.5	(0.4)	4.9	(0.4)	5.7	(0.3)	6.9	(0.3)	8.2	(0.5)	9.6	(0.7)	10.5	(0.9)	2.5	<3		7	48.4	(8.4) ^E
	4-8	425	8.6	(0.5)	6.6	(0.4)	7.0	(0.4)	7.7	(0.5)	8.5	(0.5)	9.5	(0.6)	10.5	(0.7)	11.1	(0.7)	4.0	0.0	(0.0)	12		^F
Male																								
	9-13	274	11.3	(0.5)	9.7	(0.4)	10.1	(0.5)	10.7	(0.5)	11.4	(0.5)	12.2	(0.6)	13.0	(0.7)	13.5	(0.7)	7.0	0.0	(0.0)	23	0.0	(0.0)
	14-18	297	14.9	(0.6)	8.4	(1.2)	9.6	(1.1)	11.7	(0.9)	14.5	(0.8)	17.9	(1.0)	21.8	(1.6)	24.7	(2.1)	8.5		^F	34	<3	
	19-30	249	14.8	(1.0)	9.6	(0.7)	10.4	(0.7)	12.0	(0.8)	14.0	(1.0)	16.3	(1.2)	18.6	(1.4)	20.2	(1.5)	9.4		^F	40	0.0	(0.0)
	31-50	309	13.0	(0.7)	7.4	(1.2) ^E	8.3	(1.1)	10.0	(0.9)	12.3	(0.7)	15.0	(1.0)	18.1	(1.8)	20.4	(2.5)	9.4		^F	40	<3	
	51-70	277	12.8	(0.6)	7.5	(1.3) ^E	8.4	(1.1)	10.1	(0.9)	12.3	(0.7)	15.0	(1.0)	17.9	(1.7)	19.8	(2.2)	9.4		^F	40	<3	
	>70	136	11.4	(1.1)	4.6	(1.3) ^E	5.6	(1.3) ^E	7.7	(1.2)	10.5	(1.2)	14.3	(1.8)	19.4	(3.1)	23.2	(4.2) ^E	9.4	40.2	(11.5) ^E	40	<3	
	19+	971	13.2	(0.4)	7.5	(0.7)	8.5	(0.6)	10.1	(0.5)	12.4	(0.5)	15.3	(0.6)	18.4	(1.0)	20.7	(1.4)	9.4	17.6	(4.9) ^E	40	<3	
Female																								
	9-13	265	9.4	(0.4)	6.3	(1.0)	6.9	(0.9)	7.9	(0.7)	9.2	(0.5)	10.7	(0.6)	12.3	(1.1)	13.3	(1.6)	7.0		^F	23	<3	
	14-18	290	9.1	(0.5)	5.7	(0.3)	6.3	(0.3)	7.2	(0.4)	8.6	(0.5)	10.5	(0.6)	12.4	(0.7)	13.7	(0.8)	7.3	26.0	(6.2) ^E	34	0.0	(0.0)
	19-30	197	10.1	(0.8)	6.2	(0.5)	6.9	(0.5)	8.1	(0.5)	9.5	(0.6)	11.4	(0.8)	13.3	(1.0)	14.6	(1.2)	6.8		^F	40	0.0	(0.0)
	31-50	312	10.6	(0.7)	6.3	(0.9)	6.9	(0.9)	8.3	(0.8)	10.1	(0.7)	12.3	(1.0)	14.7	(1.8)	16.5	(2.4)	6.8		^F	40	<3	
	51-70	312	9.3	(0.5)	6.2	(0.8)	6.7	(0.8)	7.8	(0.6)	9.2	(0.5)	10.7	(0.6)	12.3	(1.0)	13.4	(1.3)	6.8		^F	40	0.0	(0.0)
	>70	239	7.9	(0.4)	4.5	(0.7)	5.1	(0.7)	6.2	(0.5)	7.7	(0.4)	9.5	(0.6)	11.5	(1.1)	12.8	(1.4)	6.8	34.8	(9.3) ^E	40	0.0	(0.0)
	19+	1060	9.8	(0.3)	5.4	(0.4)	6.1	(0.4)	7.4	(0.4)	9.2	(0.3)	11.5	(0.4)	14.2	(0.8)	16.2	(1.2)	6.8	17.4	(4.3) ^E	40	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.8 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)				
Both																
	1-3	129	7.4	(0.4)	4.9 (0.6)	5.4 (0.6)	6.3 (0.5)	7.4 (0.5)	8.8 (0.6)	10.2 (0.9)	11.1 (1.1)	2.5	<3	7	60.0	(12.3) ^E
	4-8	213	8.8	(0.3)	6.0 (0.6)	6.6 (0.5)	7.5 (0.4)	8.7 (0.4)	10.0 (0.4)	11.2 (0.6)	12.1 (0.8)	4.0	<3	12	^F	
Male																
	9-13	122	12.1	(0.8)	10.2 (0.9)	10.6 (0.9)	11.3 (1.0)	12.1 (1.1)	13.1 (1.2)	14.1 (1.4)	14.7 (1.5)	7.0	<3	23	0.0	(0.0)
	14-18	150	15.0	(0.9)	8.3 (1.0)	9.5 (1.0)	11.7 (1.0)	14.5 (1.0)	17.7 (1.3)	21.2 (1.7)	23.7 (2.1)	8.5	^F	34	<3	
	19-30	106	14.7	(1.0)	8.0 (1.8) ^E	9.1 (1.7) ^E	11.2 (1.4)	14.0 (1.3)	17.3 (1.7)	20.8 (2.4)	23.2 (3.1)	9.4	^F	40	<3	
	31-50	155	13.4	(0.9)	6.6 (1.5) ^E	7.8 (1.4) ^E	10.1 (1.1)	12.9 (1.0)	16.3 (1.3)	20.0 (2.1)	22.6 (2.7)	9.4	^F	40	<3	
	51-70	122	12.6	(0.8)	9.0 (1.7) ^E	9.8 (1.5)	11.1 (1.2)	12.8 (1.1)	14.6 (1.5)	16.3 (2.4)	17.5 (3.1) ^E	9.4	^F	40	<3	
	>70	88	11.8	(0.7)	7.4 (1.0)	8.1 (0.9)	9.6 (0.8)	11.5 (0.8)	13.9 (1.1)	16.4 (1.7)	18.2 (2.2)	9.4	^F	40	<3	
	19+	471	13.3	(0.5)	7.0 (0.6)	8.2 (0.6)	10.2 (0.6)	12.9 (0.6)	16.0 (0.8)	19.5 (1.1)	21.9 (1.4)	9.4	18.1 (4.4) ^E	40	<3	
Female																
	9-13	103	9.9	(0.6)	6.9 (0.9)	7.4 (0.8)	8.4 (0.7)	9.6 (0.7)	11.1 (0.9)	12.6 (1.2)	13.7 (1.5)	7.0	^F	23	<3	
	14-18	142	10.1	(0.6)	6.4 (0.8)	7.1 (0.7)	8.3 (0.7)	9.8 (0.8)	11.5 (1.0)	13.1 (1.3)	14.1 (1.5)	7.3	^F	34	0.0	(0.0)
	19-30	111	9.2	(0.5)	5.6 (0.9)	6.2 (0.8)	7.4 (0.8)	8.8 (0.9)	10.4 (0.9)	12.0 (1.1)	13.1 (1.3)	6.8	^F	40	0.0	(0.0)
	31-50	146	11.0	(0.7)	5.4 (0.9) ^E	6.2 (0.9)	7.7 (0.8)	9.7 (0.8)	12.0 (0.9)	14.4 (1.2)	16.0 (1.6)	6.8	^F	40	<3	
	51-70	184	10.1	(0.6)	7.7 (0.4)	8.2 (0.5)	9.1 (0.6)	10.2 (0.7)	11.5 (0.9)	12.7 (1.1)	13.5 (1.2)	6.8	<3	40	0.0	(0.0)
	>70	143	10.4	(1.6)	5.4 (0.7)	6.0 (0.7)	7.1 (0.8)	8.9 (1.1)	12.1 (2.1) ^E	17.8 (4.9) ^E	^F	6.8	^F	40	^F	
	19+	584	10.3	(0.4)	6.1 (0.5)	6.8 (0.5)	8.0 (0.5)	9.6 (0.4)	11.7 (0.5)	14.1 (1.0)	15.9 (1.5)	6.8	^F	40	<3	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.9 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)				
Both																
	1-3	169	7.1	(0.3)	3.8 (0.4)	4.3 (0.4)	5.4 (0.4)	6.7 (0.4)	8.2 (0.5)	10.2 (0.7)	11.7 (1.0)	2.5	<3	7	44.1	(7.0)
	4-8	281	8.9	(0.4)	6.5 (0.3)	6.9 (0.3)	7.7 (0.4)	8.6 (0.4)	9.8 (0.5)	10.9 (0.7)	11.7 (0.8)	4.0	0.0 (0.0)	12	F	
Male																
	9-13	183	13.8	(1.5)	10.1 (1.8) ^E	10.8 (1.7)	12.0 (1.4)	13.7 (1.3)	15.6 (1.8)	17.8 (3.1) ^E	19.3 (4.3) ^E	7.0	F	23	F	
	14-18	187	14.8	(0.8)	9.9 (1.9) ^E	10.8 (1.7)	12.6 (1.3)	14.8 (1.0)	17.4 (1.3)	20.0 (2.0)	21.7 (2.6)	8.5	F	34	<3	
	19-30	223	14.4	(0.8)	9.7 (0.7)	10.6 (0.7)	12.2 (0.7)	14.1 (0.8)	16.3 (1.0)	18.4 (1.2)	19.8 (1.3)	9.4	F	40	0.0 (0.0)	
	31-50	229	13.8	(1.0)	7.1 (0.5)	8.1 (0.5)	10.0 (0.6)	12.7 (0.9)	16.4 (1.3)	20.6 (1.7)	23.4 (2.0)	9.4	19.8 (5.0) ^E	40	<3	
	51-70	197	11.7	(0.6)	6.8 (1.2) ^E	7.7 (1.1)	9.5 (0.9)	11.6 (0.7)	13.9 (0.9)	16.2 (1.4)	17.7 (1.8)	9.4	F	40	0.0 (0.0)	
	>70	72	10.9	(0.7)	8.4 (1.2)	8.9 (1.1)	9.8 (1.0)	10.8 (1.0)	11.9 (1.1)	13.0 (1.5)	13.8 (1.9)	9.4	F	40	<3	
	19+	721	13.2	(0.5)	7.6 (0.6)	8.6 (0.6)	10.4 (0.5)	12.7 (0.5)	15.6 (0.7)	19.2 (1.1)	21.6 (1.6)	9.4	16.2 (4.6) ^E	40	<3	
Female																
	9-13	165	9.2	(0.4)	6.1 (1.3) ^E	6.7 (1.1) ^E	7.9 (0.8)	9.3 (0.5)	11.1 (0.7)	13.2 (1.4)	14.7 (2.1)	7.0	F	23	<3	
	14-18	206	9.2	(0.5)	5.2 (0.7)	5.9 (0.6)	7.1 (0.5)	8.7 (0.5)	10.9 (0.8)	13.3 (1.2)	15.0 (1.6)	7.3	28.3 (8.0) ^E	34	<3	
	19-30	191	9.6	(0.7)	5.5 (1.0) ^E	6.2 (1.0)	7.4 (0.9)	9.1 (0.9)	11.5 (1.2)	14.3 (1.9)	16.6 (2.8) ^E	6.8	F	40	<3	
	31-50	258	9.5	(0.6)	7.0 (1.2) ^E	7.5 (1.1)	8.5 (1.0)	9.7 (0.9)	11.0 (1.2)	12.4 (1.7)	13.2 (2.2)	6.8	F	40	<3	
	51-70	249	9.6	(0.5)	6.2 (0.9)	6.8 (0.8)	7.9 (0.7)	9.3 (0.6)	11.0 (0.8)	12.8 (1.3)	14.0 (1.7)	6.8	F	40	0.0 (0.0)	
	>70	128	9.1	(0.5)	5.7 (0.7)	6.4 (0.7)	7.7 (0.7)	9.4 (0.7)	11.0 (0.8)	12.6 (1.1)	13.8 (1.4)	6.8	F	40	0.0 (0.0)	
	19+	826	9.5	(0.3)	6.1 (0.5)	6.7 (0.5)	7.8 (0.5)	9.3 (0.5)	11.2 (0.6)	13.4 (0.9)	14.9 (1.1)	6.8	F	40	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.10 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (<i>and SE</i>) of usual intake						EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)	
					5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)		95th (<i>SE</i>)					
Both																	
	1-3	192	7.6	(0.4)	5.3 (0.6)	5.8 (0.5)	6.6 (0.5)	7.6 (0.5)	8.8 (0.6)	10.1 (0.9)	11.0 (1.1)	2.5	<3	7	64.7	(12.6) ^E	
	4-8	321	9.1	(0.3)	5.9 (0.4)	6.5 (0.4)	7.6 (0.4)	9.1 (0.3)	10.8 (0.5)	12.6 (0.7)	13.8 (0.9)	4.0	<3	12	13.6	(4.2) ^E	
Male																	
	9-13	226	11.8	(0.5)	6.8 (0.6)	7.6 (0.6)	9.2 (0.6)	11.5 (0.6)	14.2 (0.8)	17.3 (1.2)	19.6 (1.6)	7.0	F	23	F		
	14-18	262	15.1	(1.0)	8.6 (1.1)	9.7 (1.1)	11.8 (1.0)	14.6 (1.0)	18.0 (1.3)	21.7 (2.1)	24.2 (2.7)	8.5	F	34	<3		
	19-30	197	14.7	(0.9)	7.9 (0.9)	8.8 (0.9)	10.7 (0.9)	13.7 (1.0)	17.9 (1.5)	22.5 (2.6)	26.0 (3.7)	9.4	F	40	<3		
	31-50	282	15.6	(0.8)	8.6 (1.1)	9.8 (1.0)	12.1 (1.0)	15.2 (1.0)	19.0 (1.2)	23.1 (1.9)	26.2 (2.7)	9.4	F	40	<3		
	51-70	234	13.0	(0.6)	7.9 (1.1)	8.7 (1.0)	10.2 (0.9)	12.2 (0.8)	14.7 (1.0)	17.2 (1.5)	19.0 (1.9)	9.4	F	40	0.0	(0.0)	
	>70	119	10.1	(0.6)	5.7 (0.9)	6.4 (0.8)	7.8 (0.7)	9.6 (0.7)	11.8 (0.9)	14.1 (1.3)	15.6 (1.6)	9.4	47.4	(11.4) ^E	40	0.0	(0.0)
	19+	832	14.1	(0.4)	7.8 (0.4)	8.8 (0.4)	10.7 (0.4)	13.5 (0.5)	17.0 (0.7)	20.6 (1.0)	23.1 (1.3)	9.4	14.3	(3.0) ^E	40	<3	
Female																	
	9-13	226	11.4	(1.9)	6.8 (1.1)	7.3 (1.1)	8.5 (1.2)	10.3 (1.5)	13.3 (2.4) ^E	17.6 (4.4) ^E	21.4 (6.6) ^E	7.0	F	23	F		
	14-18	242	9.2	(0.4)	5.2 (0.6)	5.9 (0.6)	7.2 (0.5)	8.8 (0.5)	10.7 (0.6)	12.7 (0.9)	14.0 (1.1)	7.3	25.9	(7.8) ^E	34	0.0	(0.0)
	19-30	208	10.0	(0.7)	6.1 (0.9)	6.8 (0.9)	8.1 (0.8)	9.8 (0.8)	12.0 (1.1)	14.3 (1.7)	16.0 (2.2)	6.8	F	40	<3		
	31-50	263	10.9	(0.6)	7.2 (1.0)	7.9 (0.8)	9.0 (0.7)	10.5 (0.7)	12.1 (1.0)	13.8 (1.4)	14.9 (1.7)	6.8	F	40	0.0	(0.0)	
	51-70	322	9.1	(0.6)	6.7 (1.0)	7.2 (0.9)	8.0 (0.8)	9.1 (0.6)	10.3 (0.8)	11.6 (1.2)	12.4 (1.6)	6.8	F	40	0.0	(0.0)	
	>70	198	8.9	(0.5)	5.0 (0.7)	5.7 (0.6)	7.1 (0.6)	8.8 (0.7)	10.7 (0.8)	12.5 (1.0)	13.7 (1.2)	6.8	F	40	0.0	(0.0)	
	19+	991	10.0	(0.3)	6.2 (0.5)	6.8 (0.4)	8.1 (0.4)	9.7 (0.3)	11.6 (0.5)	13.7 (0.7)	15.0 (0.9)	6.8	F	40	0.0	(0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.11 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR	(SE)	UL ³	% >UL	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)						
Both																
	1-3	348	7.5 (0.3)	5.5 (0.5)	5.9 (0.4)	6.6 (0.4)	7.5 (0.3)	8.5 (0.4)	9.6 (0.6)	10.2 (0.7)	2.5	0.0 (0.0)	7	64.5 (10.2)		
	4-8	554	9.3 (0.3)	6.4 (0.6)	7.0 (0.5)	8.0 (0.4)	9.3 (0.4)	10.8 (0.4)	12.3 (0.6)	13.3 (0.8)	4.0	<3	12	F		
Male	9-13	409	11.9 (0.5)	8.1 (0.6)	8.8 (0.6)	9.9 (0.6)	11.5 (0.6)	13.3 (0.7)	15.2 (1.0)	16.5 (1.2)	7.0	F	23	<3		
	14-18	414	13.5 (0.6)	8.0 (0.9)	9.2 (0.9)	11.1 (0.8)	13.6 (0.8)	16.6 (1.0)	19.4 (1.4)	21.3 (1.8)	8.5	F	34	<3		
	19-30	311	15.0 (0.7)	8.5 (0.9)	9.7 (0.9)	12.0 (0.8)	15.0 (0.9)	18.3 (1.3)	21.8 (1.8)	24.1 (2.2)	9.4	F	40	<3		
	31-50	489	13.4 (0.6)	7.8 (0.7)	8.7 (0.7)	10.4 (0.7)	12.7 (0.8)	15.7 (1.1)	19.0 (1.6)	21.3 (2.1)	9.4	F	40	<3		
	51-70	575	12.0 (0.5)	7.4 (0.9)	8.2 (0.8)	9.6 (0.6)	11.4 (0.6)	13.5 (0.7)	15.7 (1.1)	17.1 (1.4)	9.4	F	40	0.0 (0.0)		
	>70	239	10.9 (0.6)	7.8 (1.0)	8.4 (0.9)	9.4 (0.8)	10.8 (0.8)	12.3 (0.9)	13.8 (1.2)	14.8 (1.5)	9.4	F	40	0.0 (0.0)		
	19+	1614	13.1 (0.3)	7.6 (0.4)	8.5 (0.4)	10.3 (0.4)	12.6 (0.4)	15.4 (0.5)	18.5 (0.8)	20.6 (1.0)	9.4	16.8 (3.2) ^E	40	<3		
Female	9-13	355	9.2 (0.5)	5.9 (0.6)	6.5 (0.6)	7.6 (0.6)	9.1 (0.5)	10.8 (0.6)	12.6 (0.9)	13.8 (1.1)	7.0	F	23	<3		
	14-18	410	8.7 (0.4)	5.5 (0.6)	6.1 (0.6)	7.2 (0.5)	8.5 (0.4)	10.0 (0.6)	11.7 (0.9)	12.8 (1.1)	7.3	26.6 (8.3) ^E	34	0.0 (0.0)		
	19-30	384	9.6 (0.7)	6.3 (0.6)	6.9 (0.6)	8.0 (0.5)	9.3 (0.5)	11.5 (0.9)	14.3 (2.0)	16.8 (3.4) ^E	6.8	F	40	<3		
	31-50	585	9.9 (0.4)	5.3 (0.5)	6.1 (0.5)	7.6 (0.5)	9.8 (0.5)	12.1 (0.6)	14.7 (0.9)	16.4 (1.1)	6.8	16.2 (4.4) ^E	40	0.0 (0.0)		
	51-70	711	9.1 (0.4)	5.9 (0.5)	6.5 (0.5)	7.5 (0.5)	9.0 (0.5)	10.6 (0.6)	12.3 (0.9)	13.5 (1.2)	6.8	F	40	0.0 (0.0)		
	>70	342	8.5 (0.4)	4.8 (0.5)	5.4 (0.5)	6.6 (0.5)	8.2 (0.6)	10.1 (0.7)	12.0 (0.8)	13.3 (1.0)	6.8	27.8 (8.3) ^E	40	0.0 (0.0)		
	19+	2022	9.5 (0.2)	5.6 (0.3)	6.3 (0.3)	7.6 (0.3)	9.2 (0.3)	11.4 (0.4)	13.7 (0.7)	15.4 (0.9)	6.8	15.2 (3.1) ^E	40	<3		

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.12 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)	UL ³	% >UL (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)				
Both														
	1-3	622	7.1 (0.2)	4.4 (0.2)	4.9 (0.2)	5.7 (0.2)	7.0 (0.3)	8.4 (0.3)	9.9 (0.4)	10.9 (0.6)	2.5	<3	7	49.2 (5.4)
	4-8	919	8.8 (0.3)	6.6 (0.7)	7.0 (0.6)	7.7 (0.5)	8.7 (0.3)	9.7 (0.4)	10.8 (0.7)	11.5 (1.0)	4.0	<3	12	F
Male														
	9-13	579	13.0 (0.9)	10.4 (1.3)	11.0 (1.2)	12.0 (1.0)	13.3 (1.0)	14.7 (1.3)	16.2 (2.2)	17.2 (2.8)	7.0	<3	23	F
	14-18	634	14.9 (0.5)	8.2 (0.8)	9.4 (0.8)	11.7 (0.7)	14.5 (0.7)	17.9 (0.9)	21.7 (1.2)	24.3 (1.6)	8.5	F	34	<3
	19-30	578	14.5 (0.5)	9.7 (1.2)	10.6 (1.1)	12.2 (0.8)	14.1 (0.6)	16.3 (0.9)	18.5 (1.4)	20.0 (1.8)	9.4	F	40	<3
	31-50	693	13.6 (0.7)	6.9 (0.6)	8.0 (0.6)	10.0 (0.6)	12.6 (0.6)	16.1 (0.8)	20.1 (1.2)	22.8 (1.5)	9.4	20.2 (4.8) ^E	40	<3
	51-70	596	12.1 (0.4)	6.7 (0.7)	7.7 (0.7)	9.6 (0.6)	11.9 (0.5)	14.6 (0.6)	17.4 (1.0)	19.4 (1.3)	9.4	23.3 (5.7) ^E	40	<3
	>70	296	11.2 (0.5)	6.4 (0.6)	7.1 (0.6)	8.6 (0.6)	10.8 (0.6)	13.5 (0.7)	16.9 (1.2)	19.7 (1.9)	9.4	34.1 (7.8) ^E	40	<3
	19+	2163	13.2 (0.3)	7.2 (0.4)	8.2 (0.3)	10.1 (0.3)	12.6 (0.3)	15.8 (0.5)	19.5 (0.7)	22.0 (1.0)	9.4	18.5 (2.7)	40	<3
Female														
	9-13	533	9.4 (0.3)	6.3 (0.7)	6.9 (0.6)	8.0 (0.5)	9.4 (0.4)	11.0 (0.5)	12.8 (0.8)	14.0 (1.1)	7.0	F	23	<3
	14-18	638	9.3 (0.3)	5.7 (0.5)	6.4 (0.4)	7.4 (0.4)	9.0 (0.4)	10.8 (0.5)	12.7 (0.7)	14.0 (0.9)	7.3	22.7 (5.9) ^E	34	0.0 (0.0)
	19-30	499	9.7 (0.5)	5.7 (0.6)	6.4 (0.6)	7.5 (0.5)	9.0 (0.5)	11.1 (0.7)	13.5 (1.1)	15.3 (1.6)	6.8	F	40	<3
	31-50	716	10.0 (0.4)	6.1 (0.9)	6.8 (0.9)	8.0 (0.8)	9.7 (0.7)	11.8 (0.7)	14.0 (1.0)	15.5 (1.4)	6.8	F	40	0.0 (0.0)
	51-70	745	9.6 (0.3)	7.2 (0.8)	7.7 (0.6)	8.5 (0.5)	9.6 (0.4)	10.7 (0.6)	11.9 (0.9)	12.7 (1.2)	6.8	F	40	0.0 (0.0)
	>70	510	9.1 (0.5)	4.9 (0.3)	5.6 (0.3)	6.9 (0.4)	8.6 (0.4)	10.9 (0.6)	13.6 (1.1)	15.8 (1.8)	6.8	23.8 (4.6) ^E	40	<3
	19+	2470	9.7 (0.2)	5.7 (0.3)	6.4 (0.3)	7.7 (0.3)	9.3 (0.3)	11.4 (0.4)	13.8 (0.5)	15.6 (0.7)	6.8	13.8 (3.0) ^E	40	0.0 (0.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 28.13 Zinc (mg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (<i>and SE</i>) of usual intake							EAR ²	% <EAR (SE)	UL ³	% >UL (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)					
Both															
	1-3	2117	7.4 (0.1)	4.8 (0.2)	5.3 (0.2)	6.2 (0.2)	7.4 (0.2)	8.8 (0.2)	10.1 (0.2)	11.0 (0.3)	2.5	<3	7	58.9 (3.3)	
	4-8	3235	9.3 (0.2)	6.4 (0.2)	6.9 (0.2)	7.8 (0.2)	9.1 (0.2)	10.7 (0.3)	12.3 (0.4)	13.4 (0.5)	4.0	<3	12	12.3 (2.5) ^E	
Male															
	9-13	2080	12.2 (0.3)	8.1 (0.4)	8.9 (0.3)	10.2 (0.3)	12.0 (0.3)	14.2 (0.4)	16.6 (0.7)	18.2 (0.9)	7.0	<3	23	<3	
	14-18	2288	14.8 (0.3)	8.3 (0.4)	9.5 (0.3)	11.7 (0.3)	14.6 (0.4)	18.2 (0.5)	22.3 (0.8)	25.2 (1.0)	8.5	5.6 (1.3) ^E	34	<3	
	19-30	1804	14.2 (0.3)	9.1 (0.6)	10.0 (0.5)	11.7 (0.4)	13.9 (0.4)	16.5 (0.6)	19.2 (0.9)	21.0 (1.1)	9.4	^F	40	0.0 (0.0)	
	31-50	2596	13.9 (0.3)	7.9 (0.3)	8.9 (0.3)	10.9 (0.3)	13.5 (0.3)	16.7 (0.4)	20.3 (0.6)	22.9 (0.9)	9.4	13.3 (2.3) ^E	40	<3	
	51-70	2550	12.2 (0.2)	6.9 (0.3)	7.8 (0.3)	9.4 (0.3)	11.6 (0.3)	14.2 (0.3)	17.0 (0.5)	19.0 (0.7)	9.4	24.6 (3.0)	40	0.0 (0.0)	
	>70	1520	10.3 (0.2)	6.4 (0.3)	7.0 (0.3)	8.3 (0.3)	10.0 (0.3)	12.2 (0.4)	14.5 (0.5)	16.1 (0.7)	9.4	41.0 (4.9)	40	0.0 (0.0)	
	19+	8470	13.2 (0.2)	7.5 (0.2)	8.5 (0.2)	10.3 (0.2)	12.8 (0.2)	15.7 (0.2)	18.9 (0.4)	21.2 (0.5)	9.4	16.8 (1.4)	40	0.0 (0.0)	
Female															
	9-13	1980	9.8 (0.3)	5.9 (0.2)	6.6 (0.2)	7.7 (0.2)	9.2 (0.2)	11.1 (0.3)	13.4 (0.6)	15.1 (1.0)	7.0	14.6 (2.5) ^E	23	<3	
	14-18	2256	9.5 (0.2)	5.7 (0.2)	6.4 (0.2)	7.7 (0.2)	9.3 (0.2)	11.4 (0.3)	13.6 (0.4)	15.0 (0.5)	7.3	19.6 (2.6)	34	0.0 (0.0)	
	19-30	1854	9.5 (0.2)	5.8 (0.3)	6.4 (0.3)	7.5 (0.2)	9.0 (0.3)	11.0 (0.3)	13.1 (0.5)	14.6 (0.7)	6.8	14.7 (3.3) ^E	40	0.0 (0.0)	
	31-50	2686	9.9 (0.2)	5.7 (0.2)	6.4 (0.2)	7.7 (0.2)	9.5 (0.2)	11.8 (0.3)	14.2 (0.5)	15.9 (0.6)	6.8	14.2 (2.1)	40	0.0 (0.0)	
	51-70	3200	9.7 (0.2)	6.7 (0.4)	7.3 (0.4)	8.4 (0.3)	9.7 (0.2)	11.2 (0.3)	12.8 (0.4)	13.8 (0.5)	6.8	^F	40	0.0 (0.0)	
	>70	2610	8.5 (0.2)	5.0 (0.2)	5.6 (0.2)	6.8 (0.2)	8.3 (0.2)	10.2 (0.2)	12.3 (0.3)	13.7 (0.4)	6.8	25.2 (3.0)	40	0.0 (0.0)	
	19+	10350	9.6 (0.1)	5.7 (0.1)	6.4 (0.1)	7.6 (0.1)	9.3 (0.1)	11.4 (0.2)	13.6 (0.2)	15.2 (0.3)	6.8	14.0 (1.3)	40	0.0 (0.0)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

³ UL is the Tolerable Upper Intake Level. For additional detail, see footnote 11 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Appendix A: Table Footnotes

The following footnotes apply to all of the summary data tables presented in Section II of this report.

1. The survey excludes from its target population those living in the three territories, individuals living on Indian reserves or on Crown lands, residents of institutions, full-time members of the Canadian Forces and residents of certain remote regions.
2. The tables exclude pregnant and breastfeeding females, subject to another set of nutritional recommendations. The sample of pregnant and breastfeeding females is not large enough to allow for reliable estimates.
3. Sample size and mean intake are based on the first 24-hour dietary recall (first day of interview) only.
4. Intakes are based on food consumption only. Intakes from vitamin and mineral supplements are not included. Inferences about the prevalence of nutrient excess or inadequacy based on intakes from food alone may respectively underestimate or overestimate the prevalences based on total nutrient intakes from both food and supplements.
5. The intake distribution (percentiles and percentage above or below a cut-off when applicable) was adjusted to remove within-individual variability using Software for Intake Distribution Estimation (SIDE) (Iowa State University, 1996) and the method presented in Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440-1449.
6. In some cases, within-individual variance was estimated at the regional or national level and applied at the provincial level. For more details, see Section II.4: Measuring Sampling Variability with Bootstrap Replication in Volume 1 of the *Nutrient Intakes from Food* report series.
7. Bootstrapping techniques were used to produce the coefficient of variation (CV) and the standard error (SE).
8. AMDR is the Acceptable Macronutrient Distribution Range, expressed as a percentage of total energy intake. Intakes inside the range (shown in the AMDR columns) are associated with a reduced risk of chronic disease while providing adequate intakes of essential nutrients. For further information on AMDR see the Health Canada publication *Canadian Community Health Survey*,

Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data, Section 2.1.5, p. 27.

The applications of the AMDRs for essential fatty acids to group assessment are not the same as for the other macronutrients. The lower boundaries for the AMDR for linoleic and alpha-linolenic acids are not based on the same type of endpoints as the boundaries for total fat and carbohydrate. The boundaries for fat and carbohydrate are set based on evidence indicating increased risk for coronary heart diseases and the lower bound of the AMDR for both n-6 (linoleic) and n-3 (alpha-linolenic) fatty acids is based on the percent of energy from these fatty acids needed to provide the AI for these nutrients. The AI, in turn, is based on the median intake of both linoleic and alpha-linolenic acid in the United States, where essential fatty acid deficiency is non-existent in the healthy population.

Thus, by definition about half the population has intakes of these fatty acids below the AI and therefore outside the AMDR. In other words, based on the AI, one would conclude that the population is "adequate" with respect to linoleic and alpha-linolenic acids, while based on the AMDR a different conclusion (i.e. that 50% of the population has intakes below the AMDR) would be reached. Therefore, the lower bound of the AMDRs for linoleic and alpha-linolenic acids should not be used in the assessment of population intakes.

9. EAR is the Estimated Average Requirement. The level of intake at the EAR (shown in the EAR columns) is the average daily intake level that is estimated to meet the requirement, as defined by the specified indicator of adequacy, in half of the apparently healthy individuals in a DRI age–sex group. For further information on EAR see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.1, p. 23.
10. AI is the Adequate Intake. The level of intake at the AI (shown in the AI columns) is the recommended average daily intake level based on observed or experimentally determined approximations or estimates of nutrient intake by a group or groups of apparently healthy people that are assumed to be adequate. It is developed when an EAR cannot be determined. The percentage of the population having a usual intake above the AI (shown in the %>AI columns) almost certainly meets their needs. The adequacy of intakes below the AI cannot be assessed, and should not be interpreted as being inadequate. For further information on AI see the Health Canada

publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.3, p. 25.

11. UL is the Tolerable Upper Intake Level. The level of intake at the UL (shown in the UL columns) is the highest average daily intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population. For further information on UL see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.4, p. 26.
12. For a more detailed understanding of DRIs and their interpretation when assessing intakes of particular nutrients, consult the summary of the series of publications on DRIs published by the Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*.
13. In terms of precision, the estimate 0.0 with a standard error of 0.0 refers to a standard error smaller than 0.1%.

Appendix B: Iron Estimation

The distribution of iron requirements for menstruating females and some of the other age–sex groups is not normal or necessarily symmetric. Therefore, the full probability approach is required for the estimation of iron inadequacy instead of the EAR cut-point method. For all age–sex groups, the iron requirement distributions from Appendix I of the Institute of Medicine’s (IOM) report on the DRIs for iron (IOM, 2001) were used to estimate inadequacy. For the three DRI age–sex groups of menstruating females aged between 14 and 50 years, the iron requirement distributions of mixed populations, which assumes 17% oral contraceptive (OC) users and 83% non-OC users, were used to estimate inadequacy (IOM, 2001).

Tables of the risk of inadequate intake for specified ranges of the usual intake of iron, which are provided in the IOM report, were used for calculating iron inadequacy. The following summarizes how the full probability method was used to estimate iron inadequacy:

- SIDE was used to estimate the usual intake distribution of iron. A file containing the intake value at 9,999 evenly spaced percentiles was generated for each domain.
- From Appendix I of the IOM report on the DRIs for iron, Table I-3 and Table I-4 were used. For females aged 14 to 18 years and menstruating women, the tables for the mixed adolescent and adult populations were used.
- For example, for the mixed adolescent population, intakes below 4.49 mg/d are assumed to have 100% probability of inadequacy (risk=1.0). Those with intakes above or equal to 14.39 mg/d are assumed to have zero risk of inadequacy. For intakes between these two extremes, the risk of inadequacy is calculated as 100 minus the midpoint of the percentile of requirement.
- Each of the 9,999 intake values fell into one of the specified requirement ranges, each with a corresponding risk value. The corresponding risk values are 1, 0.9625, 0.925, 0.85, 0.75, 0.65, 0.55, 0.45, 0.35, 0.25, 0.15, 0.075, 0.0375 and 0. The average of these 9,999 risk values was the estimate of the iron inadequacy for that age–sex group.
- Standard errors for the estimates were calculated with the probability approach using the bootstrap method.

- For additional information on iron estimation and the probability method, consult Appendix 3 of the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, or the section ‘Assessing the Adequacy of Intakes of Groups’ in Chapter 14 of the IOM’s DRI report on iron (IOM, 2001).

Appendix C: Justification for Excluding Nutrients from Volume 2 and Volume 3

Volume 1 of the compendium contained data on 13 nutrients, including 6 nutrients expressed as a percent of total energy. There were originally 31 other nutrients scheduled to be released in future volumes of the compendium, but for a variety of reasons some of these nutrients will not be included. Decisions to omit these nutrients were made jointly by representatives from Statistics Canada and Health Canada.

Exclusions and changes to the list of nutrients that were to be included in Volumes 2 and 3 of the compendium are as follows:

Total milligrams of folic acid

Folic acid is found in small amounts in a number of foods. Most respondents consumed a small amount of folic acid, which resulted in a bimodal distribution of folic acid intake. As a result, it was very difficult to normalize the distribution, which meant that SIDE was unable to calculate usual intake.

One of the steps that SIDE uses to estimate usual intake is to transform the data into a normal distribution. Assessing SIDE's ability to perform this transformation rests on measuring the Anderson-Darling (A-D) score for normality. The A-D score is a statistic that measures how close a distribution is to a normal distribution. Any A-D score less than 0.576 is considered to be sufficiently normal for SIDE to continue without warning. Typically, SIDE will be able to transform 95% of the domains without error using the default SIDE options. The remaining 5% of domains will typically score higher than 0.576 but usually less than 1.0. Adjusting the SIDE options will usually reduce the A-D to within the limit. In the case of folic acid, more than half of the provincial domains had an A-D score above the 0.576 threshold and many domains scored higher than 2. The nature of the data simply does not allow SIDE to produce proper estimates for the usual intake of folic acid.

Total grams of alcohol

Alcohol is consumed differently than other nutrients. For most respondents, alcohol is not part of their daily intake of food, but rather is something that is consumed occasionally. In this sense, in terms of analysis, alcohol behaves more like a food than a nutrient. In order for SIDE to estimate the usual intake of foods, many recalls are needed to capture enough occurrences of the particular food. Thus, two recalls are not enough to calculate the usual intake of alcohol.

Percent of energy from alcohol

The difficulty in estimating a usual intake for alcohol causes similar problems for expressing that intake as a percent of total energy.

Caffeine

Caffeine also is consumed differently than other nutrients. The usual intake of caffeine could not be calculated due to the same issues as folic acid and alcohol. Many respondents reported zero or small levels of caffeine intake. Therefore, it is difficult for SIDE to properly model the data with only two dietary recalls.

Based on the changes above, the list of nutrients included in Volume 1 and the revised list of nutrients included in Volumes 2 and 3 are as follows:

List of Nutrients Included in the Three-Volume Set		
Volume 1	Volume 2	Volume 3
Total Energy	Folate (DFE)	Folacin
Percentage of total energy intake from fats	Iron	Linolenic acid (g, % energy)
Percentage of total energy intake from protein	Linoleic acid (g, % energy)	Moisture
Percentage of total energy intake from carbohydrates	Magnesium	Naturally occurring folate
Percentage of total energy intake from saturated fats	Niacin	Protein
Percentage of total energy intake from monounsaturated fats	Phosphorus	Total carbohydrates
Percentage of total energy intake from polyunsaturated fats	Potassium	Total fats
Total dietary fibre	Riboflavin	Total monounsaturated fats
Cholesterol	Thiamin	Total polyunsaturated fats
Vitamin A	Vitamin B ₆	Total saturated fats
Vitamin C	Vitamin B ₁₂	Total sugars
Calcium	Vitamin C by smoking status	
Sodium	Vitamin D	
	Zinc	

Appendix D: References

Department of Statistics and Center for Agricultural and Rural Development, Iowa State University: *A User's Guide to SIDE, Software for Intake Distribution Estimation Version 1.0*. Technical Report 96-TR 30. Ames, IA: Iowa State University Statistical Laboratory, 1996. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr30.pdf

Dodd KW: *A Technical Guide to C-SIDE, Software for Intake Distribution Estimation*. Technical Report 96-TR 32, Dietary Assessment Research Series Report 9. Ames, IA: Department of Statistics and Center for Agricultural and Rural Development, Iowa State University, 2006. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr32.pdf

Health Canada: *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) — A Guide to Accessing and Interpreting the Data*. Ottawa: Publications, Health Canada, 2006. Available at: www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_focus-volet_escc-eng.php

Institute of Medicine: *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D and Fluoride*. Washington, DC: National Academy Press, 1997. Available at: <http://books.nap.edu/catalog/5776.html>

Institute of Medicine: *Dietary Reference Intakes: A Risk Assessment Model for Establishing Upper Intake Levels for Nutrients*. Washington, DC: National Academy Press, 1998. Available at: <http://books.nap.edu/catalog/6432.html>

Institute of Medicine: *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin and Choline*. Washington, DC: National Academy Press, 1998. Available at: www.nap.edu/catalog.php?record_id=6015

Institute of Medicine: *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids*. Washington, DC: National Academy Press, 2000. Available at: <http://books.nap.edu/catalog/9810.html>

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academy Press, 2000. Available at: <http://books.nap.edu/catalog/9956.html>

Institute of Medicine: *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc*. Washington, DC: National Academy Press, 2001. Available at: www.nap.edu/catalog.php?record_id=10026

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Planning*. Washington, DC: National Academy Press, 2003. Available at: <http://books.nap.edu/catalog/10609.html>

Institute of Medicine: *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington, DC: National Academy Press, 2004. Available at: <http://books.nap.edu/catalog/10925.html>

Institute of Medicine: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. Washington, DC: National Academy Press, 2005. Available at: <http://books.nap.edu/catalog/10490.html>

Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: National Academy Press, 2006. Available at: www.nap.edu/topics.php?topic=380

Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440–1449

Nusser SM, Fuller WA, Guenther PM: Estimating usual dietary intake distributions: adjusting for measurement error and nonnormality in 24-hour food intake data. In L Lyberg, P Biemer, M Collins et al., eds. *Survey Measurement and Process Quality*, New York: John Wiley & Sons, 1997.

Statistics Canada: *Canadian Community Health Survey (CCHS) Cycle 2.2—Nutrition: Software for Intake Distribution Estimation (SIDE) Documentation*. Ottawa, 2007. Available at: www.statcan.ca/english/sdds/document/5049_D22_T9_V1_E.pdf