



Health
Canada

Santé
Canada

Statistics
Canada

Statistique
Canada

Canadian Community Health Survey Cycle 2.2, Nutrition (2004)

Nutrient Intakes from Food

Provincial, Regional and National Summary Data Tables
Volume 3

Revised February 2009



Canada 

Health Canada is the federal department responsible for helping the people of Canada maintain and improve their health.

We assess the safety of drugs and many consumer products, help improve the safety of food, and provide information to Canadians to help them make healthy decisions. We provide health services to First Nations people and to Inuit communities. We work with the provinces to ensure our health care system serves the needs of Canadians.

Published by authority of the Minister of Health.

**Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) —
Nutrient Intakes from Food**

Provincial, Regional and National Summary Data Tables, Volume 3

is available on Internet at the following address:

www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/index-eng.php

Également disponible en français sous le titre :

Enquête sur la santé dans les collectivités canadiennes, cycle 2.2, Nutrition (2004) —

Apports nutritionnels provenant des aliments

Tableaux sommaires provinciaux, régionaux et nationaux, Volume 3

This publication can be made available on request on
diskette, large print, audio-cassette and braille.

For further information or to obtain additional copies, please contact:

Publications

Health Canada

Ottawa, Ontario K1A 0K9

Tel.: (613) 954-5995

Fax: (613) 941-5366

E-Mail: info@hc-sc.gc.ca

© Her Majesty the Queen in Right of Canada, represented by
the Minister of Health Canada, 2008

This publication may be reproduced without permission
provided the source is fully acknowledged.

Cat.: H164-45/3-2008E-PDF

ISBN: 978-0-662-48762-3

Acknowledgements

Health Canada would like to acknowledge and thank the individuals who have contributed to this report. This publication was produced as a joint venture between Health Canada and Statistics Canada. Technical experts from Health Canada's Food Directorate and Office of Nutrition Policy and Promotion and from Statistics Canada's Health Statistics Division developed the report. Project guidance was provided by Health Canada's Food and Nutrition Surveillance System Working Group, which consists of individuals from Health Canada, the Public Health Agency of Canada and the Federal/Provincial/Territorial Group on Nutrition. External experts have advised both departments, through the CCHS 2.2 Users Group, on the need for a consistent approach to analyzing and reporting these data. The draft report was reviewed by internal experts from both departments.

Table of Contents

	Acknowledgements	i
	List of Tables.....	iii
	List of Appendices	vii
	List of Abbreviations.....	viii
I	Introduction.....	1
II	Summary Data Tables	3
	<i>(table numbering continued from Volume 2)</i>	
	29. Folic acid ($\mu\text{g/d}$): Usual intakes from food	3
	30. Linolenic acid (g/d): Usual intakes from food	17
	31. Percentage of total energy intake from linolenic acid	31
	32. Moisture (g/d): Usual intakes from food	45
	33. Naturally occurring folate ($\mu\text{g/d}$): Usual intakes from food.....	59
	34. Protein (g/d): Usual intakes from food.....	73
	35. Total carbohydrates (g/d): Usual intakes from food	87
	36. Total fats (g/d): Usual intakes from food	101
	37. Total monounsaturated fats (g/d): Usual intakes from food.....	115
	38. Total polyunsaturated fats (g/d): Usual intakes from food	129
	39. Total saturated fats (g/d): Usual intakes from food	143
	40. Total sugars (g/d): Usual intakes from food.....	157

List of Tables

29. Folic acid ($\mu\text{g/d}$): Usual intakes from food, by DRI age–sex group, household population	3
Table 29.1 Newfoundland and Labrador, 2004	4
Table 29.2 Prince Edward Island, 2004	5
Table 29.3 Nova Scotia, 2004	6
Table 29.4 New Brunswick, 2004	7
Table 29.5 Quebec, 2004	8
Table 29.6 Ontario, 2004	9
Table 29.7 Manitoba, 2004	10
Table 29.8 Saskatchewan, 2004	11
Table 29.9 Alberta, 2004	12
Table 29.10 British Columbia, 2004	13
Table 29.11 Atlantic Region, 2004	14
Table 29.12 Prairie Region, 2004	15
Table 29.13 Canada excluding territories, 2004	16
30. Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population	17
Table 30.1 Newfoundland and Labrador, 2004	18
Table 30.2 Prince Edward Island, 2004	19
Table 30.3 Nova Scotia, 2004	20
Table 30.4 New Brunswick, 2004	21
Table 30.5 Quebec, 2004	22
Table 30.6 Ontario, 2004	23
Table 30.7 Manitoba, 2004	24
Table 30.8 Saskatchewan, 2004	25
Table 30.9 Alberta, 2004	26
Table 30.10 British Columbia, 2004	27
Table 30.11 Atlantic Region, 2004	28
Table 30.12 Prairie Region, 2004	29
Table 30.13 Canada excluding territories, 2004	30
31. Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population	31
Table 31.1 Newfoundland and Labrador, 2004	32
Table 31.2 Prince Edward Island, 2004	33
Table 31.3 Nova Scotia, 2004	34
Table 31.4 New Brunswick, 2004	35
Table 31.5 Quebec, 2004	36
Table 31.6 Ontario, 2004	37
Table 31.7 Manitoba, 2004	38
Table 31.8 Saskatchewan, 2004	39
Table 31.9 Alberta, 2004	40
Table 31.10 British Columbia, 2004	41
Table 31.11 Atlantic Region, 2004	42
Table 31.12 Prairie Region, 2004	43
Table 31.13 Canada excluding territories, 2004	44

32. Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population.....	45
Table 32.1 Newfoundland and Labrador, 2004.....	46
Table 32.2 Prince Edward Island, 2004.....	47
Table 32.3 Nova Scotia, 2004.....	48
Table 32.4 New Brunswick, 2004	49
Table 32.5 Quebec, 2004.....	50
Table 32.6 Ontario, 2004	51
Table 32.7 Manitoba, 2004	52
Table 32.8 Saskatchewan, 2004	53
Table 32.9 Alberta, 2004.....	54
Table 32.10 British Columbia, 2004.....	55
Table 32.11 Atlantic Region, 2004	56
Table 32.12 Prairie Region, 2004.....	57
Table 32.13 Canada excluding territories, 2004.....	58
33. Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population.....	59
Table 33.1 Newfoundland and Labrador, 2004.....	60
Table 33.2 Prince Edward Island, 2004.....	61
Table 33.3 Nova Scotia, 2004.....	62
Table 33.4 New Brunswick, 2004	63
Table 33.5 Quebec, 2004.....	64
Table 33.6 Ontario, 2004	65
Table 33.7 Manitoba, 2004	66
Table 33.8 Saskatchewan, 2004	67
Table 33.9 Alberta, 2004.....	68
Table 33.10 British Columbia, 2004.....	69
Table 33.11 Atlantic Region, 2004	70
Table 33.12 Prairie Region, 2004.....	71
Table 33.13 Canada excluding territories, 2004.....	72
34. Protein (g/d): Usual intakes from food, by DRI age–sex group, household population.....	73
Table 34.1 Newfoundland and Labrador, 2004.....	74
Table 34.2 Prince Edward Island, 2004.....	75
Table 34.3 Nova Scotia, 2004.....	76
Table 34.4 New Brunswick, 2004	77
Table 34.5 Quebec, 2004.....	78
Table 34.6 Ontario, 2004	79
Table 34.7 Manitoba, 2004	80
Table 34.8 Saskatchewan, 2004	81
Table 34.9 Alberta, 2004.....	82
Table 34.10 British Columbia, 2004.....	83
Table 34.11 Atlantic Region, 2004	84
Table 34.12 Prairie Region, 2004.....	85
Table 34.13 Canada excluding territories, 2004.....	86

35. Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population	87
Table 35.1 Newfoundland and Labrador, 2004	88
Table 35.2 Prince Edward Island, 2004	89
Table 35.3 Nova Scotia, 2004	90
Table 35.4 New Brunswick, 2004	91
Table 35.5 Quebec, 2004	92
Table 35.6 Ontario, 2004	93
Table 35.7 Manitoba, 2004	94
Table 35.8 Saskatchewan, 2004	95
Table 35.9 Alberta, 2004	96
Table 35.10 British Columbia, 2004	97
Table 35.11 Atlantic Region, 2004	98
Table 35.12 Prairie Region, 2004	99
Table 35.13 Canada excluding territories, 2004	100
36. Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population	101
Table 36.1 Newfoundland and Labrador, 2004	102
Table 36.2 Prince Edward Island, 2004	103
Table 36.3 Nova Scotia, 2004	104
Table 36.4 New Brunswick, 2004	105
Table 36.5 Quebec, 2004	106
Table 36.6 Ontario, 2004	107
Table 36.7 Manitoba, 2004	108
Table 36.8 Saskatchewan, 2004	109
Table 36.9 Alberta, 2004	110
Table 36.10 British Columbia, 2004	111
Table 36.11 Atlantic Region, 2004	112
Table 36.12 Prairie Region, 2004	113
Table 36.13 Canada excluding territories, 2004	114
37. Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population	115
Table 37.1 Newfoundland and Labrador, 2004	116
Table 37.2 Prince Edward Island, 2004	117
Table 37.3 Nova Scotia, 2004	118
Table 37.4 New Brunswick, 2004	119
Table 37.5 Quebec, 2004	120
Table 37.6 Ontario, 2004	121
Table 37.7 Manitoba, 2004	122
Table 37.8 Saskatchewan, 2004	123
Table 37.9 Alberta, 2004	124
Table 37.10 British Columbia, 2004	125
Table 37.11 Atlantic Region, 2004	126
Table 37.12 Prairie Region, 2004	127
Table 37.13 Canada excluding territories, 2004	128

38. Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population.....	129
Table 38.1 Newfoundland and Labrador, 2004.....	130
Table 38.2 Prince Edward Island, 2004.....	131
Table 38.3 Nova Scotia, 2004.....	132
Table 38.4 New Brunswick, 2004	133
Table 38.5 Quebec, 2004.....	134
Table 38.6 Ontario, 2004	135
Table 38.7 Manitoba, 2004	136
Table 38.8 Saskatchewan, 2004	137
Table 38.9 Alberta, 2004.....	138
Table 38.10 British Columbia, 2004.....	139
Table 38.11 Atlantic Region, 2004	140
Table 38.12 Prairie Region, 2004.....	141
Table 38.13 Canada excluding territories, 2004.....	142
39. Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population.....	143
Table 39.1 Newfoundland and Labrador, 2004.....	144
Table 39.2 Prince Edward Island, 2004.....	145
Table 39.3 Nova Scotia, 2004.....	146
Table 39.4 New Brunswick, 2004	147
Table 39.5 Quebec, 2004.....	148
Table 39.6 Ontario, 2004	149
Table 39.7 Manitoba, 2004	150
Table 39.8 Saskatchewan, 2004	151
Table 39.9 Alberta, 2004.....	152
Table 39.10 British Columbia, 2004.....	153
Table 39.11 Atlantic Region, 2004	154
Table 39.12 Prairie Region, 2004.....	155
Table 39.13 Canada excluding territories, 2004.....	156
40. Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population.....	157
Table 40.1 Newfoundland and Labrador, 2004.....	158
Table 40.2 Prince Edward Island, 2004.....	159
Table 40.3 Nova Scotia, 2004.....	160
Table 40.4 New Brunswick, 2004	161
Table 40.5 Quebec, 2004.....	162
Table 40.6 Ontario, 2004	163
Table 40.7 Manitoba, 2004	164
Table 40.8 Saskatchewan, 2004	165
Table 40.9 Alberta, 2004.....	166
Table 40.10 British Columbia, 2004.....	167
Table 40.11 Atlantic Region, 2004	168
Table 40.12 Prairie Region, 2004.....	169
Table 40.13 Canada excluding territories, 2004.....	170

List of Appendices

Appendix A: Table Footnotes.....	171
Appendix B: Justification for Excluding Nutrients from Volume 2 and Volume 3.....	175
List of Nutrients Included in the Three-Volume Set.....	176
Appendix C: References.....	177

List of Abbreviations

Abbreviation	Meaning
AI	Adequate Intake
AMDR	Acceptable Macronutrient Distribution Range
CCHS	Canadian Community Health Survey
CV	coefficient of variation
d	day
DRI	Dietary Reference Intake
EAR	Estimated Average Requirement
g	gram
IOM	Institute of Medicine
kg	kilogram
µg	microgram (sometimes reported as mcg)
n	sample size
SE	standard error
SIDE	Software for Intake Distribution Estimation

I Introduction

This is Volume 3 of a three-volume set that provides summary data tables about the nutrient intakes from food obtained by Canadians in 2004. The data come from the Canadian Community Health Survey (CCHS), Cycle 2.2, Nutrition (2004). As in Volumes 1 and 2, data are provided for 14 Dietary Reference Intake (DRI) age–sex groups. For nutrients that have DRIs, the tables also compare usual intakes of these nutrients to the DRIs. Data used for producing the tables in this report were obtained from the CCHS 2.2 Share File. The nutrient intakes represent food consumption; data on nutrient intakes from vitamin and mineral supplements were not included in the production of these tables.¹

This series of products released by Health Canada is part of its ongoing support to users of the CCHS 2.2 data. It has been undertaken as a joint venture with Statistics Canada. The series is a reference for those who will use the CCHS Cycle 2.2 data and its findings to guide nutrition-related program and policy decisions. It will be of particular benefit to provincial ministries of health, researchers and graduate students, policy makers and analysts, public health professionals, epidemiologists, dietitians, the food industry, and the health media.

All three volumes consist primarily of data tables; they do not provide any interpretation or draw conclusions. To optimize the usage of this volume, we recommend that it be read in concert with Volumes 1 and 2, as well as with the report, *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) – A Guide to Accessing and Interpreting the Data* (available at www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_focus-volet_escc-eng.php), published by Health Canada in 2006. That report includes an overview of the CCHS 2.2, including a description of the survey sample, how the survey was conducted,

¹ Because supplements may make meaningful contributions to nutrient intakes, inferences about the prevalence of nutrient excess or inadequacy based on intakes from food alone may respectively underestimate or overestimate the prevalences based on total nutrient intakes from both food and supplements.

survey components and an introduction to DRIs, including examples of how to interpret the CCHS 2.2 data.

Volume 3 includes 12 sets of data tables, on 11 nutrients. Results are presented for 13 geographical areas: the 10 provinces, the Atlantic Region, the Prairie Region, and Canada excluding the territories. Data from the four Atlantic provinces and the three Prairie provinces were combined into the Atlantic Region and the Prairie Region, respectively, given the small sample sizes in these provinces.

A revised list of the data tables that appear in all three volumes is found in **Appendix B**.

Quality Assurance and Quality Control

Data quality processes were established to oversee the data analysis and production of the data tables. As the project developed, these processes also helped to monitor and address methodological issues that arose. The processes followed were jointly developed and agreed to by Health Canada and Statistics Canada. Refer to Volume 1 for a description of the methodology used to produce the tables, including the process used and the means of addressing problems encountered.

II Summary Data Tables

29. Folacin (µg/d): Usual intakes from food

Table 29.1 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	195 (14)	166 (34) ^E	169 (29) ^E	174 (20)	180 (16)	186 (27)	191 (43) ^E	195 (54) ^E
	4-8	127	295 (29)	201 (37) ^E	218 (36)	249 (34)	292 (33)	345 (41)	407 (57)	451 (74)
Male	9-13	111	346 (22)	252 (28)	267 (26)	293 (24)	329 (26)	375 (34)	426 (47)	460 (57)
	14-18	107	373 (26)	242 (37)	273 (34)	330 (30)	401 (32)	485 (44)	577 (65)	641 (83)
	19-30	77	354 (24)	250 (24)	276 (26)	320 (28)	372 (30)	425 (32)	476 (35)	507 (36)
	31-50	145	314 (19)	187 (28)	208 (26)	248 (23)	298 (23)	358 (29)	424 (39)	471 (49)
	51-70	182	320 (18)	167 (29) ^E	190 (28)	236 (24)	300 (20)	376 (26)	452 (38)	500 (46)
	>70	63	331 (26)	235 (33)	251 (32)	283 (33)	324 (36)	373 (43)	422 (55)	454 (64)
	19+	467	326 (11)	194 (16)	218 (15)	261 (14)	317 (13)	381 (16)	446 (20)	488 (25)
Female	9-13	96	260 (16)	169 (24)	185 (22)	215 (20)	252 (20)	293 (24)	335 (33)	361 (39)
	14-18	105	280 (21)	156 (23)	177 (24)	218 (27)	275 (29)	336 (32)	397 (37)	433 (42)
	19-30	91	249 (18)	169 (26)	183 (23)	208 (21)	237 (21)	269 (27)	301 (36)	321 (43)
	31-50	167	261 (17)	149 (16)	172 (16)	211 (18)	264 (23)	330 (30)	402 (39)	451 (45)
	51-70	198	293 (18)	148 (38) ^E	173 (33) ^E	222 (24)	281 (18)	353 (27)	449 (60)	532 (98) ^E
	>70	74	313 (41)	177 (19)	198 (22)	238 (28)	293 (37)	363 (51)	440 (69)	495 (82) ^E
	19+	530	274 (11)	167 (15)	187 (15)	226 (14)	276 (14)	335 (16)	400 (23)	446 (31)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.”

The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.2 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	210 (17)	114 (22) ^E	133 (21)	168 (20)	213 (22)	264 (27)	315 (34)	348 (41)
	4-8	110	257 (16)	158 (21)	179 (20)	215 (19)	258 (20)	307 (27)	355 (34)	383 (40)
Male	9-13	95	316 (22)	228 (28)	243 (29)	269 (30)	301 (32)	337 (34)	371 (37)	393 (39)
	14-18	87	387 (39)	262 (51) ^E	291 (48)	343 (45)	408 (48)	484 (74)	563 (101) ^E	616 (120) ^E
	19-30	70	405 (36)	194 (50) ^E	232 (48) ^E	306 (44)	394 (43)	496 (57)	614 (85)	698 (107)
	31-50	109	342 (21)	283 (40)	293 (36)	311 (29)	332 (29)	354 (40)	375 (57)	388 (69) ^E
	51-70	128	316 (14)	197 (23)	222 (20)	261 (17)	308 (17)	368 (22)	433 (33)	479 (43)
	>70	65	295 (14)	206 (24)	223 (22)	256 (20)	297 (19)	343 (25)	385 (35)	411 (42)
	19+	372	343 (12)	205 (15)	230 (14)	275 (14)	332 (15)	399 (19)	472 (26)	523 (33)
Female	9-13	75	266 (21)	183 (33) ^E	199 (30)	228 (28)	263 (28)	303 (33)	341 (41)	366 (47)
	14-18	81	265 (15)	132 (31) ^E	158 (27) ^E	202 (21)	249 (18)	300 (23)	357 (35)	397 (47)
	19-30	101	283 (17)	179 (23)	200 (22)	239 (21)	287 (22)	339 (27)	391 (35)	423 (41)
	31-50	116	292 (28)	150 (24)	177 (22)	225 (22)	286 (28)	361 (41)	447 (61)	509 (79)
	51-70	146	270 (13)	179 (25)	196 (22)	226 (19)	262 (17)	301 (20)	339 (26)	363 (32)
	>70	94	240 (17)	139 (19)	155 (18)	187 (18)	231 (19)	287 (25)	351 (35)	396 (44)
	19+	457	277 (13)	164 (12)	186 (12)	226 (12)	277 (15)	333 (19)	393 (27)	435 (34)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.3 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	219 (13)	148 (19)	162 (17)	189 (16)	223 (17)	263 (23)	301 (32)	324 (38)
	4-8	177	253 (13)	165 (26)	183 (23)	215 (19)	257 (17)	306 (24)	357 (37)	392 (49)
Male	9-13	111	337 (19)	222 (32)	242 (30)	280 (27)	331 (26)	391 (35)	455 (49)	499 (58)
	14-18	113	342 (25)	223 (40) ^E	245 (38)	288 (34)	342 (35)	398 (42)	451 (54)	485 (63)
	19-30	91	370 (27)	209 (44) ^E	234 (41) ^E	280 (35)	338 (31)	404 (33)	470 (43)	513 (54)
	31-50	101	381 (29)	221 (38) ^E	250 (35)	303 (31)	364 (29)	433 (35)	509 (50)	562 (64)
	51-70	134	351 (31)	236 (42) ^E	257 (39)	295 (35)	344 (35)	404 (45)	472 (64)	518 (80)
	>70	56	286 (19)	197 (35) ^E	213 (33)	242 (30)	277 (26)	315 (25)	352 (31)	375 (38)
	19+	382	360 (15)	204 (19)	230 (18)	278 (17)	341 (17)	417 (23)	501 (33)	559 (42)
Female	9-13	105	271 (16)	210 (26)	225 (23)	251 (19)	280 (19)	311 (23)	339 (30)	356 (35)
	14-18	120	272 (25)	137 (41) ^E	163 (37) ^E	209 (31)	267 (27)	334 (31)	405 (43)	454 (55)
	19-30	91	282 (19)	200 (14)	217 (16)	244 (18)	278 (23)	318 (28)	361 (34)	386 (37)
	31-50	159	288 (14)	156 (19)	181 (18)	227 (18)	284 (19)	348 (24)	406 (31)	442 (36)
	51-70	174	299 (19)	169 (26)	192 (24)	232 (21)	282 (20)	343 (28)	413 (46)	464 (59)
	>70	80	243 (14)	176 (24)	189 (22)	212 (19)	242 (20)	280 (26)	322 (38)	352 (49)
	19+	504	284 (9)	183 (15)	202 (14)	237 (12)	281 (12)	331 (15)	384 (22)	421 (27)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.4 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	201 (12)	159 (19)	168 (18)	184 (16)	203 (15)	224 (18)	244 (25)	257 (31)
	4-8	140	307 (23)	169 (30) ^E	192 (28)	237 (26)	299 (26)	377 (37)	467 (60)	532 (80)
Male	9-13	92	329 (28)	223 (28)	237 (28)	262 (29)	294 (31)	332 (34)	373 (38)	401 (41)
	14-18	107	413 (32)	264 (52) ^E	294 (48)	350 (42)	422 (38)	509 (45)	605 (67)	674 (89)
	19-30	73	475 (50)	239 (58) ^E	280 (55) ^E	361 (54)	468 (60)	593 (79)	723 (111)	815 (138) ^E
	31-50	134	365 (26)	225 (42) ^E	246 (39)	287 (33)	339 (30)	400 (37)	464 (58)	506 (77)
	51-70	131	352 (23)	231 (17)	250 (19)	290 (23)	343 (27)	397 (32)	458 (39)	505 (45)
	>70	55	323 (28)	190 (27)	216 (29)	264 (32)	318 (34)	372 (37)	426 (43)	464 (51)
	19+	393	380 (16)	220 (24)	247 (22)	297 (20)	366 (20)	450 (26)	542 (40)	606 (51)
Female	9-13	79	272 (20)	185 (21)	200 (22)	229 (24)	266 (27)	310 (29)	356 (32)	387 (35)
	14-18	104	284 (21)	186 (35) ^E	203 (32)	232 (27)	269 (25)	310 (30)	355 (43)	385 (55)
	19-30	101	291 (30)	144 (43) ^E	170 (43) ^E	220 (42) ^E	289 (41)	374 (43)	467 (51)	530 (61)
	31-50	143	303 (21)	181 (37) ^E	203 (33)	243 (27)	295 (24)	355 (33)	417 (51)	457 (66)
	51-70	193	294 (14)	160 (25)	184 (22)	226 (18)	277 (17)	336 (21)	403 (31)	451 (42)
	>70	94	256 (16)	153 (19)	170 (19)	199 (18)	235 (20)	278 (26)	325 (34)	355 (39)
	19+	531	292 (11)	155 (15)	180 (14)	225 (13)	284 (14)	353 (16)	425 (20)	474 (24)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.5 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	311	236 (11)	125 (16)	145 (15)	183 (14)	231 (15)	286 (17)	340 (22)	375 (27)
	4-8	485	318 (13)	244 (28)	260 (25)	289 (19)	324 (16)	361 (20)	399 (29)	424 (37)
Male	9-13	277	400 (19)	259 (30)	288 (28)	341 (26)	407 (26)	485 (32)	568 (43)	623 (53)
	14-18	339	435 (18)	259 (24)	292 (24)	354 (22)	436 (24)	537 (33)	654 (54)	739 (68)
	19-30	237	472 (25)	311 (37)	344 (34)	404 (29)	480 (27)	565 (35)	651 (51)	706 (63)
	31-50	423	438 (15)	288 (25)	318 (23)	372 (20)	438 (18)	513 (22)	592 (30)	645 (38)
	51-70	387	394 (16)	248 (22)	275 (20)	324 (18)	388 (19)	465 (30)	553 (51)	616 (73)
	>70	132	329 (23)	172 (41) ^E	200 (39) ^E	254 (35)	321 (34)	393 (39)	466 (51)	517 (64)
	19+	1179	423 (11)	259 (13)	291 (13)	348 (12)	420 (12)	505 (15)	596 (22)	660 (29)
	Female	9-13	281	320 (13)	200 (22)	221 (21)	259 (18)	307 (17)	361 (21)	415 (30)
14-18	321	369 (17)	225 (23)	254 (22)	306 (21)	371 (23)	441 (30)	510 (39)	553 (45)	
19-30	249	412 (34)	280 (45)	303 (42)	345 (37)	395 (36)	452 (42)	508 (55)	545 (66)	
31-50	364	362 (25)	269 (41)	289 (37)	325 (32)	369 (30)	417 (34)	467 (48)	500 (60)	
51-70	467	331 (10)	201 (15)	226 (14)	271 (12)	327 (12)	392 (16)	460 (23)	506 (29)	
>70	215	270 (13)	158 (18)	178 (18)	217 (17)	266 (18)	327 (24)	397 (36)	449 (49)	
19+	1295	351 (12)	212 (13)	237 (13)	284 (13)	345 (14)	417 (18)	496 (25)	552 (32)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.6 Folacin (µg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2,3}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	232 (8)	134 (18)	153 (15)	187 (12)	229 (9)	278 (12)	333 (22)	371 (31)
	4-8	956	296 (7)	189 (18)	209 (15)	245 (11)	290 (7)	344 (11)	400 (21)	439 (29)
Male	9-13	589	351 (9)	211 (17)	236 (15)	283 (12)	341 (10)	409 (13)	483 (22)	537 (30)
	14-18	639	422 (15)	283 (41)	311 (35)	359 (25)	420 (16)	488 (24)	557 (44)	602 (58)
	19-30	481	447 (22)	281 (44)	311 (38)	365 (28)	434 (23)	512 (37)	593 (65)	647 (87)
	31-50	709	401 (13)	271 (43)	298 (38)	343 (27)	398 (16)	462 (22)	525 (43)	563 (59)
	51-70	758	381 (11)	211 (24)	240 (21)	296 (17)	369 (13)	453 (18)	543 (33)	608 (49)
	>70	734	324 (10)	168 (12)	193 (11)	241 (10)	307 (11)	386 (13)	473 (20)	532 (25)
	19+	2682	399 (8)	223 (15)	254 (14)	312 (11)	387 (9)	475 (12)	565 (20)	626 (28)
Female	9-13	585	309 (9)	177 (20)	201 (17)	246 (12)	301 (9)	363 (14)	429 (25)	475 (34)
	14-18	645	337 (10)	203 (26)	227 (23)	273 (17)	331 (12)	397 (16)	462 (27)	505 (36)
	19-30	514	305 (10)	183 (28)	206 (24)	248 (17)	298 (12)	355 (17)	412 (30)	449 (39)
	31-50	758	330 (11)	169 (17)	193 (16)	241 (13)	307 (12)	391 (17)	488 (31)	556 (42)
	51-70	955	313 (9)	166 (12)	192 (11)	239 (9)	301 (9)	371 (13)	441 (18)	488 (22)
	>70	1345	280 (6)	167 (10)	186 (9)	220 (8)	268 (6)	324 (9)	381 (15)	419 (21)
	19+	3572	314 (6)	177 (8)	200 (7)	244 (6)	301 (6)	369 (9)	442 (13)	492 (17)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folacin.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folacin” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.7 Folacin (µg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2,3}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	209 (9)	123 (16)	138 (14)	164 (12)	198 (11)	241 (14)	290 (23)	325 (32)
	4-8	425	282 (17)	193 (28)	209 (24)	238 (20)	274 (19)	313 (27)	353 (39)	378 (48)
Male	9-13	274	332 (12)	226 (11)	246 (12)	282 (13)	327 (16)	379 (19)	433 (22)	468 (23)
	14-18	297	426 (18)	224 (22)	260 (20)	327 (16)	414 (19)	515 (30)	619 (44)	687 (53)
	19-30	249	452 (36)	263 (22)	294 (23)	351 (27)	430 (35)	532 (49)	646 (65)	723 (76)
	31-50	309	393 (19)	242 (39)	270 (35)	323 (28)	391 (24)	465 (33)	535 (48)	579 (60)
	51-70	277	345 (14)	204 (28)	230 (25)	277 (20)	336 (16)	404 (21)	472 (33)	516 (42)
	>70	136	291 (20)	156 (27) ^E	179 (25)	224 (21)	280 (19)	343 (32)	411 (44)	460 (56)
	19+	971	383 (12)	230 (25)	257 (22)	309 (17)	374 (14)	452 (20)	535 (34)	593 (46)
	Female	9-13	265	303 (18)	235 (16)	246 (17)	265 (17)	288 (18)	312 (19)	335 (21)
14-18	290	314 (14)	178 (31) ^E	201 (28)	246 (22)	302 (18)	367 (24)	433 (37)	476 (47)	
19-30	197	322 (23)	232 (36)	250 (30)	281 (22)	316 (18)	351 (25)	383 (37)	402 (44)	
31-50	312	321 (17)	200 (23)	220 (21)	258 (18)	307 (18)	366 (25)	429 (36)	470 (46)	
51-70	312	303 (15)	197 (30)	216 (27)	250 (22)	292 (18)	341 (21)	395 (34)	433 (46)	
>70	239	239 (9)	155 (23)	171 (20)	200 (15)	235 (11)	276 (15)	319 (26)	349 (35)	
19+	1060	304 (9)	182 (17)	204 (15)	243 (11)	293 (9)	352 (14)	415 (26)	459 (34)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folacin.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folacin” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.8 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	189 (9)	118 (13)	131 (12)	153 (11)	179 (11)	211 (14)	247 (21)	274 (28)
	4-8	213	275 (12)	174 (21)	193 (18)	226 (14)	269 (13)	317 (19)	365 (28)	397 (36)
Male	9-13	122	345 (35)	202 (32)	227 (31)	274 (31)	340 (36)	428 (52)	527 (79)	594 (99) ^E
	14-18	150	443 (44)	264 (51) ^E	295 (48)	353 (44)	430 (44)	521 (62)	618 (98)	683 (127) ^E
	19-30	106	435 (41)	202 (42) ^E	238 (41) ^E	308 (38)	405 (40)	525 (53)	655 (78)	744 (100)
	31-50	155	388 (24)	220 (45) ^E	248 (41)	301 (34)	379 (30)	473 (44)	568 (70)	630 (90)
	51-70	122	368 (46)	208 (33)	234 (31)	278 (30)	337 (41)	438 (85) ^E	616 (199) ^E	^F
	>70	88	316 (18)	192 (28)	213 (26)	252 (23)	304 (21)	368 (26)	438 (40)	487 (52)
	19+	471	385 (19)	187 (16)	219 (15)	281 (14)	361 (16)	467 (24)	610 (50)	723 (81)
Female	9-13	103	324 (25)	237 (35)	254 (33)	285 (32)	322 (32)	363 (38)	404 (48)	430 (55)
	14-18	142	296 (18)	175 (17)	196 (17)	235 (18)	285 (21)	348 (29)	423 (39)	480 (49)
	19-30	111	288 (19)	160 (28) ^E	183 (24)	224 (20)	278 (20)	345 (31)	419 (51)	472 (69)
	31-50	146	315 (21)	203 (39) ^E	222 (35)	257 (30)	302 (27)	359 (30)	424 (46)	471 (62)
	51-70	184	314 (13)	251 (37)	265 (32)	289 (23)	317 (16)	348 (24)	377 (41)	396 (54)
	>70	143	248 (13)	178 (23)	193 (21)	218 (18)	251 (16)	289 (19)	328 (26)	354 (32)
	19+	584	299 (10)	184 (15)	204 (15)	243 (13)	293 (13)	352 (15)	416 (20)	461 (25)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.9 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2,3}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	169	203	(10)	100	(19) ^E	117	(18)	148	(15)	190	(12)	241	(15)	300	(26)	346	(38)
	4-8	281	265	(10)	182	(25)	197	(22)	224	(16)	257	(12)	296	(17)	334	(30)	360	(39)
Male	9-13	183	364	(22)	198	(27)	224	(26)	275	(23)	347	(23)	442	(35)	556	(62)	637	(85)
	14-18	187	422	(23)	315	(51)	337	(45)	377	(35)	425	(28)	477	(35)	527	(55)	559	(72)
	19-30	223	410	(20)	382	(90) ^E	391	(76) ^E	406	(49)	422	(24)	438	(37)	453	(75)	461	(102) ^E
	31-50	229	385	(22)	228	(50) ^E	257	(46) ^E	316	(38)	399	(33)	497	(44)	595	(66)	658	(84)
	51-70	197	348	(15)	186	(33) ^E	219	(29)	277	(21)	345	(18)	421	(24)	501	(41)	558	(56)
	>70	72	314	(26)	182	(26)	201	(26)	240	(26)	296	(28)	368	(41)	464	(66)	544	(91) ^E
	19+	721	378	(11)	225	(21)	254	(20)	310	(17)	383	(15)	468	(18)	557	(27)	617	(36)
	Female	9-13	165	314	(16)	190	(37) ^E	216	(33)	264	(26)	328	(21)	405	(33)	482	(52)	530
14-18	206	316	(18)	179	(32) ^E	205	(29)	252	(25)	309	(22)	370	(25)	430	(33)	469	(41)	
19-30	191	328	(18)	130	(23) ^E	156	(24)	216	(25)	308	(24)	410	(28)	535	(42)	632	(58)	
31-50	258	298	(14)	226	(40) ^E	243	(35)	271	(27)	305	(19)	342	(23)	380	(39)	406	(52)	
51-70	249	291	(15)	182	(29)	203	(26)	240	(21)	287	(18)	338	(23)	393	(35)	430	(46)	
>70	128	260	(18)	133	(21)	153	(20)	192	(19)	246	(20)	313	(27)	386	(39)	437	(51)	
19+	826	299	(7)	159	(11)	183	(11)	232	(10)	296	(10)	365	(12)	441	(18)	498	(23)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.10 Folacin (µg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	192	256 (13)	167 (26)	184 (23)	216 (19)	257 (17)	303 (22)	352 (32)	384 (41)
	4-8	321	313 (13)	191 (18)	212 (16)	249 (14)	298 (14)	359 (20)	427 (34)	475 (47)
Male	9-13	226	366 (16)	206 (17)	233 (17)	283 (17)	350 (18)	433 (25)	526 (37)	593 (48)
	14-18	262	473 (30)	292 (56) ^E	327 (50)	389 (41)	468 (35)	559 (45)	651 (72)	712 (96)
	19-30	197	492 (29)	293 (51) ^E	327 (46)	391 (37)	475 (33)	573 (44)	674 (67)	740 (85)
	31-50	282	484 (28)	324 (48)	354 (42)	408 (33)	476 (29)	550 (40)	620 (58)	664 (71)
	51-70	234	407 (19)	232 (23)	259 (22)	312 (21)	385 (22)	477 (30)	581 (44)	654 (57)
	>70	119	358 (21)	198 (24)	227 (23)	281 (24)	349 (25)	431 (32)	521 (45)	585 (58)
	19+	832	451 (15)	253 (13)	285 (13)	349 (13)	437 (16)	540 (23)	647 (32)	720 (39)
Female	9-13	226	338 (21)	200 (22)	224 (22)	271 (22)	333 (24)	409 (30)	491 (40)	547 (48)
	14-18	242	345 (20)	155 (23)	182 (23)	239 (21)	318 (23)	416 (32)	532 (51)	614 (67)
	19-30	208	344 (22)	217 (31)	241 (29)	287 (26)	345 (26)	413 (34)	487 (49)	538 (64)
	31-50	263	385 (19)	184 (20)	214 (20)	272 (20)	352 (23)	456 (29)	567 (40)	633 (48)
	51-70	322	340 (21)	186 (25)	215 (23)	269 (22)	340 (23)	422 (31)	507 (45)	563 (57)
	>70	198	284 (19)	159 (19)	181 (20)	220 (21)	273 (24)	334 (28)	392 (32)	428 (36)
	19+	991	352 (10)	184 (11)	212 (10)	265 (10)	337 (11)	426 (16)	524 (24)	585 (29)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folacin.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folacin” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.11 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2,3}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	348	207 (7)	132 (9)	146 (9)	171 (9)	203 (9)	241 (12)	279 (16)	303 (19)
	4-8	554	280 (11)	179 (16)	198 (15)	234 (14)	282 (14)	340 (17)	405 (24)	451 (32)
Male	9-13	409	335 (13)	243 (23)	259 (21)	287 (17)	322 (15)	362 (19)	404 (30)	433 (38)
	14-18	414	374 (15)	232 (22)	261 (21)	315 (20)	384 (20)	467 (25)	557 (35)	620 (44)
	19-30	311	403 (19)	236 (20)	264 (20)	319 (20)	389 (22)	470 (27)	555 (34)	611 (41)
	31-50	489	358 (14)	210 (16)	235 (15)	280 (15)	339 (15)	407 (20)	480 (28)	529 (36)
	51-70	575	342 (16)	214 (23)	236 (21)	276 (19)	329 (18)	396 (22)	470 (34)	521 (44)
	>70	239	308 (13)	192 (17)	214 (17)	254 (17)	304 (17)	357 (18)	411 (22)	449 (27)
	19+	1614	358 (8)	205 (9)	231 (9)	278 (9)	341 (10)	418 (12)	502 (17)	560 (21)
Female	9-13	355	269 (10)	185 (15)	201 (14)	232 (13)	269 (13)	309 (15)	350 (20)	375 (23)
	14-18	410	277 (13)	141 (21)	167 (19)	212 (16)	266 (15)	330 (17)	402 (23)	447 (28)
	19-30	384	278 (13)	181 (17)	201 (16)	235 (15)	279 (16)	330 (19)	382 (24)	417 (30)
	31-50	585	287 (9)	152 (11)	176 (11)	222 (12)	284 (13)	355 (16)	431 (20)	483 (24)
	51-70	711	295 (10)	162 (11)	185 (11)	227 (10)	281 (11)	346 (14)	420 (21)	474 (28)
	>70	342	260 (11)	154 (11)	172 (11)	207 (11)	250 (13)	304 (18)	368 (25)	414 (31)
	19+	2022	284 (6)	163 (7)	185 (6)	227 (7)	280 (7)	343 (9)	412 (12)	460 (15)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.12 Folacin (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2,3}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	622	202	(7)	110	(9)	125	(8)	155	(8)	193	(8)	238	(10)	289	(15)	325	(21)
	4-8	919	270	(7)	172	(14)	189	(12)	221	(9)	262	(8)	309	(13)	356	(20)	386	(26)
Male																		
	9-13	579	353	(15)	220	(25)	244	(23)	290	(19)	352	(17)	423	(24)	501	(39)	556	(51)
	14-18	634	427	(16)	249	(22)	281	(21)	341	(19)	418	(19)	509	(25)	606	(37)	673	(48)
	19-30	578	423	(17)	266	(43)	297	(37)	354	(27)	423	(19)	495	(25)	571	(43)	624	(60)
	31-50	693	387	(15)	232	(29)	261	(27)	318	(23)	393	(22)	481	(27)	572	(40)	631	(51)
	51-70	596	351	(13)	191	(20)	222	(17)	276	(14)	343	(13)	422	(18)	521	(41)	600	(71)
	>70	296	309	(15)	182	(17)	202	(17)	240	(18)	293	(16)	361	(70) ^E	443	(81) ^E	503	(92) ^E
	19+	2163	380	(8)	211	(12)	243	(11)	299	(10)	376	(10)	471	(13)	571	(19)	645	(27)
Female																		
	9-13	533	313	(11)	256	(29)	270	(26)	294	(20)	322	(16)	353	(18)	382	(26)	401	(33)
	14-18	638	312	(12)	165	(15)	190	(14)	239	(14)	303	(15)	375	(17)	447	(21)	495	(24)
	19-30	499	320	(13)	162	(18)	188	(17)	241	(15)	309	(14)	385	(18)	464	(26)	520	(33)
	31-50	716	306	(10)	196	(26)	217	(24)	256	(19)	305	(14)	360	(15)	418	(24)	458	(32)
	51-70	745	298	(9)	189	(17)	209	(15)	247	(13)	295	(12)	349	(15)	405	(23)	443	(30)
	>70	510	252	(10)	145	(10)	165	(9)	196	(10)	243	(10)	297	(14)	370	(22)	419	(29)
	19+	2470	300	(5)	169	(7)	192	(7)	237	(7)	295	(6)	360	(8)	432	(11)	483	(14)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folacin. ³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folacin” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 29.13 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	228 (4)	130 (6)	148 (6)	181 (5)	224 (5)	273 (7)	325 (9)	361 (12)
	4-8	3235	297 (5)	184 (7)	204 (6)	242 (5)	291 (5)	349 (8)	411 (12)	454 (16)
Male	9-13	2080	363 (7)	230 (9)	254 (9)	300 (8)	358 (8)	425 (9)	496 (13)	546 (17)
	14-18	2288	429 (9)	246 (10)	280 (10)	344 (9)	425 (10)	522 (13)	627 (19)	701 (24)
	19-30	1804	451 (12)	259 (13)	294 (13)	358 (12)	440 (13)	536 (17)	639 (26)	710 (33)
	31-50	2596	415 (8)	243 (12)	275 (11)	334 (10)	410 (9)	496 (11)	583 (16)	642 (21)
	51-70	2550	380 (7)	215 (8)	244 (7)	298 (7)	366 (7)	448 (10)	543 (18)	618 (27)
	>70	1520	326 (8)	185 (8)	209 (8)	255 (9)	315 (10)	386 (13)	463 (18)	515 (23)
	19+	8470	406 (5)	226 (5)	258 (5)	317 (5)	395 (5)	487 (7)	587 (11)	656 (14)
Female	9-13	1980	313 (6)	194 (8)	216 (8)	256 (7)	307 (7)	365 (9)	425 (12)	464 (15)
	14-18	2256	336 (6)	177 (7)	204 (7)	257 (7)	327 (8)	407 (10)	491 (14)	549 (17)
	19-30	1854	336 (10)	190 (10)	216 (9)	263 (9)	323 (10)	392 (13)	466 (17)	517 (21)
	31-50	2686	337 (8)	180 (7)	205 (7)	255 (7)	321 (8)	402 (11)	495 (16)	561 (21)
	51-70	3200	317 (5)	182 (6)	207 (6)	252 (6)	310 (6)	378 (8)	448 (11)	494 (14)
	>70	2610	272 (5)	157 (6)	177 (6)	215 (5)	264 (6)	326 (8)	393 (13)	441 (17)
	19+	10350	324 (4)	177 (3)	203 (3)	249 (4)	312 (5)	387 (6)	469 (8)	526 (11)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

30. Linolenic acid (g/d): Usual intakes from food

Table 30.1 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	79	0.86 (0.09)	0.53 (0.11) ^E	0.58 (0.10) ^E	0.68 (0.09)	0.80 (0.11)	0.95 (0.15)	1.10 (0.24) ^E	1.20 (0.33) ^E	0.7	70.5	(18.2) ^E
	4-8	127	1.51 (0.12)	0.92 (0.16) ^E	1.03 (0.16)	1.24 (0.17)	1.53 (0.18)	1.89 (0.22)	2.28 (0.30)	2.56 (0.38)	0.9	95.8	(7.6)
Male	9-13	111	1.92 (0.18)	1.25 (0.25) ^E	1.37 (0.24) ^E	1.60 (0.22)	1.91 (0.23)	2.27 (0.30)	2.66 (0.43)	2.92 (0.54) ^E	1.2	96.4	(7.2)
	14-18	107	1.91 (0.15)	1.01 (0.26) ^E	1.16 (0.24) ^E	1.43 (0.22)	1.86 (0.20)	2.55 (0.31)	3.30 (0.54)	3.72 (0.67) ^E	1.6	64.5	(13.4) ^E
	19-30	77	1.93 (0.18)	1.15 (0.18)	1.29 (0.19)	1.56 (0.22)	1.90 (0.25)	2.29 (0.29)	2.66 (0.34)	2.90 (0.37)	1.6	72.1	(16.0) ^E
	31-50	145	2.13 (0.21)	1.36 (0.27) ^E	1.49 (0.25) ^E	1.72 (0.22)	2.02 (0.21)	2.37 (0.25)	2.72 (0.34)	2.95 (0.43)	1.6	83.6	(13.7)
	51-70	182	1.77 (0.20)	0.78 (0.25) ^E	0.91 (0.24) ^E	1.16 (0.22) ^E	1.53 (0.20)	2.02 (0.25)	2.57 (0.43)	2.98 (0.60) ^E	1.6	45.9	(15.0) ^E
	>70	63	1.82 (0.17)	0.94 (0.22) ^E	1.14 (0.22) ^E	1.50 (0.21)	1.96 (0.23)	2.47 (0.27)	2.97 (0.34)	3.30 (0.40)	1.6	69.7	(12.1) ^E
	19+	467	1.96 (0.11)	0.95 (0.12)	1.11 (0.11)	1.41 (0.11)	1.83 (0.11)	2.34 (0.14)	2.88 (0.21)	3.25 (0.27)	1.6	63.9	(6.7)
Female	9-13	96	1.30 (0.10)	0.92 (0.16) ^E	0.99 (0.15)	1.11 (0.13)	1.27 (0.12)	1.45 (0.15)	1.62 (0.21)	1.74 (0.25)	1.0	88.8	(12.8)
	14-18	105	1.48 (0.20)	0.64 (0.20) ^E	0.72 (0.20) ^E	0.90 (0.19) ^E	1.18 (0.19)	1.56 (0.23)	2.04 (0.36) ^E	2.40 (0.50) ^E	1.1	56.5	(18.5) ^E
	19-30	91	1.53 (0.22)	0.99 (0.22) ^E	1.08 (0.20) ^E	1.23 (0.19)	1.43 (0.19)	1.65 (0.24)	1.87 (0.34) ^E	2.02 (0.42) ^E	1.1	88.1	(17.2) ^E
	31-50	167	1.55 (0.13)	0.73 (0.16) ^E	0.84 (0.16) ^E	1.08 (0.16)	1.46 (0.17)	1.97 (0.22)	2.56 (0.39)	3.00 (0.58) ^E	1.1	74.0	(10.7)
	51-70	198	1.40 (0.13)	0.45 (0.12) ^E	0.58 (0.12) ^E	0.84 (0.13)	1.26 (0.15)	1.93 (0.21)	2.77 (0.36)	3.35 (0.48)	1.1	59.0	(8.3)
	>70	74	1.51 (0.20)	0.86 (0.17) ^E	0.98 (0.18) ^E	1.20 (0.18)	1.47 (0.21)	1.82 (0.28)	2.23 (0.40) ^E	2.52 (0.49) ^E	1.1	83.0	(14.8) ^E
	19+	530	1.50 (0.07)	0.68 (0.07)	0.80 (0.07)	1.05 (0.07)	1.43 (0.08)	1.92 (0.12)	2.48 (0.19)	2.88 (0.27)	1.1	71.4	(5.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.2 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	58	0.74 (0.06)	0.49 (0.10) ^E	0.52 (0.10) ^E	0.59 (0.09)	0.68 (0.09)	0.77 (0.09)	0.86 (0.11)	0.92 (0.14)	0.7	F	
	4-8	110	1.16 (0.07)	0.84 (0.08)	0.90 (0.09)	1.02 (0.09)	1.16 (0.10)	1.33 (0.11)	1.48 (0.12)	1.58 (0.12)	0.9	90.4	(9.8)
Male	9-13	95	1.40 (0.12)	1.08 (0.19) ^E	1.15 (0.18)	1.29 (0.16)	1.45 (0.15)	1.64 (0.16)	1.84 (0.20)	1.96 (0.23)	1.2	85.5	(18.4) ^E
	14-18	87	2.02 (0.23)	1.43 (0.35) ^E	1.54 (0.33) ^E	1.74 (0.30) ^E	2.00 (0.30)	2.29 (0.39) ^E	2.59 (0.58) ^E	2.79 (0.76) ^E	1.6	86.3	(23.0) ^E
	19-30	70	2.64 (0.23)	1.87 (0.45) ^E	2.07 (0.43) ^E	2.43 (0.39)	2.86 (0.34)	3.31 (0.35)	3.74 (0.48)	4.01 (0.62)	1.6	98.6	(8.7)
	31-50	109	2.24 (0.17)	1.52 (0.34) ^E	1.70 (0.32) ^E	2.02 (0.28)	2.45 (0.26)	2.94 (0.33)	3.45 (0.50)	3.79 (0.65) ^E	1.6	93.0	(8.6)
	51-70	128	1.91 (0.20)	0.97 (0.27) ^E	1.16 (0.26) ^E	1.51 (0.24)	1.96 (0.25)	2.52 (0.33)	3.17 (0.49)	3.64 (0.63) ^E	1.6	70.0	(13.4) ^E
	>70	65	1.55 (0.13)	0.78 (0.17) ^E	0.90 (0.17) ^E	1.13 (0.15)	1.46 (0.16)	1.88 (0.21)	2.39 (0.35)	2.76 (0.48) ^E	1.6	40.1	(12.2) ^E
	19+	372	2.16 (0.11)	1.15 (0.16)	1.34 (0.16)	1.72 (0.15)	2.23 (0.15)	2.86 (0.18)	3.53 (0.26)	4.00 (0.32)	1.6	80.5	(6.6)
Female	9-13	75	1.77 (0.33) ^E	0.89 (0.12)	1.02 (0.14)	1.26 (0.17)	1.62 (0.24)	2.08 (0.36) ^E	2.64 (0.56) ^E	3.06 (0.76) ^E	1.0	90.8	(8.5)
	14-18	81	1.44 (0.14)	0.87 (0.22) ^E	0.96 (0.21) ^E	1.13 (0.19) ^E	1.34 (0.19)	1.61 (0.21)	1.89 (0.27)	2.08 (0.32)	1.1	77.8	(15.4) ^E
	19-30	101	1.54 (0.12)	1.07 (0.22) ^E	1.19 (0.21) ^E	1.41 (0.18)	1.71 (0.19)	2.04 (0.26)	2.36 (0.43) ^E	2.57 (0.57) ^E	1.1	94.0	(7.1)
	31-50	116	1.50 (0.16)	0.80 (0.19) ^E	0.91 (0.18) ^E	1.14 (0.17)	1.45 (0.18)	1.83 (0.23)	2.25 (0.34)	2.54 (0.43) ^E	1.1	77.9	(13.0) ^E
	51-70	146	1.56 (0.13)	0.96 (0.22) ^E	1.06 (0.20) ^E	1.25 (0.17)	1.51 (0.15)	1.80 (0.18)	2.12 (0.28)	2.33 (0.37)	1.1	87.7	(12.3)
	>70	94	1.48 (0.26) ^E	0.75 (0.21) ^E	0.85 (0.21) ^E	1.04 (0.21) ^E	1.32 (0.24) ^E	1.68 (0.31) ^E	2.09 (0.45) ^E	2.39 (0.58) ^E	1.1	70.0	(20.7) ^E
	19+	457	1.52 (0.09)	0.83 (0.11)	0.94 (0.11)	1.17 (0.10)	1.49 (0.10)	1.90 (0.14)	2.33 (0.21)	2.63 (0.26)	1.1	80.1	(6.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.3 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	112	0.95 (0.12)	0.58 (0.13) ^E	0.63 (0.12) ^E	0.73 (0.12)	0.87 (0.12)	1.06 (0.15)	1.28 (0.23) ^E	1.45 (0.32) ^E	0.7	79.4	(14.0) ^E
	4-8	177	1.23 (0.11)	0.60 (0.15) ^E	0.70 (0.14) ^E	0.90 (0.12)	1.19 (0.12)	1.58 (0.18)	2.06 (0.36) ^E	2.45 (0.55) ^E	0.9	75.1	(10.3)
Male	9-13	111	1.59 (0.12)	1.05 (0.21) ^E	1.14 (0.20) ^E	1.31 (0.16)	1.52 (0.14)	1.76 (0.18)	2.00 (0.28)	2.16 (0.36)	1.2	85.6	(13.4)
	14-18	113	1.97 (0.22)	1.05 (0.32) ^E	1.20 (0.31) ^E	1.51 (0.29) ^E	1.91 (0.29)	2.35 (0.34)	2.79 (0.44)	3.09 (0.54) ^E	1.6	69.6	(16.3) ^E
	19-30	91	2.69 (0.64) ^E	^F	1.58 (0.52) ^E	2.01 (0.59) ^E	2.66 (0.76) ^E	^F	^F	^F	1.6	89.3	(17.3) ^E
	31-50	101	1.95 (0.18)	0.71 (0.20) ^E	0.87 (0.20) ^E	1.19 (0.20) ^E	1.65 (0.22)	2.24 (0.28)	2.91 (0.38)	3.37 (0.47)	1.6	52.6	(12.3) ^E
	51-70	134	2.10 (0.30)	^F	1.14 (0.32) ^E	1.50 (0.31) ^E	2.05 (0.35) ^E	2.74 (0.51) ^E	3.53 (0.76) ^E	4.11 (1.01) ^E	1.6	70.5	(17.6) ^E
	>70	56	1.41 (0.13)	0.73 (0.15) ^E	0.82 (0.15) ^E	1.01 (0.14)	1.26 (0.14)	1.57 (0.19)	1.91 (0.29)	2.15 (0.38) ^E	1.6	^F	
	19+	382	2.09 (0.17)	0.80 (0.12)	0.97 (0.11)	1.28 (0.11)	1.73 (0.14)	2.53 (0.27)	3.72 (0.63) ^E	4.73 (1.05) ^E	1.6	56.5	(6.8)
Female	9-13	105	1.30 (0.10)	0.86 (0.08)	0.94 (0.09)	1.09 (0.10)	1.28 (0.12)	1.50 (0.14)	1.73 (0.16)	1.88 (0.17)	1.0	84.9	(10.4)
	14-18	120	1.72 (0.44) ^E	0.74 (0.16) ^E	0.87 (0.18) ^E	1.12 (0.23) ^E	1.49 (0.34) ^E	2.02 (0.55) ^E	2.69 (0.87) ^E	^F	1.1	76.4	(14.4) ^E
	19-30	91	1.85 (0.22)	^F	1.01 (0.31) ^E	1.33 (0.29) ^E	1.75 (0.28)	2.25 (0.33)	2.80 (0.47) ^E	3.17 (0.60) ^E	1.1	86.5	(10.5)
	31-50	159	1.97 (0.17)	0.86 (0.20) ^E	1.04 (0.21) ^E	1.39 (0.21)	1.87 (0.24)	2.49 (0.29)	3.16 (0.39)	3.63 (0.49)	1.1	87.7	(8.2)
	51-70	174	1.59 (0.12)	0.93 (0.21) ^E	1.03 (0.19) ^E	1.23 (0.16)	1.50 (0.14)	1.83 (0.20)	2.18 (0.33)	2.42 (0.44) ^E	1.1	85.3	(11.6)
	>70	80	1.34 (0.15)	0.78 (0.12)	0.88 (0.13)	1.09 (0.16)	1.38 (0.21)	1.71 (0.25)	2.03 (0.31)	2.28 (0.37)	1.1	74.3	(16.2) ^E
	19+	504	1.76 (0.08)	0.82 (0.11)	0.95 (0.10)	1.23 (0.11)	1.65 (0.12)	2.26 (0.16)	2.94 (0.25)	3.38 (0.31)	1.1	82.5	(5.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.4 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	99	0.88 (0.11)	0.44 (0.12) ^E	0.51 (0.12) ^E	0.64 (0.11) ^E	0.83 (0.12)	1.05 (0.14)	1.28 (0.18)	1.44 (0.22)	0.7	67.5	(16.3) ^E
	4-8	140	1.31 (0.09)	0.92 (0.06)	0.99 (0.07)	1.11 (0.08)	1.26 (0.09)	1.44 (0.11)	1.62 (0.13)	1.75 (0.15)	0.9	96.0	(3.5)
Male	9-13	92	F	F	F	F	2.47 (0.82) ^E	F	F	F	1.2	95.8	(12.4)
	14-18	107	2.13 (0.21)	1.33 (0.30) ^E	1.47 (0.29) ^E	1.74 (0.28)	2.11 (0.28)	2.51 (0.31)	2.93 (0.40)	3.24 (0.50)	1.6	83.3	(16.1) ^E
	19-30	73	2.23 (0.22)	1.25 (0.32) ^E	1.44 (0.29) ^E	1.79 (0.25)	2.26 (0.26)	2.80 (0.36)	3.37 (0.51)	3.74 (0.62) ^E	1.6	83.9	(10.8)
	31-50	134	1.99 (0.18)	0.88 (0.26) ^E	1.04 (0.24) ^E	1.36 (0.22)	1.83 (0.21)	2.44 (0.28)	3.12 (0.43)	3.61 (0.57)	1.6	62.3	(13.3) ^E
	51-70	131	1.94 (0.18)	1.04 (0.11)	1.16 (0.12)	1.40 (0.15)	1.73 (0.20)	2.17 (0.24)	2.66 (0.29)	2.99 (0.33)	1.6	59.8	(12.1) ^E
	>70	55	2.18 (0.28)	1.25 (0.36) ^E	1.42 (0.36) ^E	1.77 (0.35) ^E	2.26 (0.37)	2.89 (0.45)	3.62 (0.68) ^E	4.14 (0.92) ^E	1.6	83.0	(16.1) ^E
	19+	393	2.05 (0.10)	1.07 (0.15)	1.24 (0.15)	1.57 (0.13)	2.02 (0.13)	2.57 (0.16)	3.15 (0.24)	3.55 (0.32)	1.6	73.3	(7.5)
Female	9-13	79	1.35 (0.17)	0.76 (0.17) ^E	0.86 (0.18) ^E	1.06 (0.19) ^E	1.31 (0.22)	1.63 (0.25)	1.99 (0.30)	2.25 (0.34)	1.0	80.4	(15.2) ^E
	14-18	104	1.43 (0.14)	0.88 (0.11)	0.97 (0.12)	1.15 (0.14)	1.41 (0.18)	1.73 (0.22)	2.08 (0.28)	2.33 (0.33)	1.1	79.9	(12.0)
	19-30	101	1.39 (0.13)	0.96 (0.24) ^E	1.05 (0.22) ^E	1.24 (0.20)	1.46 (0.18)	1.71 (0.21)	1.96 (0.28)	2.12 (0.34)	1.1	87.0	(14.2)
	31-50	143	1.34 (0.13)	0.60 (0.08)	0.69 (0.08)	0.89 (0.10)	1.19 (0.12)	1.60 (0.17)	2.11 (0.28)	2.52 (0.41)	1.1	57.1	(9.6) ^E
	51-70	193	1.57 (0.11)	0.82 (0.09)	0.93 (0.09)	1.14 (0.11)	1.42 (0.12)	1.77 (0.15)	2.14 (0.18)	2.41 (0.20)	1.1	78.2	(8.0)
	>70	94	1.19 (0.11)	0.44 (0.11) ^E	0.52 (0.12) ^E	0.71 (0.13) ^E	1.03 (0.13)	1.43 (0.17)	1.89 (0.24)	2.23 (0.32)	1.1	44.8	(11.2) ^E
	19+	531	1.40 (0.06)	0.76 (0.10)	0.87 (0.10)	1.06 (0.09)	1.33 (0.08)	1.67 (0.11)	2.05 (0.18)	2.32 (0.25)	1.1	71.7	(7.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.5 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	311	0.92 (0.05)	0.43 (0.06)	0.51 (0.06)	0.67 (0.06)	0.88 (0.06)	1.15 (0.10)	1.43 (0.16)	1.62 (0.21)	0.7	71.4	(6.9)
	4-8	485	1.40 (0.07)	0.85 (0.15) ^E	0.95 (0.13)	1.14 (0.11)	1.39 (0.09)	1.69 (0.12)	2.02 (0.21)	2.24 (0.28)	0.9	92.7	(4.7)
Male	9-13	277	2.10 (0.17)	1.20 (0.19)	1.37 (0.19)	1.70 (0.20)	2.15 (0.23)	2.71 (0.28)	3.33 (0.35)	3.75 (0.42)	1.2	94.9	(4.2)
	14-18	339	2.68 (0.17)	1.39 (0.26) ^E	1.61 (0.25)	2.05 (0.23)	2.66 (0.22)	3.42 (0.28)	4.27 (0.41)	4.86 (0.54)	1.6	90.4	(5.4)
	19-30	237	2.43 (0.15)	1.44 (0.15)	1.63 (0.17)	1.98 (0.20)	2.43 (0.23)	2.98 (0.28)	3.60 (0.37)	4.06 (0.48)	1.6	90.8	(5.7)
	31-50	423	2.12 (0.17)	1.00 (0.22) ^E	1.19 (0.21) ^E	1.55 (0.20)	2.05 (0.20)	2.65 (0.25)	3.30 (0.36)	3.77 (0.47)	1.6	72.6	(10.1)
	51-70	387	2.29 (0.15)	1.13 (0.16)	1.31 (0.16)	1.65 (0.16)	2.14 (0.18)	2.77 (0.24)	3.50 (0.34)	4.02 (0.44)	1.6	77.4	(8.0)
	>70	132	1.77 (0.17)	0.94 (0.25) ^E	1.10 (0.24) ^E	1.40 (0.22)	1.81 (0.23)	2.28 (0.29)	2.78 (0.40)	3.12 (0.50)	1.6	63.1	(14.9) ^E
	19+	1179	2.21 (0.09)	1.16 (0.11)	1.33 (0.11)	1.67 (0.11)	2.14 (0.12)	2.73 (0.15)	3.38 (0.21)	3.85 (0.28)	1.6	78.6	(5.4)
Female	9-13	281	1.63 (0.09)	0.84 (0.14)	0.96 (0.13)	1.20 (0.12)	1.52 (0.11)	1.92 (0.14)	2.37 (0.21)	2.69 (0.28)	1.0	88.1	(6.0)
	14-18	321	1.95 (0.18)	1.34 (0.29) ^E	1.48 (0.29) ^E	1.74 (0.28)	2.09 (0.29)	2.50 (0.35)	2.93 (0.46)	3.23 (0.58) ^E	1.1	99.2	(5.8)
	19-30	249	2.00 (0.22)	0.92 (0.10)	1.05 (0.11)	1.31 (0.14)	1.71 (0.21)	2.28 (0.32)	3.00 (0.49)	3.54 (0.65) ^E	1.1	87.6	(6.0)
	31-50	364	1.81 (0.10)	1.07 (0.08)	1.20 (0.09)	1.47 (0.11)	1.85 (0.14)	2.31 (0.18)	2.80 (0.25)	3.13 (0.30)	1.1	94.0	(2.8)
	51-70	467	1.81 (0.11)	0.88 (0.17) ^E	1.03 (0.16)	1.32 (0.14)	1.74 (0.12)	2.26 (0.16)	2.85 (0.27)	3.26 (0.35)	1.1	86.9	(6.0)
	>70	215	1.30 (0.09)	0.93 (0.16) ^E	0.99 (0.15)	1.11 (0.14)	1.26 (0.13)	1.43 (0.14)	1.60 (0.18)	1.72 (0.22)	1.1	77.0	(18.3) ^E
	19+	1295	1.78 (0.07)	1.05 (0.10)	1.18 (0.10)	1.43 (0.09)	1.76 (0.08)	2.18 (0.11)	2.63 (0.17)	2.94 (0.23)	1.1	93.3	(3.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.6 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	644	0.90 (0.05)	0.44 (0.09) ^E	0.51 (0.09) ^E	0.64 (0.08)	0.86 (0.06)	1.15 (0.10)	1.49 (0.19)	1.75 (0.27)	0.7	68.0	(10.0)
	4-8	956	1.24 (0.05)	0.58 (0.06)	0.68 (0.06)	0.87 (0.05)	1.15 (0.05)	1.53 (0.07)	2.01 (0.14)	2.40 (0.21)	0.9	72.6	(4.8)
Male	9-13	589	1.58 (0.08)	0.94 (0.16) ^E	1.05 (0.15)	1.25 (0.11)	1.52 (0.08)	1.85 (0.12)	2.20 (0.23)	2.44 (0.33)	1.2	79.0	(9.3)
	14-18	639	2.04 (0.10)	1.15 (0.06)	1.30 (0.07)	1.58 (0.08)	1.96 (0.10)	2.41 (0.14)	2.89 (0.20)	3.22 (0.24)	1.6	73.9	(5.2)
	19-30	481	2.44 (0.19)	1.21 (0.38) ^E	1.41 (0.35) ^E	1.81 (0.30)	2.36 (0.24)	3.06 (0.31)	3.84 (0.58)	4.39 (0.84) ^E	1.6	83.5	(10.8)
	31-50	709	2.14 (0.25)	0.83 (0.26) ^E	0.98 (0.25) ^E	1.30 (0.22) ^E	1.80 (0.18)	2.54 (0.25)	3.53 (0.55)	4.34 (0.87) ^E	1.6	59.5	(12.2) ^E
	51-70	758	2.27 (0.21)	0.97 (0.22) ^E	1.13 (0.21) ^E	1.46 (0.19)	1.96 (0.17)	2.68 (0.21)	3.62 (0.42)	4.38 (0.65)	1.6	67.8	(9.5)
	>70	734	1.52 (0.06)	0.56 (0.07)	0.69 (0.07)	0.94 (0.07)	1.32 (0.07)	1.83 (0.08)	2.49 (0.15)	3.04 (0.23)	1.6	34.6	(3.5)
	19+	2682	2.19 (0.12)	0.86 (0.10)	1.03 (0.10)	1.36 (0.09)	1.89 (0.11)	2.72 (0.16)	3.80 (0.33)	4.68 (0.51)	1.6	63.1	(4.7)
Female	9-13	585	1.35 (0.05)	1.00 (0.16)	1.07 (0.14)	1.20 (0.10)	1.36 (0.06)	1.54 (0.10)	1.72 (0.19)	1.84 (0.27)	1.0	95.0	(9.5)
	14-18	645	1.77 (0.11)	0.98 (0.18) ^E	1.10 (0.16)	1.35 (0.13)	1.69 (0.10)	2.12 (0.16)	2.59 (0.30)	2.92 (0.41)	1.1	90.2	(6.1)
	19-30	514	1.40 (0.07)	0.65 (0.15) ^E	0.76 (0.14) ^E	0.98 (0.12)	1.29 (0.09)	1.68 (0.11)	2.15 (0.21)	2.49 (0.31)	1.1	65.0	(9.9)
	31-50	758	1.79 (0.11)	0.75 (0.12)	0.90 (0.12)	1.20 (0.11)	1.63 (0.11)	2.20 (0.16)	2.87 (0.29)	3.35 (0.41)	1.1	80.4	(6.1)
	51-70	955	1.64 (0.08)	0.88 (0.17) ^E	1.00 (0.16)	1.24 (0.13)	1.57 (0.09)	1.99 (0.12)	2.47 (0.25)	2.81 (0.37)	1.1	84.4	(8.2)
	>70	1345	1.44 (0.05)	0.84 (0.15) ^E	0.94 (0.14)	1.13 (0.11)	1.39 (0.08)	1.71 (0.11)	2.05 (0.21)	2.29 (0.31)	1.1	78.0	(10.2)
	19+	3572	1.63 (0.05)	0.79 (0.09)	0.91 (0.08)	1.16 (0.07)	1.53 (0.06)	2.01 (0.09)	2.56 (0.16)	2.96 (0.22)	1.1	79.2	(4.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.7 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	324	0.75 (0.04)	0.42 (0.05)	0.47 (0.05)	0.57 (0.04)	0.71 (0.05)	0.88 (0.08)	1.08 (0.13)	1.23 (0.18)	0.7	51.2	(9.8) ^E
	4-8	425	1.23 (0.08)	0.72 (0.18) ^E	0.81 (0.17) ^E	0.97 (0.14)	1.18 (0.09)	1.44 (0.09)	1.71 (0.19)	1.90 (0.28)	0.9	82.0	(12.1)
Male	9-13	274	1.59 (0.13)	0.82 (0.18) ^E	0.94 (0.16) ^E	1.17 (0.14)	1.52 (0.12)	2.00 (0.19)	2.59 (0.34)	3.05 (0.51) ^E	1.2	73.2	(10.0)
	14-18	297	2.29 (0.15)	0.94 (0.21) ^E	1.15 (0.21) ^E	1.58 (0.20)	2.19 (0.19)	2.99 (0.25)	3.89 (0.40)	4.52 (0.55)	1.6	74.1	(8.0)
	19-30	249	2.60 (0.29)	1.21 (0.11)	1.41 (0.13)	1.82 (0.17)	2.48 (0.25)	3.38 (0.42)	4.46 (0.71)	5.27 (0.99) ^E	1.6	83.6	(5.2)
	31-50	309	1.78 (0.10)	0.93 (0.20) ^E	1.08 (0.18) ^E	1.36 (0.15)	1.75 (0.13)	2.23 (0.17)	2.73 (0.30)	3.11 (0.43)	1.6	59.8	(10.3) ^E
	51-70	277	1.91 (0.19)	1.09 (0.26) ^E	1.22 (0.25) ^E	1.46 (0.21)	1.79 (0.18)	2.19 (0.25)	2.62 (0.45) ^E	2.93 (0.64) ^E	1.6	64.7	(14.6) ^E
	>70	136	1.46 (0.10)	0.78 (0.17) ^E	0.88 (0.16) ^E	1.08 (0.14)	1.34 (0.12)	1.66 (0.16)	2.00 (0.26)	2.24 (0.35)	1.6		^F
	19+	971	1.97 (0.10)	1.05 (0.19) ^E	1.19 (0.18)	1.49 (0.14)	1.91 (0.12)	2.44 (0.17)	3.04 (0.33)	3.47 (0.48)	1.6	68.5	(8.9)
Female	9-13	265	1.45 (0.10)	0.84 (0.17) ^E	0.93 (0.15)	1.10 (0.12)	1.31 (0.09)	1.55 (0.14)	1.79 (0.24)	1.94 (0.31)	1.0	84.6	(10.5)
	14-18	290	1.55 (0.11)	0.71 (0.09)	0.83 (0.10)	1.10 (0.12)	1.51 (0.16)	2.02 (0.23)	2.56 (0.36)	2.94 (0.50) ^E	1.1	75.2	(7.1)
	19-30	197	1.58 (0.13)	0.75 (0.07)	0.87 (0.09)	1.12 (0.11)	1.48 (0.14)	1.92 (0.18)	2.42 (0.24)	2.77 (0.29)	1.1	76.7	(7.2)
	31-50	312	1.77 (0.14)	1.11 (0.11)	1.23 (0.12)	1.46 (0.13)	1.77 (0.16)	2.14 (0.21)	2.53 (0.27)	2.80 (0.33)	1.1	95.3	(4.2)
	51-70	312	1.60 (0.09)	0.86 (0.19) ^E	0.96 (0.18) ^E	1.18 (0.15)	1.47 (0.12)	1.82 (0.15)	2.21 (0.26)	2.46 (0.35)	1.1	81.0	(10.5)
	>70	239	1.34 (0.12)	0.58 (0.13) ^E	0.68 (0.12) ^E	0.88 (0.12)	1.16 (0.12)	1.56 (0.16)	2.02 (0.26)	2.34 (0.35)	1.1	55.3	(10.5) ^E
	19+	1060	1.62 (0.07)	0.78 (0.11)	0.91 (0.11)	1.16 (0.10)	1.53 (0.08)	2.00 (0.11)	2.53 (0.20)	2.91 (0.28)	1.1	79.2	(6.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.8 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	1.06 (0.11)	0.63 (0.15) ^E	0.71 (0.14) ^E	0.88 (0.14)	1.11 (0.16)	1.39 (0.23) ^E	1.70 (0.35) ^E	1.92 (0.45) ^E	0.7	90.8	(9.7)
	4-8	213	1.40 (0.10)	0.81 (0.12)	0.90 (0.11)	1.07 (0.10)	1.31 (0.09)	1.60 (0.13)	1.92 (0.20)	2.14 (0.27)	0.9	89.8	(6.2)
Male	9-13	122	2.23 (0.57) ^E	0.93 (0.29) ^E	1.10 (0.30) ^E	1.45 (0.33) ^E	2.02 (0.43) ^E	2.84 (0.65) ^E	3.91 (1.03) ^E	4.74 (1.38) ^E	1.2	86.1	(10.4)
	14-18	150	2.23 (0.18)	1.39 (0.14)	1.55 (0.16)	1.84 (0.20)	2.23 (0.25)	2.72 (0.32)	3.26 (0.41)	3.63 (0.48)	1.6	87.7	(8.8)
	19-30	106	2.87 (0.43)	^F	1.12 (0.35) ^E	1.63 (0.35) ^E	2.44 (0.41)	3.59 (0.62) ^E	5.00 (1.08) ^E	6.07 (1.57) ^E	1.6	76.1	(10.7)
	31-50	155	2.10 (0.16)	0.99 (0.29) ^E	1.15 (0.27) ^E	1.47 (0.23)	1.95 (0.20)	2.55 (0.26)	3.19 (0.42)	3.63 (0.56)	1.6	68.1	(13.1) ^E
	51-70	122	2.10 (0.17)	1.71 (0.24)	1.84 (0.26)	2.08 (0.28)	2.37 (0.32)	2.69 (0.35)	3.00 (0.39)	3.20 (0.42)	1.6	97.5	(11.7)
	>70	88	2.37 (0.30)	1.02 (0.31) ^E	1.20 (0.31) ^E	1.58 (0.31) ^E	2.09 (0.35) ^E	2.77 (0.50) ^E	3.58 (0.76) ^E	4.14 (1.00) ^E	1.6	74.0	(14.5) ^E
	19+	471	2.31 (0.13)	1.06 (0.15)	1.24 (0.15)	1.61 (0.15)	2.17 (0.16)	2.91 (0.22)	3.76 (0.35)	4.38 (0.48)	1.6	75.5	(6.5)
Female	9-13	103	1.55 (0.16)	1.06 (0.23) ^E	1.15 (0.22) ^E	1.34 (0.22)	1.59 (0.22)	1.88 (0.28)	2.20 (0.42) ^E	2.42 (0.56) ^E	1.0	97.0	(8.6)
	14-18	142	1.72 (0.13)	0.86 (0.17) ^E	1.01 (0.16)	1.31 (0.16)	1.73 (0.19)	2.24 (0.27)	2.78 (0.37)	3.16 (0.46)	1.1	86.0	(8.0)
	19-30	111	1.72 (0.14)	1.10 (0.28) ^E	1.22 (0.26) ^E	1.45 (0.23)	1.76 (0.22)	2.13 (0.29)	2.52 (0.45) ^E	2.79 (0.60) ^E	1.1	94.9	(9.7)
	31-50	146	1.55 (0.14)	0.90 (0.12)	1.00 (0.13)	1.17 (0.14)	1.39 (0.15)	1.64 (0.18)	1.89 (0.21)	2.05 (0.24)	1.1	81.9	(9.9)
	51-70	184	2.10 (0.22)	^F	0.98 (0.28) ^E	1.32 (0.26) ^E	1.84 (0.25)	2.57 (0.32)	3.46 (0.54)	4.14 (0.77) ^E	1.1	85.3	(8.6)
	>70	143	1.54 (0.15)	0.82 (0.26) ^E	0.95 (0.24) ^E	1.20 (0.22) ^E	1.57 (0.22)	2.07 (0.30)	2.66 (0.48) ^E	3.09 (0.65) ^E	1.1	81.4	(12.4)
	19+	584	1.73 (0.09)	0.88 (0.16) ^E	1.01 (0.15)	1.27 (0.13)	1.63 (0.12)	2.09 (0.15)	2.61 (0.25)	2.99 (0.35)	1.1	85.2	(6.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.9 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	169	0.77 (0.06)	0.48 (0.04)	0.53 (0.05)	0.61 (0.05)	0.72 (0.06)	0.85 (0.08)	0.99 (0.10)	1.08 (0.11)	0.7	55.4	(13.8) ^E
	4-8	281	1.17 (0.07)	0.69 (0.05)	0.77 (0.06)	0.91 (0.07)	1.10 (0.08)	1.33 (0.09)	1.56 (0.11)	1.71 (0.12)	0.9	76.4	(7.8)
Male	9-13	183	1.74 (0.12)	0.89 (0.09)	1.04 (0.09)	1.32 (0.11)	1.70 (0.14)	2.18 (0.18)	2.71 (0.23)	3.08 (0.28)	1.2	81.9	(6.0)
	14-18	187	2.04 (0.13)	1.31 (0.11)	1.46 (0.12)	1.72 (0.13)	2.06 (0.16)	2.45 (0.19)	2.85 (0.22)	3.12 (0.25)	1.6	82.7	(7.4)
	19-30	223	2.21 (0.21)	0.85 (0.23) ^E	1.03 (0.23) ^E	1.43 (0.22)	2.04 (0.23)	2.86 (0.32)	3.84 (0.54)	4.57 (0.75)	1.6	68.0	(9.9)
	31-50	229	2.30 (0.32)	1.29 (0.17)	1.49 (0.19)	1.87 (0.23)	2.35 (0.30)	2.90 (0.39)	3.49 (0.52)	3.91 (0.63)	1.6	86.3	(9.9)
	51-70	197	1.74 (0.16)	1.17 (0.15)	1.28 (0.16)	1.49 (0.17)	1.74 (0.19)	2.03 (0.22)	2.33 (0.24)	2.52 (0.27)	1.6	64.1	(17.2) ^E
	>70	72	1.61 (0.16)	0.86 (0.24) ^E	1.00 (0.23) ^E	1.28 (0.20)	1.65 (0.20)	2.07 (0.27)	2.50 (0.41)	2.80 (0.54) ^E	1.6	53.6	(16.5) ^E
	19+	721	2.09 (0.16)	1.22 (0.28) ^E	1.40 (0.26) ^E	1.75 (0.23)	2.18 (0.20)	2.68 (0.23)	3.22 (0.35)	3.60 (0.47)	1.6	82.1	(10.6)
Female	9-13	165	1.23 (0.10)	0.71 (0.20) ^E	0.80 (0.19) ^E	0.98 (0.17) ^E	1.24 (0.16)	1.57 (0.20)	1.96 (0.31)	2.24 (0.42) ^E	1.0	73.2	(16.4) ^E
	14-18	206	1.57 (0.12)	0.63 (0.21) ^E	0.76 (0.20) ^E	1.02 (0.17)	1.41 (0.14)	1.92 (0.20)	2.52 (0.36)	2.96 (0.51) ^E	1.1	69.8	(11.5)
	19-30	191	1.48 (0.14)	0.52 (0.16) ^E	0.63 (0.16) ^E	0.89 (0.17) ^E	1.30 (0.19)	1.86 (0.25)	2.52 (0.38)	3.00 (0.51) ^E	1.1	61.5	(11.5) ^E
	31-50	258	1.59 (0.10)	1.03 (0.08)	1.12 (0.09)	1.31 (0.11)	1.56 (0.13)	1.86 (0.17)	2.17 (0.20)	2.37 (0.22)	1.1	91.5	(7.0)
	51-70	249	1.56 (0.11)	0.57 (0.11) ^E	0.71 (0.11)	1.00 (0.12)	1.45 (0.14)	2.05 (0.19)	2.72 (0.28)	3.21 (0.39)	1.1	69.3	(7.6)
	>70	128	1.22 (0.13)	0.45 (0.10) ^E	0.55 (0.11) ^E	0.76 (0.14) ^E	1.11 (0.18)	1.62 (0.24)	2.27 (0.36)	2.79 (0.48) ^E	1.1	50.7	(11.8) ^E
	19+	826	1.52 (0.06)	0.65 (0.07)	0.77 (0.07)	1.02 (0.08)	1.43 (0.09)	1.99 (0.12)	2.58 (0.16)	2.99 (0.19)	1.1	69.7	(5.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.10 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	192	1.00 (0.10)	0.49 (0.12) ^E	0.56 (0.12) ^E	0.68 (0.11)	0.87 (0.11)	1.10 (0.15)	1.37 (0.22)	1.56 (0.29) ^E	0.7	72.6	(13.7) ^E
	4-8	321	1.36 (0.12)	0.85 (0.16) ^E	0.94 (0.14)	1.10 (0.12)	1.31 (0.11)	1.56 (0.16)	1.85 (0.27)	2.04 (0.37) ^E	0.9	92.5	(7.9)
Male	9-13	226	1.91 (0.21)	0.85 (0.13)	0.99 (0.13)	1.27 (0.14)	1.68 (0.15)	2.26 (0.22)	2.99 (0.38)	3.56 (0.59)	1.2	78.9	(7.5)
	14-18	262	2.22 (0.18)	1.21 (0.15)	1.37 (0.17)	1.68 (0.19)	2.09 (0.23)	2.57 (0.28)	3.07 (0.35)	3.41 (0.40)	1.6	79.4	(8.9)
	19-30	197	2.59 (0.17)	2.06 (0.37) ^E	2.20 (0.34)	2.45 (0.28)	2.74 (0.24)	3.05 (0.27)	3.35 (0.38)	3.54 (0.48)	1.6	99.8	(4.3)
	31-50	282	2.26 (0.14)	0.99 (0.20) ^E	1.18 (0.19)	1.56 (0.18)	2.09 (0.18)	2.76 (0.23)	3.50 (0.34)	4.02 (0.45)	1.6	73.2	(8.3)
	51-70	234	2.24 (0.19)	0.93 (0.14)	1.10 (0.15)	1.47 (0.16)	2.01 (0.19)	2.74 (0.28)	3.62 (0.43)	4.27 (0.57)	1.6	68.8	(7.9)
	>70	119	1.79 (0.13)	1.14 (0.22) ^E	1.28 (0.21)	1.53 (0.19)	1.84 (0.19)	2.17 (0.23)	2.49 (0.30)	2.70 (0.35)	1.6	69.6	(16.3) ^E
	19+	832	2.28 (0.09)	1.06 (0.09)	1.25 (0.09)	1.62 (0.10)	2.17 (0.11)	2.85 (0.14)	3.62 (0.20)	4.15 (0.26)	1.6	76.1	(4.3)
Female	9-13	226	1.47 (0.10)	0.92 (0.17) ^E	1.02 (0.16)	1.22 (0.13)	1.47 (0.12)	1.78 (0.15)	2.10 (0.23)	2.32 (0.30)	1.0	91.2	(7.0)
	14-18	242	1.58 (0.11)	0.78 (0.17) ^E	0.91 (0.16) ^E	1.16 (0.15)	1.50 (0.15)	1.90 (0.17)	2.34 (0.25)	2.63 (0.33)	1.1	79.2	(10.0)
	19-30	208	1.72 (0.11)	0.67 (0.11)	0.82 (0.11)	1.14 (0.12)	1.59 (0.15)	2.17 (0.20)	2.82 (0.28)	3.27 (0.35)	1.1	77.1	(6.7)
	31-50	263	2.06 (0.18)	0.89 (0.20) ^E	1.06 (0.19) ^E	1.42 (0.18)	1.93 (0.19)	2.61 (0.27)	3.45 (0.46)	4.09 (0.65)	1.1	88.7	(5.9)
	51-70	322	1.65 (0.12)	0.60 (0.12) ^E	0.74 (0.12)	1.04 (0.13)	1.47 (0.15)	2.09 (0.19)	2.84 (0.28)	3.37 (0.36)	1.1	71.4	(7.6)
	>70	198	1.46 (0.11)	0.72 (0.16) ^E	0.83 (0.16) ^E	1.05 (0.15)	1.35 (0.15)	1.74 (0.19)	2.16 (0.29)	2.46 (0.39)	1.1	70.8	(12.4) ^E
	19+	991	1.81 (0.09)	0.71 (0.07)	0.87 (0.07)	1.19 (0.08)	1.66 (0.09)	2.30 (0.13)	3.10 (0.20)	3.70 (0.29)	1.1	79.9	(3.7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.11 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						AI ³	% >AI	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both	1-3	348	0.89	(0.06)	0.52 (0.07)	0.58 (0.07)	0.68 (0.07)	0.84 (0.07)	1.04 (0.10)	1.28 (0.15)	1.45 (0.20)	0.7	72.5 (10.6)
	4-8	554	1.31	(0.06)	0.89 (0.10)	0.97 (0.09)	1.12 (0.08)	1.32 (0.07)	1.56 (0.10)	1.81 (0.15)	1.98 (0.20)	0.9	94.3 (4.9)
Male	9-13	409	2.25	(0.46) ^E	1.26 (0.24) ^E	1.43 (0.25) ^E	1.77 (0.28)	2.16 (0.34)	2.57 (0.51) ^E	3.03 (0.73) ^E	3.36 (0.85) ^E	1.2	96.4 (6.0)
	14-18	414	2.01	(0.12)	1.08 (0.17)	1.24 (0.17)	1.57 (0.16)	2.00 (0.16)	2.50 (0.18)	3.02 (0.24)	3.36 (0.29)	1.6	73.3 (8.9)
	19-30	311	2.37	(0.26)	1.27 (0.21)	1.44 (0.20)	1.72 (0.20)	2.17 (0.23)	2.94 (0.41)	3.99 (0.88) ^E	4.89 (1.40) ^E	1.6	82.5 (7.7)
	31-50	489	2.02	(0.10)	0.94 (0.12)	1.10 (0.12)	1.42 (0.12)	1.86 (0.13)	2.41 (0.16)	3.04 (0.22)	3.47 (0.28)	1.6	64.6 (7.4)
	51-70	575	1.97	(0.15)	0.94 (0.15)	1.09 (0.14)	1.39 (0.14)	1.81 (0.14)	2.37 (0.20)	3.02 (0.31)	3.49 (0.41)	1.6	62.5 (8.6)
	>70	239	1.75	(0.10)	0.82 (0.12)	0.97 (0.12)	1.28 (0.12)	1.71 (0.14)	2.26 (0.19)	2.88 (0.31)	3.32 (0.45)	1.6	56.4 (8.2)
	19+	1614	2.05	(0.08)	0.92 (0.06)	1.09 (0.06)	1.41 (0.06)	1.87 (0.07)	2.53 (0.11)	3.34 (0.22)	3.97 (0.35)	1.6	64.3 (3.8)
Female	9-13	355	1.35	(0.07)	0.81 (0.07)	0.90 (0.08)	1.07 (0.08)	1.29 (0.09)	1.57 (0.11)	1.88 (0.13)	2.09 (0.15)	1.0	81.5 (7.0)
	14-18	410	1.56	(0.19)	0.75 (0.13) ^E	0.86 (0.13)	1.08 (0.12)	1.39 (0.14)	1.79 (0.18)	2.25 (0.29)	2.59 (0.38)	1.1	73.1 (9.5)
	19-30	384	1.62	(0.11)	1.09 (0.16)	1.19 (0.15)	1.38 (0.14)	1.64 (0.14)	1.93 (0.18)	2.24 (0.27)	2.46 (0.34)	1.1	94.6 (5.9)
	31-50	585	1.65	(0.09)	0.73 (0.09)	0.86 (0.09)	1.13 (0.09)	1.52 (0.11)	2.04 (0.14)	2.65 (0.22)	3.10 (0.29)	1.1	77.2 (5.8)
	51-70	711	1.54	(0.07)	0.78 (0.08)	0.89 (0.08)	1.12 (0.08)	1.44 (0.08)	1.85 (0.11)	2.28 (0.17)	2.58 (0.22)	1.1	76.5 (6.1)
	>70	342	1.33	(0.09)	0.66 (0.09)	0.76 (0.09)	0.98 (0.10)	1.28 (0.11)	1.66 (0.14)	2.10 (0.19)	2.41 (0.25)	1.1	64.8 (9.1)
	19+	2022	1.57	(0.05)	0.79 (0.05)	0.91 (0.05)	1.15 (0.05)	1.50 (0.06)	1.96 (0.07)	2.48 (0.11)	2.85 (0.15)	1.1	78.8 (3.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.12 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	622	0.82 (0.04)	0.51 (0.07)	0.56 (0.07)	0.65 (0.06)	0.78 (0.05)	0.93 (0.06)	1.10 (0.09)	1.22 (0.12)	0.7	65.4	(11.9) ^E
	4-8	919	1.22 (0.05)	0.70 (0.10)	0.78 (0.09)	0.94 (0.07)	1.15 (0.06)	1.41 (0.08)	1.69 (0.15)	1.88 (0.20)	0.9	79.3	(7.8)
Male	9-13	579	1.80 (0.13)	0.88 (0.17) ^E	1.01 (0.17)	1.30 (0.15)	1.73 (0.14)	2.30 (0.22)	2.98 (0.41)	3.50 (0.54)	1.2	80.4	(7.5)
	14-18	634	2.13 (0.09)	1.32 (0.23) ^E	1.48 (0.21)	1.77 (0.17)	2.16 (0.13)	2.63 (0.16)	3.11 (0.27)	3.44 (0.36)	1.6	84.4	(8.2)
	19-30	578	2.39 (0.16)	1.02 (0.20) ^E	1.23 (0.20)	1.64 (0.19)	2.24 (0.18)	3.04 (0.25)	4.00 (0.42)	4.71 (0.60)	1.6	76.7	(7.5)
	31-50	693	2.16 (0.21)	1.20 (0.31) ^E	1.37 (0.29) ^E	1.77 (0.26)	2.25 (0.24)	2.68 (0.25)	3.16 (0.38)	3.51 (0.52)	1.6	82.1	(11.7)
	51-70	596	1.85 (0.11)	1.21 (0.25) ^E	1.34 (0.23) ^E	1.57 (0.19)	1.87 (0.15)	2.24 (0.18)	2.62 (0.30)	2.87 (0.41)	1.6	72.5	(13.8) ^E
	>70	296	1.75 (0.12)	0.78 (0.12)	0.92 (0.12)	1.21 (0.12)	1.63 (0.14)	2.15 (0.18)	2.75 (0.27)	3.21 (0.36)	1.6	51.8	(8.2)
	19+	2163	2.10 (0.10)	0.94 (0.05)	1.11 (0.05)	1.48 (0.07)	2.01 (0.09)	2.72 (0.14)	3.56 (0.21)	4.18 (0.27)	1.6	69.5	(3.5)
Female	9-13	533	1.34 (0.07)	0.74 (0.14) ^E	0.84 (0.13)	1.03 (0.12)	1.30 (0.11)	1.64 (0.13)	2.01 (0.19)	2.28 (0.26)	1.0	78.0	(9.8)
	14-18	638	1.59 (0.08)	0.71 (0.09)	0.84 (0.09)	1.10 (0.09)	1.47 (0.09)	1.96 (0.13)	2.55 (0.20)	2.96 (0.26)	1.1	74.7	(5.7)
	19-30	499	1.54 (0.09)	0.72 (0.14) ^E	0.84 (0.13)	1.09 (0.12)	1.44 (0.13)	1.88 (0.16)	2.37 (0.25)	2.71 (0.32)	1.1	74.2	(8.6)
	31-50	716	1.62 (0.08)	1.01 (0.19) ^E	1.11 (0.18)	1.31 (0.14)	1.58 (0.11)	1.91 (0.13)	2.26 (0.23)	2.51 (0.31)	1.1	90.6	(8.4)
	51-70	745	1.67 (0.08)	0.71 (0.09)	0.84 (0.09)	1.11 (0.09)	1.53 (0.10)	2.09 (0.13)	2.74 (0.20)	3.21 (0.28)	1.1	75.6	(5.5)
	>70	510	1.33 (0.08)	0.56 (0.07)	0.66 (0.07)	0.88 (0.09)	1.24 (0.11)	1.74 (0.14)	2.34 (0.20)	2.76 (0.26)	1.1	59.3	(7.1)
	19+	2470	1.58 (0.04)	0.73 (0.05)	0.85 (0.05)	1.10 (0.06)	1.49 (0.06)	2.00 (0.07)	2.58 (0.11)	2.99 (0.15)	1.1	75.3	(3.7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 30.13 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	2117	0.90 (0.03)	0.46 (0.03)	0.53 (0.03)	0.65 (0.03)	0.84 (0.03)	1.07 (0.05)	1.36 (0.08)	1.58 (0.11)	0.7	68.7	(4.1)
	4-8	3235	1.29 (0.03)	0.69 (0.04)	0.78 (0.04)	0.97 (0.03)	1.23 (0.03)	1.56 (0.05)	1.95 (0.08)	2.24 (0.12)	0.9	81.1	(2.9)
Male	9-13	2080	1.83 (0.07)	0.95 (0.06)	1.08 (0.06)	1.35 (0.06)	1.73 (0.07)	2.23 (0.10)	2.81 (0.16)	3.24 (0.22)	1.2	84.0	(3.2)
	14-18	2288	2.22 (0.07)	1.27 (0.10)	1.43 (0.09)	1.75 (0.08)	2.17 (0.08)	2.70 (0.10)	3.26 (0.16)	3.64 (0.20)	1.6	82.9	(4.2)
	19-30	1804	2.45 (0.09)	1.24 (0.10)	1.44 (0.10)	1.83 (0.10)	2.39 (0.11)	3.10 (0.16)	3.91 (0.27)	4.50 (0.37)	1.6	84.4	(3.8)
	31-50	2596	2.15 (0.11)	0.92 (0.09)	1.09 (0.09)	1.45 (0.09)	1.96 (0.09)	2.66 (0.14)	3.54 (0.28)	4.23 (0.42)	1.6	67.4	(4.4)
	51-70	2550	2.18 (0.09)	0.96 (0.07)	1.13 (0.07)	1.48 (0.07)	1.96 (0.07)	2.65 (0.11)	3.54 (0.20)	4.21 (0.29)	1.6	68.6	(3.7)
	>70	1520	1.67 (0.05)	0.75 (0.05)	0.89 (0.05)	1.17 (0.06)	1.57 (0.06)	2.08 (0.09)	2.69 (0.13)	3.14 (0.17)	1.6	47.9	(3.9)
	19+	8470	2.18 (0.05)	0.95 (0.04)	1.13 (0.04)	1.49 (0.04)	2.00 (0.05)	2.69 (0.07)	3.64 (0.17)	4.42 (0.29)	1.6	69.6	(2.1)
Female	9-13	1980	1.43 (0.03)	0.84 (0.06)	0.94 (0.05)	1.13 (0.05)	1.38 (0.04)	1.68 (0.05)	2.01 (0.08)	2.24 (0.10)	1.0	86.0	(3.5)
	14-18	2256	1.74 (0.06)	0.89 (0.07)	1.03 (0.07)	1.29 (0.07)	1.67 (0.07)	2.16 (0.10)	2.71 (0.15)	3.10 (0.20)	1.1	86.5	(3.5)
	19-30	1854	1.62 (0.06)	0.77 (0.06)	0.89 (0.06)	1.14 (0.06)	1.48 (0.06)	1.92 (0.09)	2.42 (0.15)	2.78 (0.20)	1.1	77.7	(4.2)
	31-50	2686	1.79 (0.06)	0.91 (0.08)	1.04 (0.07)	1.33 (0.07)	1.73 (0.07)	2.24 (0.09)	2.82 (0.15)	3.23 (0.21)	1.1	87.5	(3.4)
	51-70	3200	1.68 (0.04)	0.78 (0.05)	0.91 (0.05)	1.19 (0.05)	1.59 (0.05)	2.11 (0.07)	2.72 (0.10)	3.15 (0.14)	1.1	80.3	(3.0)
	>70	2610	1.38 (0.04)	0.71 (0.05)	0.82 (0.05)	1.03 (0.05)	1.32 (0.05)	1.69 (0.07)	2.11 (0.10)	2.40 (0.14)	1.1	69.0	(4.3)
	19+	10350	1.68 (0.03)	0.80 (0.03)	0.93 (0.03)	1.20 (0.03)	1.58 (0.03)	2.08 (0.05)	2.67 (0.08)	3.10 (0.10)	1.1	81.3	(1.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

31. Percentage of total energy intake from linolenic acid

Table 31.1 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	0.49 (0.06)	0.31 (0.06) ^E	0.33 (0.06) ^E	0.38 (0.06)	0.46 (0.06)	0.55 (0.09)	0.66 (0.13) ^E	0.76 (0.18) ^E
	4-8	127	0.62 (0.04)	0.41 (0.05)	0.45 (0.05)	0.52 (0.05)	0.62 (0.05)	0.73 (0.06)	0.86 (0.09)	0.95 (0.11)
Male	9-13	111	0.64 (0.04)	0.47 (0.07)	0.50 (0.07)	0.57 (0.06)	0.64 (0.05)	0.71 (0.06)	0.79 (0.08)	0.83 (0.09)
	14-18	107	0.62 (0.04)	0.32 (0.06) ^E	0.37 (0.06)	0.47 (0.05)	0.59 (0.04)	0.74 (0.06)	0.89 (0.09)	0.99 (0.11)
	19-30	77	0.67 (0.06)	0.39 (0.09) ^E	0.43 (0.09) ^E	0.51 (0.09) ^E	0.60 (0.09)	0.71 (0.10)	0.82 (0.11)	0.90 (0.12)
	31-50	145	0.76 (0.07)	0.52 (0.09) ^E	0.56 (0.08)	0.63 (0.08)	0.72 (0.07)	0.83 (0.09)	0.94 (0.13)	1.02 (0.17) ^E
	51-70	182	0.70 (0.05)	0.53 (0.10) ^E	0.55 (0.09)	0.60 (0.07)	0.65 (0.05)	0.71 (0.08)	0.77 (0.15) ^E	0.81 (0.20) ^E
	>70	63	0.79 (0.08)	0.48 (0.10) ^E	0.54 (0.09) ^E	0.66 (0.08)	0.81 (0.08)	0.98 (0.10)	1.13 (0.15)	1.22 (0.19)
	19+	467	0.73 (0.03)	0.40 (0.04)	0.45 (0.04)	0.54 (0.03)	0.67 (0.03)	0.84 (0.05)	1.02 (0.07)	1.14 (0.09)
Female	9-13	96	0.56 (0.03)	0.44 (0.04)	0.46 (0.04)	0.49 (0.04)	0.54 (0.04)	0.58 (0.05)	0.63 (0.05)	0.66 (0.05)
	14-18	105	0.58 (0.06)	0.36 (0.04)	0.39 (0.04)	0.43 (0.04)	0.50 (0.05)	0.57 (0.06)	0.66 (0.08)	0.72 (0.09)
	19-30	91	0.70 (0.05)	0.59 (0.06)	0.61 (0.07)	0.65 (0.07)	0.70 (0.07)	0.75 (0.08)	0.79 (0.08)	0.82 (0.08)
	31-50	167	0.81 (0.06)	0.64 (0.12) ^E	0.68 (0.11)	0.74 (0.09)	0.82 (0.08)	0.91 (0.09)	1.00 (0.14)	1.06 (0.20) ^E
	51-70	198	0.78 (0.06)	0.51 (0.10) ^E	0.56 (0.09)	0.65 (0.08)	0.78 (0.07)	0.94 (0.09)	1.12 (0.15)	1.24 (0.19)
	>70	74	0.76 (0.05)	0.53 (0.10) ^E	0.58 (0.09)	0.67 (0.08)	0.79 (0.08)	0.92 (0.09)	1.04 (0.12)	1.12 (0.15)
	19+	530	0.78 (0.03)	0.56 (0.07)	0.60 (0.07)	0.68 (0.05)	0.79 (0.04)	0.92 (0.06)	1.05 (0.10)	1.14 (0.13)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.2 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Prince Edward Island, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	0.45 (0.04)	0.26 (0.06) ^E	0.29 (0.05) ^E	0.34 (0.05)	0.39 (0.04)	0.46 (0.05)	0.54 (0.06)	0.59 (0.07)
	4-8	110	0.56 (0.02)	0.45 (0.05)	0.47 (0.04)	0.51 (0.04)	0.56 (0.03)	0.62 (0.05)	0.67 (0.07)	0.70 (0.09)
Male	9-13	95	0.55 (0.04)	0.43 (0.06)	0.46 (0.06)	0.51 (0.05)	0.57 (0.05)	0.64 (0.06)	0.71 (0.07)	0.76 (0.08)
	14-18	87	0.63 (0.05)	0.44 (0.08) ^E	0.47 (0.07)	0.53 (0.07)	0.60 (0.07)	0.68 (0.08)	0.76 (0.10)	0.81 (0.11)
	19-30	70	0.80 (0.07)	0.65 (0.15) ^E	0.67 (0.13) ^E	0.70 (0.09)	0.74 (0.08)	0.78 (0.14) ^E	0.82 (0.23) ^E	^F
	31-50	109	0.84 (0.06)	0.57 (0.06)	0.62 (0.06)	0.74 (0.08)	0.89 (0.10)	1.08 (0.12)	1.27 (0.15)	1.41 (0.18)
	51-70	128	0.79 (0.07)	0.46 (0.05)	0.52 (0.05)	0.64 (0.06)	0.80 (0.08)	0.99 (0.11)	1.22 (0.14)	1.38 (0.16)
	>70	65	0.78 (0.06)	0.48 (0.08) ^E	0.53 (0.08)	0.63 (0.07)	0.76 (0.07)	0.91 (0.09)	1.08 (0.14)	1.20 (0.19)
	19+	372	0.81 (0.04)	0.52 (0.07)	0.58 (0.07)	0.68 (0.06)	0.83 (0.06)	1.01 (0.07)	1.21 (0.10)	1.34 (0.13)
Female	9-13	75	0.69 (0.07)	0.56 (0.04)	0.58 (0.05)	0.63 (0.05)	0.69 (0.07)	0.76 (0.09)	0.84 (0.11)	0.89 (0.13)
	14-18	81	0.67 (0.06)	0.45 (0.07)	0.48 (0.08)	0.55 (0.09)	0.63 (0.10)	0.73 (0.11)	0.84 (0.13)	0.90 (0.14)
	19-30	101	0.70 (0.07)	0.59 (0.07)	0.61 (0.07)	0.66 (0.08)	0.72 (0.10)	0.79 (0.13)	0.86 (0.15) ^E	0.90 (0.18) ^E
	31-50	116	0.74 (0.06)	0.52 (0.08)	0.56 (0.08)	0.63 (0.07)	0.72 (0.07)	0.83 (0.09)	0.94 (0.13)	1.01 (0.16)
	51-70	146	0.80 (0.05)	0.57 (0.04)	0.61 (0.05)	0.68 (0.05)	0.77 (0.06)	0.87 (0.07)	0.97 (0.08)	1.03 (0.09)
	>70	94	0.85 (0.11)	0.48 (0.07)	0.53 (0.08)	0.65 (0.11)	0.81 (0.15) ^E	1.02 (0.21) ^E	1.26 (0.28) ^E	1.43 (0.34) ^E
	19+	457	0.76 (0.04)	0.54 (0.06)	0.58 (0.06)	0.65 (0.05)	0.75 (0.04)	0.85 (0.06)	0.97 (0.09)	1.05 (0.11)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.3 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Nova Scotia, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	0.60 (0.10)	0.39 (0.04)	0.42 (0.04)	0.47 (0.05)	0.56 (0.07)	0.67 (0.12) ^E	0.80 (0.19) ^E	0.90 (0.26) ^E
	4-8	177	0.58 (0.05)	0.33 (0.05)	0.37 (0.04)	0.44 (0.04)	0.54 (0.04)	0.70 (0.07)	0.90 (0.14)	1.06 (0.23) ^E
Male	9-13	111	0.58 (0.03)	0.42 (0.07) ^E	0.45 (0.07)	0.50 (0.05)	0.56 (0.04)	0.62 (0.05)	0.68 (0.08)	0.71 (0.11)
	14-18	113	0.62 (0.05)	0.38 (0.08) ^E	0.42 (0.07) ^E	0.50 (0.07)	0.60 (0.07)	0.73 (0.08)	0.85 (0.10)	0.92 (0.12)
	19-30	91	0.91 (0.27) ^E	0.46 (0.06)	0.50 (0.07)	0.59 (0.11) ^E	0.76 (0.20) ^E	F	F	F
	31-50	101	0.63 (0.05)	0.35 (0.06)	0.39 (0.05)	0.46 (0.05)	0.56 (0.05)	0.67 (0.07)	0.79 (0.09)	0.87 (0.12)
	51-70	134	0.79 (0.08)	0.39 (0.08) ^E	0.45 (0.08) ^E	0.56 (0.08)	0.73 (0.09)	0.97 (0.13)	1.26 (0.20)	1.48 (0.26) ^E
	>70	56	0.68 (0.04)	0.39 (0.07) ^E	0.44 (0.06)	0.53 (0.05)	0.63 (0.04)	0.74 (0.06)	0.85 (0.08)	0.93 (0.10)
	19+	382	0.74 (0.06)	0.36 (0.03)	0.41 (0.03)	0.51 (0.03)	0.64 (0.04)	0.84 (0.07)	1.09 (0.16)	1.28 (0.28) ^E
Female	9-13	105	0.57 (0.03)	0.47 (0.03)	0.49 (0.03)	0.52 (0.03)	0.55 (0.04)	0.59 (0.04)	0.62 (0.04)	0.65 (0.04)
	14-18	120	0.73 (0.18) ^E	0.39 (0.11) ^E	0.43 (0.11) ^E	0.50 (0.11) ^E	0.60 (0.13) ^E	0.76 (0.17) ^E	0.98 (0.24) ^E	1.17 (0.32) ^E
	19-30	91	0.82 (0.11)	0.65 (0.10)	0.68 (0.10)	0.72 (0.11)	0.77 (0.12)	0.83 (0.13)	0.88 (0.15)	0.92 (0.16) ^E
	31-50	159	0.93 (0.08)	0.50 (0.11) ^E	0.56 (0.11) ^E	0.69 (0.09)	0.86 (0.09)	1.08 (0.12)	1.33 (0.20)	1.51 (0.26) ^E
	51-70	174	0.83 (0.05)	0.50 (0.09) ^E	0.55 (0.08)	0.66 (0.06)	0.80 (0.06)	0.97 (0.09)	1.14 (0.14)	1.26 (0.19)
	>70	80	0.76 (0.08)	0.53 (0.11) ^E	0.58 (0.11) ^E	0.66 (0.10)	0.76 (0.10)	0.89 (0.12)	1.02 (0.17) ^E	1.11 (0.22) ^E
	19+	504	0.86 (0.04)	0.51 (0.06)	0.56 (0.06)	0.67 (0.05)	0.82 (0.04)	1.00 (0.06)	1.19 (0.09)	1.32 (0.12)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.4 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, New Brunswick, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	0.49 (0.05)	0.27 (0.05) ^E	0.30 (0.05) ^E	0.37 (0.05)	0.46 (0.05)	0.55 (0.06)	0.66 (0.08)	0.74 (0.09)
	4-8	140	0.59 (0.04)	0.42 (0.02)	0.45 (0.03)	0.49 (0.03)	0.55 (0.04)	0.62 (0.05)	0.69 (0.06)	0.73 (0.08)
Male	9-13	92	1.05 (0.34) ^E	0.48 (0.14) ^E	0.55 (0.14) ^E	0.66 (0.16) ^E	0.82 (0.20) ^E	1.04 (0.28) ^E	1.30 (0.37) ^E	1.48 (0.45) ^E
	14-18	107	0.64 (0.07)	0.46 (0.09) ^E	0.49 (0.08)	0.54 (0.08)	0.61 (0.07)	0.67 (0.08)	0.74 (0.11)	0.79 (0.13)
	19-30	73	0.70 (0.09)	0.43 (0.09) ^E	0.47 (0.08) ^E	0.55 (0.08)	0.65 (0.08)	0.76 (0.09)	0.88 (0.12)	0.95 (0.15)
	31-50	134	0.68 (0.05)	0.45 (0.08) ^E	0.49 (0.07)	0.57 (0.06)	0.66 (0.05)	0.79 (0.10)	^F	^F
	51-70	131	0.80 (0.06)	0.51 (0.10) ^E	0.56 (0.09)	0.64 (0.08)	0.75 (0.07)	0.88 (0.09)	1.00 (0.13)	1.09 (0.17)
	>70	55	0.88 (0.07)	0.55 (0.12) ^E	0.62 (0.12) ^E	0.76 (0.10)	0.93 (0.10)	1.13 (0.12)	1.34 (0.17)	1.48 (0.23)
	19+	393	0.73 (0.03)	0.44 (0.05)	0.49 (0.04)	0.59 (0.04)	0.72 (0.03)	0.87 (0.05)	1.03 (0.08)	1.14 (0.10)
Female	9-13	79	0.57 (0.06)	0.41 (0.06)	0.44 (0.06)	0.48 (0.07)	0.54 (0.07)	0.60 (0.08)	0.66 (0.09)	0.70 (0.10)
	14-18	104	0.66 (0.06)	0.46 (0.05)	0.50 (0.05)	0.57 (0.06)	0.67 (0.07)	0.78 (0.09)	0.90 (0.11)	0.98 (0.13)
	19-30	101	0.72 (0.07)	0.60 (0.07)	0.62 (0.07)	0.67 (0.08)	0.72 (0.08)	0.77 (0.09)	0.83 (0.10)	0.86 (0.10)
	31-50	143	0.70 (0.06)	0.47 (0.04)	0.50 (0.05)	0.57 (0.05)	0.66 (0.06)	0.77 (0.07)	0.88 (0.09)	0.95 (0.11)
	51-70	193	0.86 (0.05)	0.62 (0.05)	0.66 (0.05)	0.73 (0.06)	0.83 (0.06)	0.94 (0.07)	1.05 (0.08)	1.12 (0.08)
	>70	94	0.74 (0.06)	0.43 (0.08) ^E	0.47 (0.08) ^E	0.56 (0.08)	0.68 (0.09)	0.86 (0.10)	1.05 (0.13)	1.18 (0.16)
	19+	531	0.75 (0.03)	0.53 (0.03)	0.57 (0.03)	0.65 (0.03)	0.74 (0.04)	0.85 (0.05)	0.96 (0.06)	1.03 (0.07)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.5 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Quebec, 2004^{1–3}

Sex	Age (years)	Percentiles (and <i>SE</i>) of usual intake								
		n	Mean (<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)	95th (<i>SE</i>)
Both	1-3	311	0.49 (0.02)	0.33 (0.03)	0.36 (0.03)	0.41 (0.02)	0.48 (0.02)	0.56 (0.03)	0.64 (0.05)	0.70 (0.06)
	4-8	485	0.61 (0.03)	0.45 (0.02)	0.48 (0.02)	0.53 (0.03)	0.59 (0.03)	0.67 (0.04)	0.74 (0.04)	0.79 (0.05)
Male	9-13	277	0.68 (0.04)	0.43 (0.05)	0.48 (0.05)	0.56 (0.05)	0.68 (0.05)	0.82 (0.07)	0.98 (0.11)	1.09 (0.13)
	14-18	339	0.74 (0.04)	0.43 (0.06)	0.48 (0.06)	0.58 (0.05)	0.71 (0.05)	0.86 (0.07)	1.02 (0.10)	1.13 (0.12)
	19-30	237	0.78 (0.05)	0.50 (0.04)	0.55 (0.05)	0.65 (0.06)	0.78 (0.07)	0.94 (0.10)	1.13 (0.14)	1.26 (0.18)
	31-50	423	0.68 (0.03)	0.53 (0.07)	0.56 (0.07)	0.62 (0.05)	0.69 (0.04)	0.76 (0.05)	0.83 (0.07)	0.88 (0.09)
	51-70	387	0.86 (0.04)	0.56 (0.07)	0.61 (0.07)	0.71 (0.06)	0.84 (0.05)	0.99 (0.07)	1.14 (0.10)	1.24 (0.13)
	>70	132	0.79 (0.05)	0.50 (0.05)	0.55 (0.05)	0.65 (0.06)	0.78 (0.07)	0.94 (0.09)	1.11 (0.11)	1.24 (0.14)
	19+	1179	0.76 (0.02)	0.58 (0.06)	0.61 (0.05)	0.68 (0.04)	0.76 (0.03)	0.86 (0.04)	0.95 (0.06)	1.02 (0.08)
Female	9-13	281	0.66 (0.03)	0.44 (0.06)	0.48 (0.05)	0.55 (0.04)	0.64 (0.04)	0.75 (0.05)	0.86 (0.08)	0.93 (0.10)
	14-18	321	0.76 (0.07)	0.65 (0.12) ^E	0.67 (0.11) ^E	0.72 (0.10)	0.79 (0.09)	0.85 (0.10)	0.92 (0.13)	0.97 (0.15)
	19-30	249	0.79 (0.06)	0.55 (0.09)	0.59 (0.09)	0.65 (0.08)	0.74 (0.07)	0.85 (0.09)	0.96 (0.13)	1.04 (0.16)
	31-50	364	0.80 (0.04)	0.62 (0.08)	0.65 (0.07)	0.72 (0.06)	0.81 (0.05)	0.90 (0.06)	1.00 (0.10)	1.06 (0.13)
	51-70	467	0.87 (0.04)	0.54 (0.09)	0.60 (0.08)	0.71 (0.06)	0.85 (0.05)	1.02 (0.07)	1.22 (0.12)	1.35 (0.16)
	>70	215	0.77 (0.05)	0.56 (0.08)	0.60 (0.08)	0.67 (0.07)	0.76 (0.06)	0.86 (0.08)	0.96 (0.10)	1.03 (0.13)
	19+	1295	0.81 (0.03)	0.55 (0.04)	0.60 (0.04)	0.69 (0.03)	0.80 (0.03)	0.94 (0.04)	1.09 (0.06)	1.19 (0.08)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.6 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Ontario, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	0.53 (0.02)	0.34 (0.01)	0.37 (0.01)	0.42 (0.02)	0.50 (0.02)	0.61 (0.03)	0.73 (0.05)	0.82 (0.07)
	4-8	956	0.59 (0.02)	0.33 (0.03)	0.37 (0.03)	0.44 (0.02)	0.55 (0.02)	0.69 (0.03)	0.87 (0.05)	1.01 (0.08)
Male	9-13	589	0.60 (0.02)	0.38 (0.02)	0.42 (0.02)	0.48 (0.02)	0.57 (0.02)	0.67 (0.03)	0.78 (0.04)	0.85 (0.04)
	14-18	639	0.69 (0.05)	0.44 (0.02)	0.47 (0.02)	0.54 (0.02)	0.63 (0.03)	0.74 (0.05)	0.87 (0.09)	0.95 (0.13)
	19-30	481	0.79 (0.04)	0.42 (0.09) ^E	0.48 (0.08) ^E	0.59 (0.07)	0.76 (0.05)	0.96 (0.07)	1.19 (0.13)	1.36 (0.19)
	31-50	709	0.74 (0.05)	0.47 (0.09) ^E	0.51 (0.08)	0.60 (0.07)	0.71 (0.05)	0.85 (0.07)	0.99 (0.12)	1.10 (0.17)
	51-70	758	0.86 (0.04)	0.44 (0.06)	0.50 (0.06)	0.62 (0.05)	0.80 (0.04)	1.03 (0.06)	1.30 (0.11)	1.50 (0.16)
	>70	734	0.75 (0.03)	0.34 (0.03)	0.40 (0.03)	0.51 (0.03)	0.66 (0.03)	0.88 (0.04)	1.15 (0.06)	1.37 (0.09)
	19+	2682	0.78 (0.02)	0.38 (0.03)	0.44 (0.03)	0.55 (0.03)	0.72 (0.03)	0.95 (0.04)	1.22 (0.07)	1.42 (0.10)
Female	9-13	585	0.61 (0.02)	0.47 (0.02)	0.50 (0.02)	0.55 (0.02)	0.61 (0.03)	0.69 (0.03)	0.78 (0.05)	0.83 (0.05)
	14-18	645	0.77 (0.04)	0.50 (0.02)	0.55 (0.02)	0.63 (0.03)	0.73 (0.04)	0.85 (0.05)	0.98 (0.07)	1.07 (0.09)
	19-30	514	0.70 (0.03)	0.47 (0.07)	0.51 (0.07)	0.58 (0.05)	0.67 (0.04)	0.77 (0.05)	0.88 (0.09)	0.95 (0.13)
	31-50	758	0.86 (0.04)	0.45 (0.06)	0.51 (0.06)	0.63 (0.05)	0.80 (0.04)	1.03 (0.06)	1.30 (0.12)	1.48 (0.16)
	51-70	955	0.84 (0.03)	0.54 (0.09)	0.60 (0.08)	0.70 (0.06)	0.83 (0.04)	0.98 (0.05)	1.15 (0.11)	1.26 (0.15)
	>70	1345	0.81 (0.03)	0.52 (0.07)	0.56 (0.07)	0.66 (0.05)	0.78 (0.03)	0.94 (0.05)	1.12 (0.10)	1.24 (0.14)
	19+	3572	0.82 (0.02)	0.47 (0.04)	0.53 (0.04)	0.64 (0.03)	0.79 (0.03)	0.97 (0.03)	1.17 (0.06)	1.31 (0.09)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.7 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Manitoba, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	0.48 (0.02)	0.42 (0.02)	0.43 (0.02)	0.45 (0.03)	0.47 (0.03)	0.49 (0.03)	0.51 (0.03)	0.52 (0.03)
	4-8	425	0.61 (0.06)	0.43 (0.04)	0.46 (0.05)	0.52 (0.05)	0.59 (0.06)	0.67 (0.07)	0.76 (0.08)	0.82 (0.08)
Male	9-13	274	0.58 (0.04)	0.37 (0.03)	0.40 (0.03)	0.47 (0.03)	0.57 (0.04)	0.68 (0.04)	0.81 (0.06)	0.90 (0.07)
	14-18	297	0.66 (0.03)	0.35 (0.06)	0.41 (0.05)	0.50 (0.04)	0.63 (0.03)	0.79 (0.05)	0.96 (0.08)	1.07 (0.11)
	19-30	249	0.83 (0.06)	0.53 (0.04)	0.59 (0.04)	0.70 (0.05)	0.84 (0.06)	1.01 (0.08)	1.19 (0.10)	1.31 (0.12)
	31-50	309	0.71 (0.04)	0.40 (0.07) ^E	0.45 (0.07)	0.55 (0.06)	0.69 (0.05)	0.86 (0.06)	1.04 (0.09)	1.15 (0.12)
	51-70	277	0.78 (0.05)	0.48 (0.09) ^E	0.53 (0.08)	0.62 (0.06)	0.74 (0.05)	0.89 (0.07)	1.06 (0.13)	1.18 (0.19)
	>70	136	0.74 (0.04)	0.43 (0.08) ^E	0.48 (0.07)	0.57 (0.06)	0.70 (0.05)	0.85 (0.08)	1.01 (0.12)	1.11 (0.15)
	19+	971	0.76 (0.02)	0.45 (0.04)	0.50 (0.04)	0.60 (0.03)	0.74 (0.03)	0.91 (0.04)	1.09 (0.07)	1.22 (0.10)
Female	9-13	265	0.65 (0.04)	0.42 (0.04)	0.45 (0.04)	0.52 (0.05)	0.60 (0.06)	0.70 (0.07)	0.80 (0.09)	0.86 (0.10)
	14-18	290	0.67 (0.04)	0.44 (0.04)	0.48 (0.04)	0.55 (0.05)	0.65 (0.06)	0.77 (0.06)	0.89 (0.08)	0.96 (0.09)
	19-30	197	0.68 (0.05)	0.42 (0.04)	0.46 (0.04)	0.54 (0.05)	0.65 (0.06)	0.79 (0.07)	0.93 (0.09)	1.02 (0.11)
	31-50	312	0.84 (0.05)	0.57 (0.04)	0.62 (0.04)	0.71 (0.05)	0.83 (0.06)	0.97 (0.07)	1.11 (0.08)	1.21 (0.10)
	51-70	312	0.90 (0.06)	0.47 (0.10) ^E	0.53 (0.10) ^E	0.65 (0.08)	0.81 (0.07)	1.02 (0.09)	1.24 (0.16)	1.40 (0.22)
	>70	239	0.84 (0.09)	0.39 (0.06)	0.45 (0.06)	0.55 (0.06)	0.72 (0.07)	0.95 (0.10)	1.24 (0.18)	1.47 (0.25) ^E
	19+	1060	0.82 (0.03)	0.45 (0.05)	0.51 (0.05)	0.62 (0.04)	0.78 (0.04)	0.97 (0.05)	1.18 (0.09)	1.33 (0.12)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.8 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Saskatchewan, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	0.64 (0.07)	0.44 (0.08) ^E	0.48 (0.08)	0.55 (0.08)	0.65 (0.09)	0.76 (0.12)	0.90 (0.17) ^E	0.99 (0.21) ^E
	4-8	213	0.61 (0.03)	0.47 (0.02)	0.49 (0.02)	0.54 (0.02)	0.59 (0.03)	0.65 (0.03)	0.72 (0.04)	0.76 (0.04)
Male	9-13	122	0.75 (0.10)	0.41 (0.08) ^E	0.46 (0.08) ^E	0.56 (0.08)	0.70 (0.09)	0.88 (0.14)	1.09 (0.21) ^E	1.24 (0.27) ^E
	14-18	150	0.67 (0.04)	0.48 (0.04)	0.51 (0.04)	0.58 (0.05)	0.66 (0.06)	0.76 (0.07)	0.86 (0.09)	0.93 (0.11)
	19-30	106	0.88 (0.11)	0.54 (0.07)	0.59 (0.08)	0.70 (0.10)	0.83 (0.12)	1.00 (0.15)	1.19 (0.19)	1.32 (0.22) ^E
	31-50	155	0.77 (0.07)	0.56 (0.11) ^E	0.59 (0.10)	0.65 (0.08)	0.73 (0.07)	0.82 (0.09)	0.92 (0.14)	0.98 (0.19) ^E
	51-70	122	0.89 (0.08)	0.62 (0.07)	0.68 (0.08)	0.80 (0.10)	0.95 (0.12)	1.11 (0.14)	1.27 (0.17)	1.37 (0.19)
	>70	88	1.05 (0.13)	0.49 (0.12) ^E	0.56 (0.12) ^E	0.69 (0.12) ^E	0.90 (0.14)	1.20 (0.20)	1.57 (0.31) ^E	1.85 (0.41) ^E
	19+	471	0.86 (0.04)	0.54 (0.10) ^E	0.60 (0.09)	0.71 (0.07)	0.84 (0.06)	1.01 (0.07)	1.19 (0.11)	1.30 (0.15)
Female	9-13	103	0.70 (0.09)	0.41 (0.09) ^E	0.46 (0.09) ^E	0.55 (0.08)	0.68 (0.09)	0.86 (0.13)	1.09 (0.22) ^E	1.27 (0.32) ^E
	14-18	142	0.72 (0.06)	0.49 (0.05)	0.53 (0.05)	0.61 (0.06)	0.72 (0.07)	0.85 (0.09)	0.98 (0.12)	1.07 (0.14)
	19-30	111	0.80 (0.06)	0.65 (0.12) ^E	0.69 (0.12) ^E	0.76 (0.10)	0.85 (0.10)	0.95 (0.11)	1.05 (0.14)	1.12 (0.18)
	31-50	146	0.73 (0.06)	0.46 (0.06)	0.50 (0.06)	0.58 (0.06)	0.68 (0.07)	0.80 (0.08)	0.93 (0.09)	1.01 (0.11)
	51-70	184	0.99 (0.09)	0.51 (0.13) ^E	0.58 (0.12) ^E	0.71 (0.11)	0.89 (0.11)	1.15 (0.14)	1.46 (0.23)	1.70 (0.31) ^E
	>70	143	0.84 (0.06)	0.58 (0.10) ^E	0.63 (0.10)	0.72 (0.09)	0.85 (0.08)	1.01 (0.10)	1.18 (0.17)	1.29 (0.23) ^E
	19+	584	0.83 (0.04)	0.51 (0.08)	0.56 (0.08)	0.66 (0.07)	0.80 (0.05)	0.97 (0.06)	1.17 (0.10)	1.31 (0.15)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.9 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Alberta, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	169	0.50 (0.03)	0.43 (0.03)	0.44 (0.03)	0.46 (0.03)	0.48 (0.03)	0.50 (0.03)	0.52 (0.04)	0.53 (0.04)
	4-8	281	0.57 (0.03)	0.42 (0.03)	0.45 (0.03)	0.50 (0.03)	0.55 (0.03)	0.62 (0.04)	0.68 (0.04)	0.72 (0.05)
Male	9-13	183	0.61 (0.03)	0.38 (0.07) ^E	0.42 (0.06)	0.50 (0.05)	0.60 (0.03)	0.72 (0.05)	0.84 (0.09)	0.92 (0.12)
	14-18	187	0.66 (0.04)	0.49 (0.03)	0.52 (0.04)	0.58 (0.04)	0.66 (0.05)	0.75 (0.05)	0.85 (0.06)	0.92 (0.07)
	19-30	223	0.78 (0.08)	0.38 (0.09) ^E	0.44 (0.09) ^E	0.55 (0.08)	0.72 (0.07)	0.95 (0.13)	1.21 (0.26) ^E	1.40 (0.39) ^E
	31-50	229	0.78 (0.08)	0.50 (0.05)	0.55 (0.06)	0.64 (0.07)	0.75 (0.08)	0.89 (0.09)	1.03 (0.11)	1.12 (0.12)
	51-70	197	0.69 (0.05)	0.41 (0.05)	0.46 (0.05)	0.55 (0.06)	0.67 (0.06)	0.82 (0.07)	0.98 (0.09)	1.08 (0.10)
	>70	72	0.78 (0.07)	0.51 (0.11) ^E	0.56 (0.10) ^E	0.67 (0.10)	0.81 (0.09)	0.96 (0.11)	1.12 (0.16)	1.23 (0.21) ^E
	19+	721	0.76 (0.04)	0.57 (0.09)	0.61 (0.08)	0.68 (0.07)	0.77 (0.05)	0.88 (0.06)	0.99 (0.11)	1.07 (0.15)
Female	9-13	165	0.54 (0.03)	0.44 (0.08) ^E	0.46 (0.07)	0.50 (0.06)	0.55 (0.04)	0.60 (0.05)	0.65 (0.08)	0.68 (0.11)
	14-18	206	0.71 (0.05)	0.39 (0.08) ^E	0.43 (0.07)	0.53 (0.06)	0.65 (0.05)	0.81 (0.07)	1.00 (0.11)	1.12 (0.16)
	19-30	191	0.67 (0.05)	0.33 (0.07) ^E	0.38 (0.07) ^E	0.47 (0.06)	0.61 (0.05)	0.79 (0.07)	0.99 (0.12)	1.13 (0.15)
	31-50	258	0.81 (0.04)	0.54 (0.03)	0.58 (0.04)	0.67 (0.04)	0.77 (0.05)	0.89 (0.06)	1.02 (0.07)	1.10 (0.07)
	51-70	249	0.83 (0.05)	0.37 (0.07) ^E	0.44 (0.06)	0.58 (0.06)	0.78 (0.06)	1.05 (0.08)	1.36 (0.14)	1.59 (0.20)
	>70	128	0.74 (0.07)	0.41 (0.08) ^E	0.46 (0.08) ^E	0.57 (0.09)	0.71 (0.09)	0.90 (0.11)	1.11 (0.15)	1.25 (0.18)
	19+	826	0.78 (0.02)	0.41 (0.03)	0.46 (0.03)	0.57 (0.03)	0.73 (0.03)	0.93 (0.04)	1.15 (0.06)	1.29 (0.08)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.10 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, British Columbia, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	192	0.59 (0.05)	0.31 (0.05) ^E	0.35 (0.05)	0.42 (0.05)	0.52 (0.05)	0.65 (0.08)	0.82 (0.12)	0.95 (0.17) ^E
	4-8	321	0.63 (0.04)	0.44 (0.03)	0.47 (0.03)	0.53 (0.03)	0.61 (0.04)	0.70 (0.05)	0.81 (0.07)	0.87 (0.08)
Male	9-13	226	0.68 (0.06)	0.39 (0.05)	0.42 (0.05)	0.50 (0.05)	0.61 (0.04)	0.74 (0.06)	0.88 (0.10)	0.98 (0.14)
	14-18	262	0.65 (0.03)	0.43 (0.04)	0.47 (0.04)	0.53 (0.05)	0.61 (0.05)	0.70 (0.06)	0.79 (0.06)	0.85 (0.07)
	19-30	197	0.79 (0.05)	0.55 (0.09)	0.61 (0.08)	0.70 (0.07)	0.81 (0.07)	0.94 (0.08)	1.06 (0.11)	1.13 (0.14)
	31-50	282	0.71 (0.04)	0.42 (0.03)	0.46 (0.04)	0.54 (0.04)	0.66 (0.05)	0.79 (0.05)	0.93 (0.06)	1.02 (0.07)
	51-70	234	0.80 (0.04)	0.46 (0.06)	0.52 (0.06)	0.62 (0.05)	0.76 (0.05)	0.93 (0.07)	1.12 (0.10)	1.24 (0.13)
	>70	119	0.81 (0.05)	0.52 (0.05)	0.59 (0.06)	0.70 (0.06)	0.84 (0.07)	0.98 (0.08)	1.12 (0.09)	1.22 (0.11)
	19+	832	0.76 (0.02)	0.43 (0.04)	0.48 (0.04)	0.59 (0.03)	0.73 (0.03)	0.91 (0.04)	1.08 (0.06)	1.20 (0.08)
Female	9-13	226	0.62 (0.04)	0.47 (0.04)	0.50 (0.04)	0.55 (0.04)	0.61 (0.05)	0.69 (0.05)	0.76 (0.06)	0.81 (0.06)
	14-18	242	0.68 (0.04)	0.50 (0.09) ^E	0.53 (0.08)	0.60 (0.07)	0.68 (0.06)	0.77 (0.06)	0.85 (0.10)	0.91 (0.12)
	19-30	208	0.79 (0.05)	0.43 (0.06)	0.48 (0.06)	0.59 (0.05)	0.74 (0.06)	0.92 (0.08)	1.13 (0.12)	1.28 (0.16)
	31-50	263	0.89 (0.07)	0.58 (0.11) ^E	0.64 (0.10)	0.75 (0.09)	0.89 (0.08)	1.07 (0.10)	1.25 (0.16)	1.38 (0.21)
	51-70	322	0.85 (0.05)	0.40 (0.09) ^E	0.47 (0.09) ^E	0.61 (0.07)	0.80 (0.06)	1.04 (0.08)	1.30 (0.13)	1.49 (0.18)
	>70	198	0.81 (0.05)	0.52 (0.05)	0.57 (0.06)	0.67 (0.06)	0.80 (0.07)	0.95 (0.09)	1.12 (0.11)	1.24 (0.13)
	19+	991	0.85 (0.03)	0.48 (0.04)	0.54 (0.04)	0.66 (0.04)	0.82 (0.04)	1.02 (0.05)	1.26 (0.08)	1.42 (0.10)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.11 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Atlantic Region, 2004^{1–3}

Sex	Age (years)	Percentiles (and <i>SE</i>) of usual intake								
		n	Mean (<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)	95th (<i>SE</i>)
Both	1-3	348	0.53 (0.04)	0.34 (0.04)	0.37 (0.04)	0.42 (0.04)	0.49 (0.04)	0.60 (0.06)	0.73 (0.12)	0.83 (0.17) ^E
	4-8	554	0.59 (0.03)	0.41 (0.04)	0.44 (0.04)	0.49 (0.03)	0.57 (0.03)	0.67 (0.04)	0.77 (0.06)	0.84 (0.09)
Male	9-13	409	0.74 (0.11)	0.43 (0.04)	0.48 (0.05)	0.57 (0.06)	0.69 (0.09)	0.86 (0.12)	1.05 (0.18) ^E	1.19 (0.22) ^E
	14-18	414	0.63 (0.03)	0.39 (0.04)	0.43 (0.04)	0.51 (0.04)	0.61 (0.04)	0.72 (0.04)	0.83 (0.05)	0.91 (0.06)
	19-30	311	0.78 (0.11)	0.45 (0.07)	0.50 (0.06)	0.56 (0.06)	0.67 (0.07)	0.87 (0.14)	1.18 (0.33) ^E	^F
	31-50	489	0.69 (0.03)	0.41 (0.04)	0.45 (0.04)	0.54 (0.03)	0.65 (0.03)	0.78 (0.04)	0.93 (0.07)	1.04 (0.09)
	51-70	575	0.77 (0.04)	0.44 (0.05)	0.49 (0.05)	0.59 (0.04)	0.72 (0.04)	0.90 (0.06)	1.11 (0.09)	1.25 (0.12)
	>70	239	0.77 (0.03)	0.47 (0.05)	0.53 (0.04)	0.63 (0.04)	0.77 (0.04)	0.93 (0.05)	1.09 (0.06)	1.20 (0.08)
	19+	1614	0.74 (0.03)	0.42 (0.02)	0.47 (0.02)	0.56 (0.02)	0.69 (0.02)	0.86 (0.04)	1.07 (0.07)	1.22 (0.11)
Female	9-13	355	0.58 (0.02)	0.44 (0.05)	0.46 (0.04)	0.50 (0.03)	0.55 (0.03)	0.61 (0.04)	0.67 (0.06)	0.71 (0.08)
	14-18	410	0.67 (0.08)	0.40 (0.07)	0.43 (0.06)	0.50 (0.06)	0.60 (0.06)	0.72 (0.07)	0.87 (0.11)	0.98 (0.14)
	19-30	384	0.76 (0.05)	0.61 (0.08)	0.64 (0.07)	0.69 (0.06)	0.76 (0.06)	0.82 (0.08)	0.89 (0.11)	0.93 (0.13)
	31-50	585	0.82 (0.04)	0.53 (0.07)	0.57 (0.06)	0.66 (0.05)	0.78 (0.05)	0.92 (0.06)	1.07 (0.10)	1.17 (0.14)
	51-70	711	0.82 (0.03)	0.60 (0.08)	0.64 (0.07)	0.72 (0.05)	0.81 (0.04)	0.92 (0.05)	1.03 (0.08)	1.11 (0.11)
	>70	342	0.76 (0.04)	0.50 (0.07)	0.55 (0.07)	0.64 (0.06)	0.76 (0.05)	0.90 (0.06)	1.05 (0.08)	1.15 (0.11)
	19+	2022	0.80 (0.02)	0.56 (0.04)	0.60 (0.03)	0.68 (0.03)	0.79 (0.02)	0.92 (0.03)	1.04 (0.05)	1.13 (0.07)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.12 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Prairie Region, 2004^{1–3}

Sex	Age (years)	Percentiles (and <i>SE</i>) of usual intake								
		n	Mean (<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)	95th (<i>SE</i>)
Both	1-3	622	0.52 (0.02)	0.43 (0.06)	0.45 (0.05)	0.47 (0.04)	0.50 (0.03)	0.54 (0.03)	0.57 (0.05)	0.59 (0.07)
	4-8	919	0.58 (0.02)	0.43 (0.05)	0.45 (0.04)	0.50 (0.03)	0.57 (0.03)	0.64 (0.03)	0.71 (0.05)	0.76 (0.07)
Male	9-13	579	0.63 (0.03)	0.38 (0.06)	0.42 (0.05)	0.50 (0.04)	0.61 (0.03)	0.75 (0.05)	0.89 (0.08)	0.99 (0.12)
	14-18	634	0.66 (0.02)	0.48 (0.06)	0.51 (0.06)	0.58 (0.04)	0.66 (0.03)	0.75 (0.04)	0.85 (0.07)	0.91 (0.09)
	19-30	578	0.81 (0.06)	0.47 (0.09) ^E	0.53 (0.08)	0.64 (0.07)	0.79 (0.06)	0.97 (0.08)	1.18 (0.13)	1.33 (0.18)
	31-50	693	0.77 (0.06)	0.50 (0.09) ^E	0.55 (0.08)	0.64 (0.07)	0.75 (0.06)	0.88 (0.07)	1.01 (0.10)	1.10 (0.14)
	51-70	596	0.75 (0.04)	0.45 (0.03)	0.51 (0.03)	0.60 (0.04)	0.73 (0.04)	0.90 (0.05)	1.08 (0.07)	1.20 (0.08)
	>70	296	0.83 (0.05)	0.45 (0.05)	0.51 (0.05)	0.62 (0.05)	0.77 (0.06)	0.98 (0.08)	1.23 (0.11)	1.42 (0.15)
	19+	2163	0.78 (0.03)	0.48 (0.04)	0.53 (0.04)	0.63 (0.04)	0.77 (0.03)	0.94 (0.04)	1.14 (0.06)	1.27 (0.08)
Female	9-13	533	0.59 (0.03)	0.39 (0.06)	0.42 (0.05)	0.48 (0.04)	0.57 (0.03)	0.68 (0.04)	0.81 (0.08)	0.90 (0.12)
	14-18	638	0.70 (0.03)	0.44 (0.05)	0.48 (0.05)	0.56 (0.04)	0.67 (0.04)	0.80 (0.05)	0.94 (0.08)	1.03 (0.10)
	19-30	499	0.69 (0.03)	0.42 (0.06)	0.46 (0.06)	0.55 (0.05)	0.66 (0.04)	0.80 (0.05)	0.95 (0.08)	1.05 (0.11)
	31-50	716	0.80 (0.03)	0.52 (0.08)	0.57 (0.07)	0.65 (0.06)	0.77 (0.04)	0.90 (0.05)	1.04 (0.09)	1.13 (0.12)
	51-70	745	0.88 (0.03)	0.41 (0.05)	0.48 (0.05)	0.61 (0.04)	0.80 (0.04)	1.06 (0.06)	1.37 (0.10)	1.60 (0.14)
	>70	510	0.79 (0.04)	0.45 (0.05)	0.50 (0.05)	0.60 (0.05)	0.75 (0.05)	0.95 (0.07)	1.18 (0.10)	1.34 (0.14)
	19+	2470	0.80 (0.02)	0.44 (0.03)	0.49 (0.02)	0.60 (0.02)	0.75 (0.02)	0.94 (0.03)	1.16 (0.04)	1.31 (0.06)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.13 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Canada excluding territories, 2004^{1–3}

Sex	Age (years)	Percentiles (and <i>SE</i>) of usual intake								
		n	Mean (<i>SE</i>)	5th (<i>SE</i>)	10th (<i>SE</i>)	25th (<i>SE</i>)	50th (<i>SE</i>)	75th (<i>SE</i>)	90th (<i>SE</i>)	95th (<i>SE</i>)
Both	1-3	2117	0.52 (0.01)	0.34 (0.02)	0.37 (0.02)	0.42 (0.02)	0.49 (0.01)	0.58 (0.02)	0.69 (0.04)	0.78 (0.06)
	4-8	3235	0.60 (0.01)	0.43 (0.03)	0.46 (0.03)	0.51 (0.02)	0.58 (0.01)	0.66 (0.02)	0.75 (0.04)	0.81 (0.05)
Male	9-13	2080	0.64 (0.02)	0.41 (0.02)	0.45 (0.02)	0.52 (0.02)	0.62 (0.02)	0.74 (0.03)	0.87 (0.04)	0.97 (0.06)
	14-18	2288	0.68 (0.02)	0.45 (0.03)	0.49 (0.03)	0.56 (0.02)	0.65 (0.02)	0.76 (0.03)	0.88 (0.05)	0.96 (0.07)
	19-30	1804	0.79 (0.02)	0.48 (0.04)	0.53 (0.03)	0.64 (0.03)	0.78 (0.03)	0.97 (0.05)	1.18 (0.07)	1.32 (0.10)
	31-50	2596	0.72 (0.02)	0.42 (0.03)	0.47 (0.03)	0.56 (0.02)	0.68 (0.02)	0.85 (0.04)	1.05 (0.06)	1.18 (0.08)
	51-70	2550	0.83 (0.02)	0.48 (0.03)	0.54 (0.03)	0.65 (0.02)	0.79 (0.02)	0.98 (0.03)	1.18 (0.05)	1.33 (0.07)
	>70	1520	0.78 (0.02)	0.45 (0.02)	0.51 (0.02)	0.61 (0.02)	0.75 (0.03)	0.93 (0.03)	1.13 (0.05)	1.27 (0.06)
	19+	8470	0.77 (0.01)	0.44 (0.01)	0.49 (0.01)	0.60 (0.01)	0.74 (0.02)	0.92 (0.02)	1.13 (0.03)	1.29 (0.05)
Female	9-13	1980	0.62 (0.01)	0.46 (0.04)	0.49 (0.03)	0.54 (0.02)	0.61 (0.02)	0.69 (0.02)	0.76 (0.04)	0.82 (0.06)
	14-18	2256	0.74 (0.02)	0.48 (0.04)	0.52 (0.04)	0.60 (0.03)	0.71 (0.03)	0.84 (0.04)	0.99 (0.06)	1.09 (0.08)
	19-30	1854	0.74 (0.02)	0.46 (0.03)	0.50 (0.03)	0.58 (0.03)	0.69 (0.03)	0.83 (0.03)	0.98 (0.06)	1.08 (0.07)
	31-50	2686	0.84 (0.02)	0.53 (0.04)	0.58 (0.04)	0.68 (0.03)	0.81 (0.03)	0.97 (0.03)	1.15 (0.06)	1.27 (0.08)
	51-70	3200	0.85 (0.02)	0.52 (0.03)	0.57 (0.03)	0.69 (0.03)	0.84 (0.02)	1.02 (0.03)	1.21 (0.05)	1.35 (0.06)
	>70	2610	0.79 (0.02)	0.51 (0.03)	0.56 (0.03)	0.65 (0.03)	0.78 (0.03)	0.93 (0.03)	1.09 (0.05)	1.20 (0.07)
	19+	10350	0.82 (0.01)	0.50 (0.02)	0.55 (0.02)	0.65 (0.02)	0.79 (0.01)	0.96 (0.02)	1.14 (0.03)	1.26 (0.03)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

32. Moisture (g/d): Usual intakes from food

Table 32.1 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake									AI ³	% >AI (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	79	1319 (66)	872 (96)	956 (90)	1108 (80)	1295 (77)	1502 (99)	1706 (139)	1837 (168)	1300	49.4 (12.3) ^E
	4-8	127	1513 (70)	880 (90)	997 (86)	1213 (80)	1482 (89)	1781 (115)	2076 (147)	2265 (173)	1700	30.9 (8.6) ^E
Male	9-13	111	2049 (154)	1184 (135)	1304 (133)	1539 (134)	1876 (164)	2349 (259)	2933 (389)	3330 (475)	2400	F
	14-18	107	2704 (159)	1707 (265)	1913 (243)	2302 (208)	2805 (194)	3391 (259)	3998 (397)	4401 (511)	3300	F
	19-30	77	3153 (220)	2662 (460) ^E	2796 (421)	3033 (356)	3307 (301)	3587 (295)	3838 (347)	3985 (401)	3700	F
	31-50	145	2554 (112)	1676 (204)	1830 (184)	2109 (154)	2457 (142)	2860 (173)	3285 (246)	3574 (312)	3700	F
	51-70	182	2548 (134)	1262 (142)	1420 (143)	1749 (142)	2242 (144)	2930 (192)	3783 (323)	4429 (445)	3700	F
	>70	63	2453 (196)	1500 (197)	1659 (199)	1972 (213)	2401 (255)	2941 (334)	3545 (443)	3970 (534)	3700	F
	19+	467	2667 (80)	1590 (93)	1769 (87)	2091 (87)	2552 (99)	3158 (134)	3803 (191)	4228 (234)	3700	11.7 (3.2) ^E
Female	9-13	96	1595 (102)	1006 (123)	1111 (113)	1292 (102)	1516 (109)	1796 (160)	2137 (256)	2401 (338)	2100	F
	14-18	105	1892 (125)	1345 (159)	1455 (147)	1664 (130)	1921 (136)	2215 (197)	2533 (283)	2753 (343)	2300	F
	19-30	91	2255 (118)	1377 (115)	1543 (111)	1842 (116)	2213 (139)	2654 (174)	3149 (222)	3503 (262)	2700	23.1 (7.2) ^E
	31-50	167	2197 (117)	1361 (133)	1510 (135)	1792 (138)	2155 (142)	2580 (180)	3021 (242)	3315 (285)	2700	F
	51-70	198	2249 (152)	1432 (216)	1580 (194)	1845 (163)	2178 (148)	2575 (179)	3020 (280)	3343 (383)	2700	F
	>70	74	1837 (109)	1062 (117)	1188 (118)	1417 (118)	1704 (130)	2037 (145)	2389 (152)	2628 (159)	2700	F
	19+	530	2183 (73)	1350 (66)	1502 (65)	1779 (68)	2127 (78)	2545 (105)	3010 (158)	3341 (209)	2700	18.6 (4.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.2 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	58	1414 (100)	1022 (103)	1100 (97)	1236 (94)	1404 (107)	1601 (140)	1820 (191)	1975 (235)	1300	65.8 (14.8) ^E
	4-8	110	1460 (85)	917 (97)	1013 (96)	1190 (97)	1410 (101)	1658 (110)	1905 (131)	2065 (153)	1700	F
Male	9-13	95	1974 (121)	1135 (154)	1277 (156)	1565 (159)	1966 (165)	2450 (223)	2945 (272)	3262 (317)	2400	27.1 (9.0) ^E
	14-18	87	2444 (156)	1483 (149)	1636 (161)	1963 (188)	2447 (229)	3061 (285)	3705 (351)	4121 (403)	3300	F
	19-30	70	3040 (180)	1993 (330)	2205 (297)	2595 (242)	3051 (220)	3543 (273)	4023 (374)	4308 (438)	3700	F
	31-50	109	2638 (155)	1665 (284) ^E	1807 (276)	2119 (245)	2572 (222)	3086 (272)	3648 (390)	4055 (503)	3700	F
	51-70	128	2624 (167)	1926 (235)	2033 (219)	2244 (191)	2525 (175)	2858 (213)	3207 (319)	3440 (417)	3700	F
	>70	65	2251 (116)	1448 (127)	1612 (120)	1909 (117)	2284 (137)	2728 (190)	3216 (260)	3557 (319)	3700	F
	19+	372	2680 (93)	1668 (101)	1825 (105)	2155 (109)	2626 (118)	3181 (151)	3768 (210)	4172 (260)	3700	11.2 (3.7) ^E
Female	9-13	75	1781 (115)	1347 (184)	1446 (171)	1626 (156)	1847 (166)	2094 (206)	2339 (264)	2498 (310)	2100	F
	14-18	81	2072 (116)	1132 (173)	1283 (160)	1568 (141)	1938 (139)	2369 (175)	2815 (248)	3110 (310)	2300	28.3 (9.3) ^E
	19-30	101	2868 (208)	1367 (210)	1647 (205)	2158 (202)	2799 (224)	3563 (313)	4417 (470)	5024 (593)	2700	53.9 (9.3) ^E
	31-50	116	2698 (153)	1599 (197)	1781 (196)	2140 (195)	2639 (207)	3275 (246)	3995 (316)	4506 (382)	2700	47.1 (10.4) ^E
	51-70	146	2619 (105)	1557 (137)	1744 (128)	2074 (117)	2503 (123)	3041 (160)	3636 (209)	4040 (246)	2700	39.5 (7.1) ^E
	>70	94	2190 (120)	1618 (217)	1742 (205)	1975 (187)	2269 (183)	2594 (208)	2906 (257)	3099 (295)	2700	F
	19+	457	2643 (85)	1497 (80)	1706 (81)	2091 (87)	2583 (101)	3184 (133)	3866 (179)	4358 (217)	2700	44.3 (5.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.3 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	112	1300 (47)	937 (80)	1002 (74)	1125 (64)	1278 (57)	1449 (67)	1627 (99)	1751 (131)	1300	46.4 (10.3) ^E
	4-8	177	1492 (65)	906 (126)	1021 (115)	1232 (98)	1497 (85)	1795 (90)	2094 (120)	2287 (147)	1700	32.0 (7.4) ^E
Male	9-13	111	2013 (89)	1298 (151)	1435 (140)	1685 (124)	1992 (126)	2333 (160)	2669 (222)	2885 (274)	2400	F
	14-18	113	2384 (128)	1490 (144)	1654 (136)	1940 (133)	2310 (155)	2787 (201)	3333 (274)	3714 (341)	3300	F
	19-30	91	3601 (294)	2418 (344)	2632 (315)	3030 (276)	3578 (291)	4301 (448)	5146 (737)	5753 (982) ^E	3700	F
	31-50	101	3118 (171)	1926 (251)	2161 (234)	2589 (210)	3119 (200)	3708 (235)	4290 (322)	4663 (407)	3700	F
	51-70	134	2726 (169)	1650 (150)	1834 (145)	2166 (143)	2586 (158)	3091 (194)	3657 (261)	4064 (324)	3700	F
	>70	56	2376 (218)	1864 (272)	1979 (246)	2184 (210)	2440 (201)	2737 (259)	3052 (381)	3268 (489)	3700	F
	19+	382	3027 (104)	1828 (127)	2043 (115)	2442 (101)	2957 (105)	3588 (149)	4302 (238)	4816 (322)	3700	21.8 (4.6) ^E
Female	9-13	105	1768 (107)	1239 (196)	1340 (165)	1529 (137)	1772 (144)	2048 (167)	2322 (189)	2498 (206)	2100	F
	14-18	120	2037 (122)	1180 (145)	1366 (130)	1639 (124)	1939 (161)	2436 (222)	2997 (299)	3324 (343)	2300	30.3 (10.0) ^E
	19-30	91	2857 (203)	1806 (317) ^E	2011 (306)	2392 (280)	2860 (262)	3375 (280)	3920 (367)	4311 (467)	2700	58.9 (15.7) ^E
	31-50	159	2906 (167)	1491 (159)	1714 (156)	2147 (152)	2696 (167)	3337 (227)	4132 (379)	4808 (566)	2700	49.8 (7.9)
	51-70	174	2462 (108)	1291 (119)	1492 (105)	1858 (97)	2311 (109)	2817 (130)	3385 (175)	3817 (227)	2700	29.8 (5.4) ^E
	>70	80	2148 (92)	1351 (143)	1527 (133)	1850 (121)	2241 (133)	2571 (160)	3057 (238)	3439 (317)	2700	F
	19+	504	2669 (80)	1416 (72)	1620 (70)	2011 (74)	2531 (86)	3128 (112)	3811 (165)	4359 (247)	2700	42.0 (4.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.4 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	99	1594 (91)	1101 (161)	1192 (153)	1378 (135)	1635 (123)	1941 (156)	2248 (219)	2441 (263)	1300	81.9 (10.2)
	4-8	140	1614 (92)	1227 (161)	1308 (144)	1448 (119)	1616 (105)	1802 (130)	1987 (186)	2108 (229)	1700	F
Male	9-13	92	1793 (136)	1256 (185)	1340 (176)	1499 (163)	1712 (168)	1974 (215)	2265 (309)	2470 (395)	2400	F
	14-18	107	2686 (240)	1407 (180)	1597 (187)	1975 (208)	2534 (241)	3341 (342)	4420 (580)	5297 (804)	3300	25.9 (8.3) ^E
	19-30	73	3531 (365)	3203 (637) ^E	3307 (591) ^E	3485 (510)	3691 (429)	3906 (408)	4106 (498)	4229 (608)	3700	F
	31-50	134	3306 (222)	2260 (291)	2429 (277)	2751 (256)	3180 (252)	3700 (295)	4262 (399)	4647 (502)	3700	F
	51-70	131	2746 (169)	1481 (168)	1669 (169)	2039 (183)	2604 (210)	3321 (254)	4057 (348)	4574 (448)	3700	F
	>70	55	2205 (162)	1195 (166)	1378 (163)	1672 (168)	2067 (204)	2652 (265)	3273 (336)	3625 (370)	3700	F
	19+	393	3088 (109)	1777 (148)	1975 (152)	2408 (149)	3047 (143)	3795 (163)	4602 (220)	5185 (283)	3700	27.6 (4.5)
Female	9-13	79	1835 (93)	1170 (140)	1301 (124)	1525 (109)	1796 (117)	2154 (152)	2636 (225)	3029 (314)	2100	27.8 (8.8) ^E
	14-18	104	2201 (150)	1315 (134)	1488 (143)	1820 (164)	2243 (180)	2711 (214)	3194 (266)	3528 (316)	2300	46.6 (11.4) ^E
	19-30	101	2490 (150)	1328 (164)	1507 (177)	1897 (188)	2419 (192)	2979 (195)	3568 (250)	4015 (349)	2700	36.5 (9.0) ^E
	31-50	143	2914 (138)	1645 (254)	1880 (219)	2305 (163)	2789 (142)	3299 (193)	3915 (350)	4441 (516)	2700	54.9 (9.0)
	51-70	193	2647 (137)	1517 (136)	1683 (133)	2015 (126)	2484 (134)	3097 (233)	3808 (341)	4322 (423)	2700	39.8 (7.4) ^E
	>70	94	2010 (105)	1207 (102)	1331 (99)	1552 (108)	1857 (133)	2229 (162)	2610 (198)	2869 (232)	2700	F
	19+	531	2637 (78)	1397 (70)	1594 (71)	1985 (75)	2513 (87)	3152 (114)	3884 (179)	4435 (253)	2700	41.6 (4.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.5 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	311	1407 (47)	826 (78)	949 (74)	1156 (68)	1395 (63)	1657 (65)	1934 (84)	2124 (106)	1300	60.4 (7.1)
	4-8	485	1610 (59)	987 (50)	1098 (50)	1308 (52)	1578 (61)	1896 (85)	2244 (126)	2494 (162)	1700	39.3 (5.9)
Male	9-13	277	2163 (92)	1330 (139)	1503 (127)	1805 (115)	2177 (118)	2623 (142)	3124 (195)	3484 (248)	2400	36.1 (7.3) ^E
	14-18	339	2881 (150)	1636 (138)	1826 (151)	2258 (168)	2917 (187)	3703 (231)	4574 (316)	5204 (394)	3300	36.6 (6.9) ^E
	19-30	237	3489 (173)	2067 (234)	2345 (222)	2864 (198)	3526 (199)	4286 (251)	5058 (348)	5563 (430)	3700	43.6 (7.9) ^E
	31-50	423	2981 (81)	1952 (117)	2138 (106)	2475 (96)	2898 (99)	3400 (123)	3949 (175)	4335 (225)	3700	15.4 (4.0) ^E
	51-70	387	2805 (81)	1715 (114)	1938 (109)	2334 (103)	2744 (104)	3227 (137)	3782 (182)	4107 (207)	3700	11.6 (3.8) ^E
	>70	132	2325 (164)	1461 (227)	1635 (210)	1955 (188)	2359 (188)	2818 (226)	3281 (283)	3582 (325)	3700	F
	19+	1179	2985 (57)	1750 (74)	1984 (69)	2389 (65)	2898 (70)	3550 (87)	4234 (110)	4681 (135)	3700	20.8 (2.4)
Female	9-13	281	1811 (111)	1103 (88)	1205 (80)	1395 (79)	1677 (93)	2051 (141)	2477 (230)	2798 (319)	2100	F
	14-18	321	2154 (108)	1240 (117)	1403 (115)	1704 (119)	2092 (132)	2567 (161)	3104 (214)	3490 (267)	2300	37.7 (8.1) ^E
	19-30	249	2631 (97)	1733 (99)	1900 (100)	2212 (108)	2620 (127)	3106 (159)	3636 (208)	3996 (245)	2700	45.2 (7.9) ^E
	31-50	364	2794 (104)	1698 (106)	1887 (101)	2241 (98)	2700 (107)	3236 (139)	3856 (205)	4332 (277)	2700	50.0 (6.2)
	51-70	467	2511 (73)	1455 (83)	1649 (78)	2003 (76)	2451 (84)	2988 (106)	3589 (153)	4017 (198)	2700	37.0 (4.5)
	>70	215	2006 (91)	1321 (113)	1437 (110)	1661 (111)	1958 (115)	2321 (139)	2716 (199)	2987 (257)	2700	F
	19+	1295	2581 (51)	1551 (46)	1727 (47)	2071 (51)	2537 (61)	3084 (77)	3695 (113)	4143 (150)	2700	41.5 (3.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.6 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						AI ³	% >AI	(SE)	
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)				95th (SE)
Both	1-3	644	1418	(33)	851 (68)	962 (58)	1158 (46)	1395 (38)	1670 (46)	1971 (75)	2181 (103)	1300	60.3	(4.5)
	4-8	956	1588	(30)	1007 (52)	1112 (46)	1302 (39)	1549 (36)	1857 (46)	2190 (73)	2412 (97)	1700	36.4	(3.2)
Male	9-13	589	1984	(54)	1350 (132)	1461 (116)	1668 (86)	1935 (59)	2248 (86)	2574 (158)	2792 (216)	2400	F	
	14-18	639	2592	(65)	1527 (137)	1734 (120)	2111 (94)	2580 (80)	3126 (100)	3708 (164)	4107 (225)	3300	19.3	(3.7) ^E
	19-30	481	3035	(89)	1834 (193)	2059 (165)	2463 (122)	2965 (97)	3554 (138)	4191 (241)	4636 (329)	3700	20.5	(5.0) ^E
	31-50	709	2961	(74)	1764 (184)	1962 (177)	2387 (134)	2947 (88)	3500 (115)	4090 (206)	4535 (285)	3700	18.4	(4.2) ^E
	51-70	758	2815	(64)	1764 (108)	1948 (97)	2284 (83)	2729 (73)	3251 (96)	3787 (152)	4160 (210)	3700	11.7	(3.0) ^E
	>70	734	2198	(61)	1328 (64)	1487 (58)	1776 (51)	2122 (53)	2552 (74)	3060 (127)	3433 (195)	3700	F	
	19+	2682	2870	(43)	1646 (63)	1864 (56)	2263 (50)	2803 (51)	3410 (58)	4041 (96)	4529 (137)	3700	16.4	(1.8)
Female	9-13	585	1739	(44)	1016 (65)	1137 (60)	1369 (51)	1674 (48)	2027 (66)	2382 (98)	2612 (124)	2100	21.1	(3.7) ^E
	14-18	645	2151	(53)	1213 (76)	1377 (72)	1698 (64)	2105 (61)	2607 (83)	3220 (144)	3687 (212)	2300	38.8	(3.5)
	19-30	514	2431	(66)	1297 (79)	1493 (76)	1861 (72)	2345 (77)	2933 (99)	3585 (142)	4043 (182)	2700	33.6	(3.6)
	31-50	758	2774	(81)	1554 (87)	1752 (82)	2125 (74)	2634 (78)	3293 (112)	4054 (182)	4603 (257)	2700	47.0	(3.8)
	51-70	955	2535	(51)	1485 (81)	1674 (72)	2016 (63)	2446 (60)	2959 (72)	3531 (109)	3938 (148)	2700	36.3	(3.3)
	>70	1345	2132	(37)	1265 (51)	1420 (46)	1698 (40)	2046 (40)	2455 (51)	2898 (74)	3206 (99)	2700	15.3	(2.0)
	19+	3572	2565	(40)	1431 (35)	1622 (33)	1976 (33)	2446 (38)	3046 (56)	3730 (91)	4219 (129)	2700	38.0	(1.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.7 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	324	1388 (46)	799 (79)	899 (75)	1093 (68)	1349 (66)	1652 (78)	1984 (116)	2222 (142)	1300	54.8 (7.2)
	4-8	425	1504 (37)	945 (111)	1039 (97)	1217 (68)	1450 (40)	1727 (69)	2022 (134)	2221 (183)	1700	27.0 (6.4) ^E
Male	9-13	274	1918 (64)	1233 (172)	1368 (141)	1602 (91)	1876 (58)	2172 (99)	2480 (180)	2695 (246)	2400	^F
	14-18	297	2595 (111)	1467 (104)	1660 (105)	2018 (114)	2498 (138)	3125 (216)	3850 (305)	4341 (364)	3300	20.2 (5.6) ^E
	19-30	249	3145 (143)	1679 (210)	1889 (199)	2315 (171)	2909 (147)	3636 (201)	4399 (329)	4902 (429)	3700	23.3 (6.1) ^E
	31-50	309	3150 (137)	1736 (167)	1997 (154)	2460 (133)	3032 (133)	3735 (222)	4571 (373)	5200 (472)	3700	26.0 (5.5) ^E
	51-70	277	2567 (81)	1524 (118)	1710 (103)	2028 (91)	2434 (91)	2944 (114)	3490 (174)	3846 (217)	3700	^F
	>70	136	2317 (121)	1256 (148)	1413 (147)	1738 (141)	2200 (143)	2758 (188)	3321 (266)	3678 (328)	3700	^F
	19+	971	2908 (72)	1581 (70)	1794 (68)	2201 (68)	2750 (73)	3411 (103)	4169 (163)	4718 (214)	3700	17.8 (2.7)
Female	9-13	265	1806 (67)	1166 (136)	1286 (116)	1494 (88)	1760 (72)	2084 (103)	2456 (191)	2735 (280)	2100	24.1 (7.1) ^E
	14-18	290	2033 (78)	1061 (87)	1192 (89)	1455 (100)	1856 (108)	2424 (137)	3096 (204)	3565 (268)	2300	29.3 (5.2) ^E
	19-30	197	2626 (137)	1461 (203)	1691 (180)	2087 (150)	2568 (149)	3126 (202)	3714 (296)	4126 (380)	2700	43.3 (8.8) ^E
	31-50	312	2717 (117)	1441 (130)	1643 (122)	2002 (119)	2509 (125)	3170 (165)	3947 (250)	4504 (326)	2700	41.5 (6.0)
	51-70	312	2536 (102)	1464 (137)	1625 (133)	1939 (127)	2375 (126)	2932 (146)	3545 (207)	3973 (272)	2700	34.0 (6.5) ^E
	>70	239	2090 (76)	1186 (92)	1331 (88)	1614 (86)	2002 (90)	2484 (108)	3019 (156)	3393 (209)	2700	17.5 (3.9) ^E
	19+	1060	2556 (64)	1379 (61)	1570 (60)	1932 (61)	2409 (69)	3013 (89)	3699 (124)	4187 (159)	2700	36.5 (3.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.8 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	129	1461 (64)	910 (76)	1010 (73)	1186 (77)	1458 (88)	1784 (124)	2073 (161)	2258 (188)	1300	64.1 (8.2)
	4-8	213	1655 (57)	1137 (122)	1240 (108)	1427 (85)	1660 (69)	1920 (87)	2179 (132)	2347 (168)	1700	45.6 (8.9) ^E
Male	9-13	122	1968 (112)	1195 (155)	1319 (157)	1586 (161)	1958 (158)	2368 (159)	2754 (196)	3002 (239)	2400	^F
	14-18	150	3075 (161)	1760 (247)	1995 (229)	2447 (197)	3052 (191)	3780 (270)	4558 (430)	5087 (564)	3300	40.3 (9.2) ^E
	19-30	106	3161 (204)	1677 (255)	1901 (245)	2334 (225)	2918 (209)	3640 (251)	4446 (370)	5015 (478)	3700	23.4 (6.9) ^E
	31-50	155	2951 (178)	1608 (227)	1802 (220)	2191 (214)	2745 (220)	3467 (269)	4304 (403)	4911 (528)	3700	^F
	51-70	122	2672 (131)	1802 (220)	1955 (195)	2240 (158)	2607 (155)	3035 (235)	3481 (377)	3779 (494)	3700	^F
	>70	88	2499 (110)	1509 (194)	1713 (171)	2082 (143)	2516 (136)	2972 (152)	3448 (217)	3789 (298)	3700	^F
	19+	471	2872 (97)	1587 (77)	1770 (84)	2161 (96)	2729 (114)	3365 (146)	4146 (224)	4774 (310)	3700	16.7 (3.5) ^E
Female	9-13	103	2017 (117)	1331 (143)	1463 (137)	1702 (131)	2005 (140)	2379 (173)	2811 (233)	3129 (290)	2100	42.6 (11.1) ^E
	14-18	142	2428 (131)	1518 (155)	1706 (155)	2047 (158)	2478 (175)	2988 (212)	3544 (276)	3932 (336)	2300	60.5 (9.6)
	19-30	111	2867 (149)	1670 (248)	1851 (235)	2207 (213)	2710 (205)	3360 (262)	4063 (386)	4517 (481)	2700	50.5 (11.6) ^E
	31-50	146	3328 (187)	2216 (357)	2444 (322)	2835 (270)	3284 (230)	3745 (238)	4168 (295)	4424 (345)	2700	81.1 (11.1)
	51-70	184	2810 (107)	1941 (214)	2113 (189)	2432 (147)	2837 (123)	3305 (172)	3785 (273)	4103 (350)	2700	58.6 (9.5)
	>70	143	2333 (89)	1494 (141)	1667 (130)	1973 (118)	2338 (119)	2729 (143)	3102 (177)	3335 (203)	2700	26.6 (7.7) ^E
	19+	584	2938 (83)	1818 (103)	2020 (102)	2401 (102)	2905 (109)	3506 (133)	4110 (169)	4481 (193)	2700	60.2 (5.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.9 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	169	1329 (48)	856 (85)	941 (76)	1104 (60)	1319 (55)	1578 (79)	1855 (126)	2043 (167)	1300	52.1 (7.1)
	4-8	281	1590 (51)	955 (94)	1063 (85)	1265 (67)	1529 (58)	1841 (81)	2168 (129)	2388 (167)	1700	35.1 (5.8)
Male	9-13	183	2216 (148)	1370 (201)	1509 (181)	1771 (150)	2132 (143)	2619 (231)	3230 (429)	3707 (611)	2400	^F
	14-18	187	2851 (127)	1935 (267)	2119 (236)	2459 (178)	2898 (138)	3423 (208)	3997 (380)	4400 (533)	3300	29.9 (9.3) ^E
	19-30	223	3189 (113)	2037 (209)	2244 (187)	2622 (147)	3094 (129)	3635 (170)	4199 (256)	4577 (330)	3700	22.7 (6.5) ^E
	31-50	229	3101 (156)	1947 (200)	2168 (191)	2582 (183)	3113 (197)	3731 (240)	4367 (310)	4789 (373)	3700	26.0 (8.1) ^E
	51-70	197	2634 (120)	1442 (166)	1658 (149)	2050 (129)	2550 (132)	3169 (192)	3890 (298)	4420 (383)	3700	^F
	>70	72	2087 (136)	1201 (217) ^E	1349 (205)	1632 (186)	2009 (170)	2463 (193)	2949 (255)	3279 (311)	3700	^F
	19+	721	2937 (83)	1754 (112)	1974 (111)	2371 (104)	2886 (105)	3524 (117)	4192 (132)	4621 (148)	3700	20.0 (3.1)
Female	9-13	165	1834 (77)	1261 (162)	1364 (147)	1561 (119)	1824 (102)	2144 (145)	2491 (240)	2730 (319)	2100	^F
	14-18	206	2145 (88)	1269 (120)	1426 (116)	1719 (108)	2095 (111)	2530 (140)	2975 (189)	3267 (235)	2300	37.1 (7.4) ^E
	19-30	191	2628 (142)	1434 (185)	1621 (186)	2019 (174)	2514 (172)	3084 (200)	3765 (288)	4264 (378)	2700	40.6 (8.9) ^E
	31-50	258	2745 (111)	1657 (205)	1861 (192)	2254 (169)	2771 (169)	3363 (212)	3943 (282)	4305 (334)	2700	53.5 (9.4) ^E
	51-70	249	2525 (96)	1643 (162)	1799 (142)	2076 (115)	2438 (104)	2888 (149)	3385 (246)	3724 (322)	2700	34.0 (7.2) ^E
	>70	128	2199 (111)	1401 (146)	1545 (144)	1832 (167)	2220 (152)	2642 (222)	3080 (264)	3382 (306)	2700	^F
	19+	826	2607 (56)	1534 (94)	1728 (90)	2096 (82)	2577 (80)	3146 (92)	3741 (120)	4138 (147)	2700	43.8 (4.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.10 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						AI ³	% >AI	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	192	1485	(55)	932 (103)	1037 (101)	1257 (85)	1538 (82)	1738 (82)	1964 (111)	2131 (151)	1300	71.7 (8.1)
	4-8	321	1516	(58)	893 (64)	992 (61)	1186 (59)	1449 (68)	1776 (96)	2137 (144)	2390 (187)	1700	29.8 (6.4) ^E
Male	9-13	226	2175	(138)	1250 (142)	1388 (141)	1661 (147)	2050 (168)	2568 (211)	3182 (304)	3634 (405)	2400	31.7 (9.1) ^E
	14-18	262	2946	(160)	1537 (137)	1788 (138)	2266 (153)	2886 (183)	3617 (234)	4452 (331)	5085 (424)	3300	34.5 (6.9) ^E
	19-30	197	3646	(210)	2005 (265)	2284 (254)	2823 (240)	3544 (250)	4413 (321)	5341 (455)	5972 (569)	3700	44.8 (9.1) ^E
	31-50	282	3616	(115)	2216 (131)	2463 (128)	2929 (126)	3534 (135)	4242 (174)	4980 (235)	5472 (283)	3700	43.3 (6.0)
	51-70	234	2935	(142)	1683 (162)	1878 (163)	2256 (167)	2765 (180)	3389 (222)	4070 (303)	4541 (376)	3700	^F
	>70	119	2653	(161)	1271 (228) ^E	1501 (214)	1943 (191)	2526 (199)	3215 (256)	3932 (354)	4408 (441)	3700	^F
	19+	832	3337	(77)	1792 (87)	2056 (87)	2562 (87)	3231 (93)	4028 (120)	4867 (162)	5430 (197)	3700	34.1 (3.3)
Female	9-13	226	2108	(126)	1220 (130)	1362 (132)	1652 (131)	2061 (131)	2578 (170)	3176 (271)	3619 (370)	2100	47.7 (8.1) ^E
	14-18	242	2285	(94)	1146 (111)	1326 (110)	1682 (109)	2171 (111)	2780 (139)	3448 (188)	3911 (230)	2300	43.8 (5.7)
	19-30	208	2825	(171)	1464 (192)	1655 (198)	2104 (197)	2760 (214)	3478 (253)	4281 (361)	4917 (492)	2700	52.3 (9.0) ^E
	31-50	263	3087	(149)	1862 (164)	2088 (156)	2455 (161)	2979 (161)	3581 (254)	4218 (365)	4768 (446)	2700	63.1 (7.9)
	51-70	322	2748	(108)	1468 (136)	1691 (135)	2122 (132)	2700 (136)	3397 (173)	4139 (249)	4642 (316)	2700	50.0 (6.2)
	>70	198	2628	(128)	1531 (166)	1734 (164)	2117 (160)	2610 (163)	3180 (187)	3766 (240)	4152 (288)	2700	45.5 (8.5) ^E
	19+	991	2884	(75)	1582 (81)	1819 (80)	2257 (79)	2816 (87)	3497 (113)	4274 (177)	4840 (238)	2700	55.2 (3.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.11 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						AI ³	% >AI	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both	1-3	348	1408	(39)	947 (47)	1031 (46)	1190 (46)	1399 (48)	1651 (63)	1921 (93)	2106 (119)	1300	61.9 (5.7)
	4-8	554	1534	(44)	899 (53)	1017 (50)	1236 (49)	1514 (53)	1835 (64)	2170 (77)	2395 (91)	1700	34.4 (4.5)
Male	9-13	409	1948	(62)	1176 (80)	1304 (78)	1553 (76)	1882 (80)	2271 (98)	2690 (138)	2983 (179)	2400	19.1 (4.5) ^E
	14-18	414	2554	(96)	1476 (80)	1664 (82)	2026 (91)	2520 (110)	3159 (149)	3921 (233)	4496 (323)	3300	21.2 (4.2) ^E
	19-30	311	3445	(171)	2412 (242)	2631 (228)	3039 (209)	3562 (211)	4169 (264)	4797 (362)	5216 (451)	3700	43.5 (10.6) ^E
	31-50	489	3023	(105)	1850 (121)	2062 (117)	2456 (114)	2961 (118)	3558 (139)	4198 (185)	4637 (228)	3700	20.7 (4.3) ^E
	51-70	575	2685	(91)	1555 (91)	1729 (88)	2064 (89)	2555 (99)	3147 (124)	3766 (171)	4218 (222)	3700	11.1 (2.8) ^E
	>70	239	2329	(110)	1416 (110)	1608 (99)	1911 (90)	2258 (107)	2763 (155)	3403 (234)	3825 (280)	3700	^F
	19+	1614	2945	(56)	1741 (59)	1943 (57)	2332 (56)	2873 (62)	3548 (82)	4284 (117)	4800 (149)	3700	20.9 (2.2)
Female	9-13	355	1754	(55)	1203 (66)	1310 (63)	1500 (61)	1744 (68)	2045 (86)	2380 (115)	2613 (142)	2100	21.7 (5.3) ^E
	14-18	410	2058	(71)	1227 (82)	1375 (78)	1656 (75)	2027 (86)	2475 (115)	2965 (161)	3306 (196)	2300	33.5 (5.4)
	19-30	384	2609	(98)	1479 (94)	1659 (104)	2043 (119)	2569 (135)	3128 (146)	3751 (182)	4240 (236)	2700	43.6 (6.5)
	31-50	585	2732	(83)	1470 (75)	1680 (78)	2090 (79)	2618 (87)	3223 (115)	3916 (183)	4457 (260)	2700	46.1 (4.2)
	51-70	711	2481	(68)	1332 (69)	1525 (60)	1869 (57)	2314 (65)	2877 (96)	3550 (156)	4066 (218)	2700	31.5 (3.4)
	>70	342	2044	(55)	1277 (76)	1418 (69)	1672 (66)	2011 (74)	2401 (96)	2783 (126)	3030 (149)	2700	12.4 (3.4) ^E
	19+	2022	2549	(42)	1390 (36)	1575 (38)	1953 (43)	2435 (47)	3013 (61)	3698 (92)	4205 (129)	2700	37.0 (2.3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.12 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						AI ³	% >AI	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	622	1365	(33)	863 (37)	949 (37)	1113 (38)	1339 (40)	1621 (51)	1923 (69)	2128 (84)	1300	54.2 (4.3)
	4-8	919	1583	(34)	995 (58)	1098 (53)	1287 (43)	1530 (39)	1824 (51)	2137 (79)	2348 (102)	1700	34.2 (3.8)
Male	9-13	579	2106	(92)	1268 (81)	1410 (77)	1678 (75)	2033 (88)	2482 (136)	3024 (232)	3442 (326)	2400	28.6 (6.3) ^E
	14-18	634	2840	(86)	1664 (108)	1883 (107)	2312 (99)	2857 (95)	3471 (134)	4199 (228)	4773 (317)	3300	30.8 (4.4)
	19-30	578	3176	(85)	1759 (117)	1995 (112)	2446 (96)	3008 (95)	3672 (124)	4410 (169)	4918 (210)	3700	24.2 (3.6)
	31-50	693	3086	(107)	1846 (121)	2072 (122)	2494 (127)	3043 (140)	3702 (155)	4416 (182)	4911 (214)	3700	25.0 (4.6) ^E
	51-70	596	2626	(78)	1527 (90)	1726 (83)	2073 (77)	2521 (86)	3102 (118)	3759 (176)	4214 (220)	3700	10.9 (2.7) ^E
	>70	296	2242	(82)	1238 (113)	1402 (113)	1726 (106)	2154 (101)	2666 (116)	3183 (141)	3531 (171)	3700	^F
	19+	2163	2919	(55)	1627 (62)	1850 (64)	2273 (62)	2811 (67)	3494 (83)	4278 (107)	4803 (126)	3700	19.9 (2.0)
Female	9-13	533	1861	(54)	1153 (70)	1278 (66)	1511 (63)	1825 (70)	2203 (94)	2618 (140)	2924 (184)	2100	30.9 (5.5) ^E
	14-18	638	2174	(62)	1227 (67)	1396 (69)	1708 (71)	2106 (78)	2583 (95)	3109 (121)	3478 (147)	2300	38.6 (4.6)
	19-30	499	2671	(96)	1448 (106)	1652 (110)	2048 (112)	2560 (113)	3173 (134)	3860 (178)	4348 (219)	2700	43.4 (5.5)
	31-50	716	2838	(81)	1623 (125)	1845 (117)	2260 (109)	2808 (115)	3449 (136)	4083 (169)	4483 (195)	2700	54.9 (5.5)
	51-70	745	2581	(65)	1670 (84)	1830 (77)	2122 (70)	2504 (73)	2976 (96)	3493 (142)	3844 (180)	2700	38.3 (4.5)
	>70	510	2202	(63)	1338 (64)	1486 (69)	1786 (79)	2195 (85)	2638 (96)	3079 (118)	3371 (137)	2700	22.2 (4.3) ^E
	19+	2470	2657	(39)	1518 (54)	1722 (51)	2099 (50)	2593 (54)	3192 (62)	3837 (80)	4273 (97)	2700	44.8 (2.7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.13 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	2117	1412 (20)	858 (25)	958 (23)	1148 (22)	1400 (24)	1680 (29)	1968 (38)	2167 (48)	1300	60.1 (2.4)
	4-8	3235	1579 (21)	956 (21)	1067 (19)	1270 (19)	1534 (23)	1858 (31)	2214 (44)	2460 (58)	1700	35.8 (2.0)
Male	9-13	2080	2067 (40)	1271 (45)	1413 (43)	1677 (41)	2018 (45)	2438 (61)	2895 (87)	3211 (109)	2400	26.8 (2.9)
	14-18	2288	2743 (52)	1531 (50)	1748 (49)	2160 (54)	2715 (62)	3397 (76)	4153 (107)	4699 (139)	3300	27.9 (2.3)
	19-30	1804	3274 (63)	1891 (80)	2137 (75)	2593 (69)	3187 (71)	3904 (88)	4681 (129)	5218 (165)	3700	31.0 (2.6)
	31-50	2596	3076 (42)	1826 (55)	2046 (54)	2468 (50)	3008 (51)	3636 (63)	4318 (85)	4786 (106)	3700	23.1 (1.9)
	51-70	2550	2790 (41)	1631 (43)	1835 (42)	2210 (43)	2687 (50)	3252 (63)	3862 (87)	4286 (109)	3700	12.9 (1.6)
	>70	1520	2311 (56)	1315 (57)	1492 (54)	1818 (54)	2246 (58)	2762 (76)	3327 (105)	3731 (132)	3700	5.3 (1.2) ^E
	19+	8470	2974 (27)	1677 (32)	1905 (29)	2327 (29)	2884 (32)	3544 (39)	4273 (53)	4782 (67)	3700	20.8 (1.1)
Female	9-13	1980	1823 (37)	1087 (33)	1210 (31)	1438 (29)	1746 (34)	2126 (48)	2540 (71)	2826 (95)	2100	26.3 (2.5)
	14-18	2256	2166 (37)	1216 (36)	1380 (37)	1701 (38)	2101 (41)	2603 (54)	3201 (78)	3616 (102)	2300	38.5 (2.4)
	19-30	1854	2581 (44)	1437 (48)	1633 (47)	2002 (48)	2499 (55)	3105 (66)	3764 (93)	4233 (124)	2700	40.6 (2.6)
	31-50	2686	2827 (46)	1607 (38)	1813 (38)	2201 (40)	2722 (48)	3364 (69)	4082 (104)	4603 (137)	2700	51.0 (2.2)
	51-70	3200	2560 (32)	1493 (39)	1677 (37)	2023 (36)	2479 (37)	3024 (46)	3636 (66)	4076 (88)	2700	38.6 (1.9)
	>70	2610	2173 (33)	1312 (37)	1453 (37)	1742 (38)	2119 (39)	2561 (48)	3045 (68)	3379 (83)	2700	19.5 (1.9)
	19+	10350	2624 (23)	1480 (20)	1678 (20)	2046 (22)	2538 (25)	3138 (33)	3816 (52)	4312 (70)	2700	42.1 (1.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

33. Naturally occurring folate (µg/d): Usual intakes from food

Table 33.1 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	79	114 (9)	79 (13)	84 (12)	93 (10)	104 (9)	115 (12)	126 (17)	133 (21)
	4-8	127	159 (12)	92 (15)	103 (15)	124 (14)	154 (14)	194 (19)	243 (32)	281 (48) ^E
Male										
	9-13	111	210 (17)	134 (19)	147 (18)	172 (17)	205 (18)	247 (25)	296 (37)	331 (47)
	14-18	107	208 (15)	100 (15)	120 (14)	158 (14)	208 (18)	274 (27)	357 (43)	423 (59)
	19-30	77	184 (14)	104 (18) ^E	122 (17)	153 (16)	188 (16)	225 (19)	262 (23)	286 (27)
	31-50	145	208 (15)	126 (22) ^E	140 (20)	166 (17)	199 (16)	239 (21)	285 (33)	318 (44)
	51-70	182	222 (14)	110 (20) ^E	127 (19)	160 (17)	202 (15)	252 (19)	310 (29)	351 (39)
	>70	63	215 (20)	139 (15)	150 (15)	170 (16)	197 (21)	233 (30)	277 (44)	310 (56) ^E
	19+	467	208 (8)	114 (10)	130 (9)	160 (8)	198 (8)	243 (10)	295 (15)	332 (19)
Female										
	9-13	96	141 (7)	100 (14)	108 (13)	122 (11)	139 (10)	158 (12)	177 (14)	189 (17)
	14-18	105	164 (15)	98 (28) ^E	111 (26) ^E	135 (23) ^E	166 (21)	200 (23)	236 (30)	260 (38)
	19-30	91	162 (17)	77 (16) ^E	88 (15) ^E	111 (15)	143 (18)	184 (26)	232 (38)	265 (47) ^E
	31-50	167	158 (12)	93 (17) ^E	104 (16)	125 (16)	154 (17)	191 (20)	233 (27)	262 (34)
	51-70	198	202 (16)	102 (22) ^E	117 (20) ^E	145 (17)	185 (15)	240 (22)	317 (49)	387 (82) ^E
	>70	74	202 (37) ^E	152 (32) ^E	162 (31) ^E	181 (31) ^E	204 (33)	233 (42) ^E	265 (57) ^E	287 (70) ^E
	19+	530	177 (8)	94 (8)	107 (8)	132 (8)	167 (10)	213 (13)	270 (20)	316 (30)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.2 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	58	133	(16)	62	(12) ^E	76	(11)	99	(12)	126	(16)	161	(23)	204	(33)	236	(44) ^E
	4-8	110	148	(13)	96	(16) ^E	106	(16)	123	(15)	145	(16)	171	(21)	199	(28)	218	(34)
	9-13	95	180	(12)	127	(19)	138	(18)	155	(17)	176	(16)	199	(18)	222	(21)	237	(25)
Male	14-18	87	229	(24)	129	(25) ^E	149	(24)	188	(22)	235	(27)	293	(41)	358	(61) ^E	404	(77) ^E
	19-30	70	241	(18)	126	(27) ^E	146	(26) ^E	188	(24)	241	(23)	295	(26)	349	(37)	389	(49)
	31-50	109	221	(14)	172	(20)	181	(18)	197	(15)	215	(16)	235	(21)	254	(29)	267	(36)
	51-70	128	224	(13)	132	(15)	148	(16)	178	(16)	219	(16)	269	(21)	323	(29)	362	(38)
	>70	65	214	(12)	157	(21)	169	(19)	191	(17)	219	(15)	250	(18)	280	(25)	298	(29)
	19+	372	225	(8)	142	(10)	158	(10)	186	(9)	222	(9)	264	(11)	307	(15)	336	(19)
	9-13	75	155	(11)	98	(16)	108	(15)	125	(14)	147	(15)	171	(18)	195	(24)	211	(29)
	14-18	81	162	(11)	80	(19) ^E	95	(17) ^E	123	(14)	156	(14)	187	(17)	219	(24)	244	(31)
Female	19-30	101	173	(12)	89	(12)	103	(12)	130	(12)	166	(14)	207	(17)	251	(24)	280	(30)
	31-50	116	186	(22)	76	(13) ^E	93	(13)	125	(15)	166	(20)	226	(32)	310	(53) ^E	374	(72) ^E
	51-70	146	195	(9)	112	(14)	127	(13)	154	(12)	187	(12)	224	(15)	263	(21)	290	(26)
	>70	94	168	(11)	93	(11)	105	(10)	128	(9)	161	(11)	202	(16)	248	(22)	281	(28)
	19+	457	184	(9)	90	(6)	106	(7)	136	(8)	175	(10)	223	(14)	278	(21)	318	(27)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.3 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	131	(10)	68 (14) ^E	79 (13)	98 (11)	124 (12)	158 (18)	194 (27)	217 (34)
	4-8	177	153	(10)	82 (12)	95 (12)	119 (11)	151 (11)	193 (16)	245 (28)	284 (40)
Male	9-13	111	201	(18)	105 (24) ^E	118 (23) ^E	145 (21)	185 (21)	239 (28)	302 (46)	348 (60) ^E
	14-18	113	205	(13)	132 (22) ^E	146 (21)	171 (20)	203 (20)	242 (21)	281 (26)	308 (31)
	19-30	91	232	(23)	117 (28) ^E	135 (26) ^E	167 (23)	207 (23)	255 (28)	310 (41)	348 (53)
	31-50	101	242	(14)	128 (21) ^E	149 (19)	186 (16)	227 (16)	273 (20)	328 (30)	368 (39)
	51-70	134	231	(22)	139 (29) ^E	156 (26)	186 (22)	219 (20)	259 (28)	310 (47)	350 (64) ^E
	>70	56	190	(16)	128 (22) ^E	138 (22)	157 (22)	180 (22)	207 (22)	234 (25)	251 (29)
	19+	382	232	(9)	120 (11)	137 (11)	171 (10)	214 (9)	263 (12)	318 (20)	361 (27)
	Female	9-13	105	168	(15)	121 (19)	131 (18)	149 (16)	172 (17)	196 (21)	218 (26)
14-18	120	155	(15)	^F	80 (23) ^E	111 (19) ^E	149 (18)	198 (22)	262 (37)	312 (53) ^E	
19-30	91	178	(16)	128 (23) ^E	137 (21)	152 (17)	169 (18)	189 (28)	213 (37) ^E	230 (44) ^E	
31-50	159	200	(10)	106 (14)	123 (14)	156 (13)	199 (14)	250 (19)	302 (26)	337 (32)	
51-70	174	211	(15)	98 (11)	116 (10)	146 (10)	186 (12)	238 (17)	314 (33)	389 (52)	
>70	80	174	(10)	109 (13)	121 (12)	146 (10)	170 (11)	200 (17)	248 (28)	286 (39)	
19+	504	196	(7)	108 (8)	123 (8)	150 (8)	187 (9)	236 (11)	292 (16)	335 (22)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.4 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	135 (9)	78 (14) ^E	89 (13)	108 (12)	134 (11)	164 (15)	196 (23)	217 (30)
	4-8	140	195 (23)	86 (20) ^E	100 (20) ^E	127 (22) ^E	179 (22)	253 (37)	349 (71) ^E	436 (101) ^E
Male	9-13	92	173 (14)	107 (25) ^E	116 (23) ^E	136 (20)	162 (17)	193 (21)	226 (31)	249 (40)
	14-18	107	216 (14)	139 (27) ^E	155 (24)	183 (20)	217 (18)	257 (23)	299 (35)	327 (46)
	19-30	73	266 (23)	139 (28) ^E	162 (28) ^E	209 (29)	269 (32)	339 (38)	416 (52)	472 (66)
	31-50	134	232 (19)	133 (25) ^E	148 (24)	177 (21)	217 (22)	271 (30)	339 (52)	392 (75) ^E
	51-70	131	251 (23)	167 (28) ^E	181 (27)	207 (24)	242 (24)	283 (31)	327 (46)	356 (59)
	>70	55	225 (19)	106 (23) ^E	122 (24) ^E	162 (22)	210 (22)	270 (28)	342 (41)	391 (60)
	19+	393	244 (11)	128 (12)	146 (12)	182 (12)	233 (13)	299 (17)	378 (27)	438 (37)
Female	9-13	79	159 (17)	79 (22) ^E	90 (23) ^E	114 (22) ^E	149 (21)	193 (25)	242 (36)	276 (46) ^E
	14-18	104	161 (13)	106 (20) ^E	116 (18)	133 (16)	155 (15)	180 (18)	208 (26)	228 (33)
	19-30	101	194 (24)	95 (29) ^E	110 (29) ^E	141 (29) ^E	187 (30)	247 (36)	315 (49)	362 (61) ^E
	31-50	143	199 (17)	108 (25) ^E	123 (24) ^E	151 (21)	190 (20)	235 (25)	283 (41)	316 (57) ^E
	51-70	193	197 (12)	91 (13)	107 (13)	136 (12)	176 (13)	227 (17)	290 (26)	337 (36)
	>70	94	166 (12)	88 (15) ^E	99 (15)	119 (14)	147 (15)	181 (19)	216 (27)	241 (33)
	19+	531	193 (9)	87 (8)	102 (9)	134 (9)	180 (10)	237 (12)	301 (16)	349 (21)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.5 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	311	148	(8)	72	86	112	143	180	220	247
	4-8	485	187	(9)	114	128	153	184	222	264	293
Male	9-13	277	225	(13)	146	161	189	225	266	309	338
	14-18	339	240	(12)	118	139	181	239	307	383	440
	19-30	237	286	(14)	177	200	243	295	353	412	452
	31-50	423	272	(9)	164	185	220	266	323	384	428
	51-70	387	259	(14)	141	161	199	252	319	404	476
	>70	132	215	(16)	98	116	154	204	263	329	380
	19+	1179	266	(7)	149	171	210	262	327	400	454
	9-13	281	185	(9)	102	115	140	173	212	253	281
Female	14-18	321	211	(11)	130	145	173	208	248	289	316
	19-30	249	255	(26)	145	161	192	235	286	344	384
	31-50	364	241	(24)	148	163	193	237	289	345	386
	51-70	467	227	(9)	120	139	174	219	277	344	391
	>70	215	189	(13)	98	113	141	180	235	307	366
	19+	1295	233	(11)	127	144	175	226	287	357	410

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.6 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	143	(5)	67 (8)	80 (7)	104 (6)	137 (5)	177 (7)	223 (12)	256 (17)
	4-8	956	167	(4)	89 (7)	103 (6)	127 (5)	159 (5)	199 (6)	245 (10)	279 (15)
Male	9-13	589	205	(7)	116 (9)	130 (9)	158 (8)	196 (7)	243 (10)	294 (16)	330 (23)
	14-18	639	240	(8)	168 (24)	182 (20)	208 (15)	241 (10)	278 (14)	315 (26)	340 (35)
	19-30	481	261	(12)	127 (20)	148 (18)	189 (15)	244 (13)	314 (19)	395 (35)	454 (49)
	31-50	709	249	(9)	187 (30)	199 (26)	221 (19)	248 (11)	277 (16)	306 (33)	325 (46)
	51-70	758	264	(8)	121 (10)	144 (10)	189 (9)	248 (9)	320 (12)	402 (21)	462 (30)
	>70	734	223	(8)	109 (8)	127 (8)	160 (7)	206 (8)	266 (11)	336 (18)	385 (24)
	19+	2682	253	(5)	134 (8)	154 (7)	190 (6)	240 (6)	302 (8)	371 (13)	419 (19)
	Female	9-13	585	178	(6)	92 (8)	106 (7)	133 (6)	168 (6)	211 (9)	260 (16)
14-18	645	198	(7)	95 (12)	112 (11)	143 (10)	186 (8)	241 (10)	303 (16)	346 (22)	
19-30	514	191	(8)	97 (15)	112 (14)	142 (11)	181 (10)	228 (12)	278 (19)	311 (24)	
31-50	758	226	(9)	96 (8)	113 (8)	148 (8)	201 (9)	276 (14)	371 (25)	445 (37)	
51-70	955	217	(6)	104 (8)	123 (8)	159 (7)	206 (7)	264 (9)	329 (14)	376 (19)	
>70	1345	199	(5)	106 (8)	121 (7)	151 (6)	191 (5)	236 (7)	282 (11)	315 (16)	
19+	3572	213	(5)	104 (4)	120 (4)	154 (4)	200 (5)	258 (6)	325 (10)	375 (14)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.7 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	123	(8)	60 (6)	70 (6)	89 (5)	115 (7)	147 (11)	187 (19)	217 (27)
	4-8	425	152	(8)	76 (11)	88 (10)	111 (8)	143 (7)	182 (11)	226 (19)	258 (27)
Male	9-13	274	178	(9)	123 (18)	134 (16)	153 (13)	177 (11)	203 (14)	230 (22)	249 (28)
	14-18	297	243	(12)	126 (16)	146 (15)	182 (13)	232 (14)	296 (21)	367 (33)	414 (43)
	19-30	249	264	(22)	134 (12)	154 (13)	192 (15)	244 (20)	315 (29)	396 (41)	451 (50)
	31-50	309	246	(15)	125 (20)	144 (19)	181 (18)	236 (17)	305 (25)	378 (38)	428 (49)
	51-70	277	226	(10)	126 (21)	143 (19)	175 (15)	217 (12)	267 (16)	319 (28)	354 (37)
	>70	136	201	(17)	86 (14)	104 (13)	140 (13)	187 (14)	242 (18)	302 (28)	346 (37)
	19+	971	240	(8)	126 (13)	145 (12)	180 (10)	228 (9)	288 (13)	354 (22)	400 (30)
	Female	9-13	265	161	(10)	101 (9)	111 (10)	130 (11)	153 (12)	177 (13)	200 (14)
14-18	290	177	(10)	83 (14) ^E	96 (14)	123 (13)	161 (13)	211 (17)	266 (27)	307 (35)	
19-30	197	208	(17)	126 (27) ^E	142 (25) ^E	172 (21)	210 (19)	250 (24)	287 (31)	309 (37)	
31-50	312	209	(14)	101 (11)	115 (11)	147 (11)	192 (13)	248 (18)	312 (28)	361 (37)	
51-70	312	219	(12)	126 (21) ^E	141 (20)	170 (16)	208 (14)	252 (18)	300 (30)	334 (40)	
>70	239	170	(9)	94 (18) ^E	107 (16)	132 (12)	165 (9)	207 (13)	251 (23)	282 (32)	
19+	1060	206	(7)	103 (7)	119 (7)	151 (7)	196 (7)	251 (10)	312 (16)	356 (21)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.8 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	129	121	(6)	68	(10)	78	(9)	94	(8)	114	(8)	139	(10)	172	(15)	200	(22)
	4-8	213	149	(8)	89	(12)	100	(11)	119	(9)	143	(9)	172	(12)	202	(17)	221	(22)
Male	9-13	122	189	(13)	122	(19)	133	(18)	156	(16)	187	(15)	225	(22)	267	(35)	295	(47)
	14-18	150	260	(46) ^E	157	(39) ^E	178	(40) ^E	218	(44) ^E	260	(48) ^E	301	(57) ^E	346	(83) ^E	378	(108) ^E
	19-30	106	255	(21)	99	(29) ^E	122	(28) ^E	171	(25)	241	(25)	332	(35)	435	(60)	508	(83)
	31-50	155	258	(21)	137	(38) ^E	158	(35) ^E	199	(31)	255	(27)	325	(34)	401	(53)	454	(70)
	51-70	122	279	(46)	142	(32) ^E	163	(31) ^E	198	(32)	246	(43) ^E	336	(78) ^E	F		F	
	>70	88	245	(16)	130	(21)	147	(19)	180	(17)	226	(17)	287	(25)	361	(42)	417	(59)
	19+	471	262	(15)	120	(13)	143	(12)	187	(11)	243	(12)	321	(19)	427	(41)	517	(72)
Female	9-13	103	193	(23)	124	(23) ^E	135	(23) ^E	155	(23)	182	(26)	216	(32)	251	(41)	275	(48) ^E
	14-18	142	166	(13)	86	(15) ^E	99	(14)	124	(14)	158	(16)	203	(22)	258	(35)	300	(45)
	19-30	111	184	(14)	91	(15) ^E	107	(14)	136	(13)	173	(15)	222	(22)	285	(38)	335	(54)
	31-50	146	196	(12)	114	(24) ^E	126	(22) ^E	148	(19)	177	(17)	216	(19)	260	(29)	293	(40)
	51-70	184	223	(12)	152	(26) ^E	167	(23)	191	(18)	222	(16)	256	(21)	293	(33)	318	(44)
	>70	143	176	(8)	111	(13)	123	(12)	147	(10)	177	(10)	211	(13)	246	(18)	269	(23)
	19+	584	198	(7)	106	(10)	122	(9)	150	(9)	187	(9)	234	(11)	289	(15)	330	(21)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.9 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	169	122 (9)	47 (15) ^E	59 (13) ^E	83 (11)	115 (10)	156 (13)	205 (23)	243 (33)
	4-8	281	156 (7)	96 (14)	106 (13)	124 (10)	150 (8)	181 (12)	216 (22)	240 (31)
Male	9-13	183	225 (16)	100 (22) ^E	118 (21) ^E	154 (19)	206 (17)	277 (24)	370 (48)	445 (71)
	14-18	187	248 (20)	145 (33) ^E	162 (30) ^E	196 (24)	240 (20)	296 (27)	358 (46)	403 (62)
	19-30	223	246 (16)	170 (34) ^E	186 (31)	214 (24)	245 (18)	279 (22)	314 (38)	337 (52)
	31-50	229	240 (12)	130 (19)	149 (18)	188 (16)	241 (17)	305 (22)	375 (33)	424 (42)
	51-70	197	229 (12)	131 (23) ^E	149 (21)	182 (17)	225 (15)	273 (19)	323 (30)	355 (40)
	>70	72	198 (16)	119 (18)	133 (17)	158 (17)	190 (18)	229 (22)	271 (31)	301 (40)
	19+	721	236 (7)	128 (11)	147 (11)	184 (10)	233 (10)	292 (11)	357 (16)	403 (21)
	Female	9-13	165	178 (10)	116 (22) ^E	129 (20)	153 (17)	183 (14)	219 (18)	257 (28)
14-18	206	185 (11)	83 (15) ^E	99 (15)	131 (14)	174 (14)	223 (15)	275 (21)	313 (29)	
19-30	191	202 (16)	100 (20) ^E	113 (20) ^E	141 (20)	183 (20)	239 (25)	303 (38)	347 (50)	
31-50	258	195 (11)	154 (28) ^E	163 (25)	179 (19)	199 (15)	221 (20)	244 (34)	259 (45) ^E	
51-70	249	200 (10)	111 (17)	127 (14)	154 (11)	189 (10)	232 (15)	281 (27)	317 (38)	
>70	128	168 (10)	93 (10)	106 (10)	130 (10)	161 (13)	197 (17)	238 (24)	268 (29)	
19+	826	195 (5)	104 (9)	120 (9)	147 (8)	190 (7)	238 (8)	291 (13)	330 (18)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.10 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	192	165	(10)	94	(14)	107	(13)	131	(12)	162	(12)	198	(16)	235	(23)	260	(29)
	4-8	321	179	(6)	120	(16)	131	(14)	151	(11)	177	(8)	207	(12)	238	(20)	259	(26)
Male																		
	9-13	226	216	(12)	110	(13)	127	(13)	163	(13)	211	(15)	269	(18)	333	(25)	377	(31)
	14-18	262	285	(23)	128	(27) ^E	154	(26) ^E	204	(25)	274	(25)	361	(38)	460	(70)	530	(82)
	19-30	197	299	(25)	165	(31) ^E	186	(30)	227	(27)	282	(26)	350	(32)	423	(48)	472	(62)
	31-50	282	297	(17)	170	(26)	192	(23)	232	(20)	284	(19)	348	(26)	412	(38)	450	(47)
	51-70	234	280	(16)	148	(19)	169	(17)	209	(16)	260	(17)	325	(23)	401	(35)	456	(47)
	>70	119	257	(17)	121	(18)	142	(18)	186	(20)	245	(23)	318	(29)	397	(36)	451	(43)
	19+	832	289	(10)	148	(9)	170	(9)	213	(9)	270	(11)	344	(16)	427	(22)	484	(28)
Female																		
	9-13	226	198	(12)	129	(17)	142	(16)	165	(15)	195	(15)	231	(18)	267	(26)	292	(32)
	14-18	242	201	(16)	90	(15)	106	(14)	136	(15)	182	(16)	247	(24)	324	(40)	383	(55)
	19-30	208	221	(17)	126	(20)	142	(19)	171	(18)	213	(20)	269	(27)	336	(42)	384	(56)
	31-50	263	248	(13)	118	(14)	137	(14)	173	(15)	224	(17)	289	(23)	361	(32)	409	(38)
	51-70	322	234	(10)	129	(18)	149	(16)	186	(14)	233	(13)	288	(18)	344	(27)	381	(33)
	>70	198	196	(9)	106	(11)	122	(11)	153	(12)	192	(13)	231	(14)	269	(17)	295	(20)
	19+	991	233	(7)	119	(7)	137	(7)	171	(7)	220	(9)	282	(12)	350	(17)	396	(21)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.11 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	348	129 (5)	67 (5)	76 (5)	94 (5)	121 (6)	155 (9)	189 (12)	212 (15)
	4-8	554	167 (9)	91 (7)	103 (7)	127 (7)	160 (10)	206 (15)	265 (23)	315 (32)
Male	9-13	409	192 (10)	116 (12)	128 (12)	152 (11)	184 (11)	222 (15)	266 (23)	297 (30)
	14-18	414	211 (8)	118 (11)	136 (10)	169 (10)	211 (11)	262 (13)	318 (17)	358 (20)
	19-30	311	233 (12)	120 (13)	138 (13)	173 (12)	221 (13)	279 (18)	340 (26)	382 (33)
	31-50	489	230 (9)	133 (12)	149 (11)	179 (10)	219 (10)	264 (13)	314 (20)	350 (27)
	51-70	575	235 (12)	124 (12)	142 (11)	177 (11)	219 (12)	273 (18)	347 (31)	403 (42)
	>70	239	208 (10)	123 (11)	137 (11)	163 (11)	199 (12)	240 (13)	286 (17)	319 (21)
	19+	1614	230 (5)	122 (5)	140 (5)	173 (5)	217 (6)	271 (8)	333 (12)	379 (16)
Female	9-13	355	158 (8)	94 (9)	106 (9)	127 (9)	154 (10)	189 (13)	225 (17)	251 (20)
	14-18	410	159 (8)	79 (14) ^E	94 (13)	120 (11)	153 (10)	193 (12)	241 (18)	276 (25)
	19-30	384	180 (11)	100 (10)	113 (10)	139 (11)	172 (13)	215 (17)	265 (22)	300 (26)
	31-50	585	189 (7)	90 (7)	105 (8)	138 (8)	180 (10)	235 (13)	301 (17)	348 (21)
	51-70	711	204 (8)	98 (7)	115 (6)	145 (6)	185 (7)	237 (10)	305 (18)	362 (27)
	>70	342	177 (9)	96 (7)	109 (7)	133 (7)	166 (9)	205 (13)	254 (22)	295 (31)
	19+	2022	190 (4)	94 (4)	109 (4)	139 (4)	179 (5)	232 (7)	295 (10)	343 (13)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.12 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	622	122	(5)	52 (7)	63 (7)	86 (5)	114 (6)	151 (8)	196 (13)	230 (19)
	4-8	919	154	(5)	88 (6)	98 (6)	119 (5)	147 (6)	181 (8)	218 (12)	245 (16)
Male	9-13	579	208	(10)	122 (18)	137 (17)	166 (14)	203 (11)	249 (16)	301 (28)	339 (39)
	14-18	634	249	(15)	140 (18)	157 (17)	191 (15)	237 (14)	297 (20)	370 (36)	424 (50)
	19-30	578	251	(12)	135 (25) ^E	156 (22)	195 (17)	242 (14)	299 (18)	364 (32)	412 (44)
	31-50	693	244	(9)	125 (11)	146 (11)	187 (11)	242 (12)	310 (15)	387 (21)	441 (26)
	51-70	596	238	(11)	123 (16)	143 (14)	179 (11)	227 (10)	280 (14)	360 (34)	429 (62)
	>70	296	210	(10)	104 (12)	121 (12)	152 (11)	194 (11)	247 (13)	312 (20)	362 (29)
	19+	2163	241	(5)	123 (6)	144 (6)	182 (6)	233 (7)	297 (8)	375 (14)	432 (20)
Female	9-13	533	177	(8)	115 (16)	126 (15)	148 (12)	176 (10)	209 (13)	243 (22)	267 (27)
	14-18	638	180	(7)	81 (7)	96 (7)	126 (8)	167 (9)	218 (11)	276 (14)	317 (18)
	19-30	499	200	(11)	103 (12)	118 (12)	147 (12)	189 (13)	242 (16)	300 (23)	340 (28)
	31-50	716	198	(8)	118 (17)	132 (16)	158 (13)	192 (11)	235 (12)	281 (18)	314 (25)
	51-70	745	209	(7)	118 (10)	133 (9)	162 (8)	199 (7)	246 (11)	299 (18)	337 (25)
	>70	510	170	(6)	95 (6)	108 (6)	131 (6)	165 (7)	207 (9)	250 (12)	281 (16)
	19+	2470	198	(4)	103 (5)	118 (5)	148 (5)	190 (5)	242 (6)	299 (9)	342 (12)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 33.13 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	141 (3)	66 (4)	79 (4)	103 (3)	135 (4)	175 (4)	220 (7)	252 (9)
	4-8	3235	171 (3)	95 (4)	108 (3)	132 (3)	164 (4)	204 (5)	249 (7)	282 (9)
Male	9-13	2080	210 (5)	120 (5)	135 (5)	164 (5)	204 (6)	251 (7)	303 (10)	340 (13)
	14-18	2288	245 (6)	127 (6)	147 (6)	187 (6)	239 (7)	304 (9)	381 (14)	437 (19)
	19-30	1804	268 (7)	135 (8)	157 (7)	200 (7)	257 (8)	326 (10)	403 (15)	458 (20)
	31-50	2596	259 (5)	141 (6)	161 (6)	199 (5)	248 (6)	311 (8)	381 (11)	430 (15)
	51-70	2550	259 (6)	129 (5)	150 (5)	191 (5)	244 (5)	310 (8)	396 (16)	463 (26)
	>70	1520	223 (6)	110 (5)	128 (5)	162 (6)	209 (7)	269 (9)	336 (13)	384 (16)
	19+	8470	258 (3)	131 (3)	153 (3)	192 (3)	245 (3)	312 (5)	389 (7)	445 (10)
Female	9-13	1980	181 (4)	101 (4)	115 (4)	140 (4)	173 (4)	213 (6)	257 (9)	287 (11)
	14-18	2256	195 (4)	96 (4)	112 (4)	143 (4)	185 (5)	238 (7)	299 (10)	341 (12)
	19-30	1854	211 (8)	105 (6)	121 (7)	152 (7)	196 (8)	252 (10)	316 (14)	362 (18)
	31-50	2686	225 (7)	109 (4)	125 (4)	160 (5)	208 (6)	275 (10)	354 (16)	415 (22)
	51-70	3200	220 (4)	113 (4)	132 (4)	166 (4)	211 (4)	266 (5)	329 (8)	374 (11)
	>70	2610	190 (4)	98 (4)	114 (4)	143 (4)	182 (5)	232 (7)	291 (11)	333 (15)
	19+	10350	217 (3)	107 (2)	124 (2)	158 (3)	203 (3)	265 (5)	335 (8)	387 (10)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

34. Protein (g/d): Usual intakes from food

Table 34.1 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	57 (4)	47 (7)	48 (6)	52 (5)	55 (4)	59 (7)	62 (10)	65 (12) ^E
	4-8	127	71 (5)	51 (7)	54 (7)	61 (6)	69 (6)	78 (7)	88 (10)	95 (14)
	9-13	111	96 (7)	71 (10)	76 (9)	84 (8)	95 (8)	107 (10)	118 (15)	126 (19)
Male	14-18	107	98 (6)	70 (11)	75 (10)	85 (9)	98 (8)	113 (9)	128 (14)	138 (17)
	19-30	77	105 (5)	72 (7)	79 (7)	92 (7)	107 (7)	122 (7)	136 (9)	145 (10)
	31-50	145	111 (11)	63 (12) ^E	71 (11)	86 (9)	105 (11)	126 (16)	151 (23)	168 (29) ^E
	51-70	182	86 (4)	56 (5)	61 (5)	71 (5)	82 (5)	95 (6)	107 (6)	114 (6)
	>70	63	101 (8)	76 (9)	82 (10)	95 (12)	112 (14)	130 (17)	148 (20)	161 (23)
	19+	467	102 (5)	71 (8)	77 (7)	88 (6)	101 (5)	115 (6)	129 (9)	138 (11)
	9-13	96	70 (5)	45 (9) ^E	50 (8)	58 (7)	69 (6)	82 (7)	95 (9)	103 (11)
	14-18	105	80 (11)	51 (12) ^E	56 (11) ^E	66 (10)	79 (10)	95 (13)	114 (20) ^E	127 (25) ^E
Female	19-30	91	66 (4)	43 (6)	47 (5)	54 (5)	62 (5)	72 (7)	81 (9)	87 (11)
	31-50	167	72 (5)	41 (6)	47 (6)	57 (6)	71 (7)	87 (8)	104 (10)	115 (12)
	51-70	198	76 (5)	46 (9) ^E	52 (8)	61 (6)	73 (6)	88 (8)	105 (13)	117 (17)
	>70	74	71 (6)	46 (4)	50 (4)	58 (5)	69 (6)	82 (8)	95 (11)	104 (12)
	19+	530	72 (3)	45 (4)	50 (4)	59 (3)	71 (3)	85 (4)	99 (6)	109 (7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.2 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	57 (5)	45 (5)	47 (5)	52 (5)	58 (6)	63 (6)	69 (7)	72 (7)
	4-8	110	64 (3)	47 (3)	50 (4)	56 (4)	63 (4)	70 (4)	77 (4)	82 (4)
Male	9-13	95	84 (5)	62 (10)	66 (9)	73 (9)	82 (8)	92 (9)	102 (10)	108 (11)
	14-18	87	105 (6)	64 (12) ^E	72 (11)	87 (9)	106 (9)	127 (11)	150 (16)	166 (21)
	19-30	70	118 (8)	72 (13) ^E	82 (12)	101 (10)	123 (9)	146 (11)	168 (15)	182 (17)
	31-50	109	96 (6)	62 (9)	68 (9)	79 (7)	94 (7)	110 (9)	127 (13)	138 (16)
	51-70	128	91 (5)	54 (6)	60 (6)	72 (6)	87 (6)	106 (8)	127 (11)	141 (14)
	>70	65	67 (4)	48 (4)	51 (4)	56 (5)	63 (5)	70 (6)	78 (7)	82 (7)
	19+	372	96 (3)	57 (4)	64 (4)	77 (4)	94 (4)	115 (5)	136 (7)	151 (9)
Female	9-13	75	72 (7)	47 (10) ^E	52 (9) ^E	61 (9)	72 (10)	84 (12)	96 (15)	105 (18) ^E
	14-18	81	68 (5)	42 (8) ^E	47 (7)	56 (6)	66 (6)	77 (7)	87 (9)	93 (10)
	19-30	101	79 (8)	45 (7)	52 (7)	66 (8)	83 (10)	102 (14)	120 (18)	130 (20)
	31-50	116	72 (5)	47 (8) ^E	52 (7)	61 (7)	72 (7)	83 (8)	93 (9)	100 (11)
	51-70	146	73 (3)	44 (5)	50 (5)	60 (5)	73 (5)	87 (5)	102 (7)	111 (8)
	>70	94	57 (3)	37 (6)	41 (5)	47 (4)	54 (4)	63 (5)	72 (7)	77 (9)
	19+	457	72 (3)	45 (3)	51 (3)	60 (3)	72 (4)	85 (5)	98 (6)	106 (7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.3 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	56 (4)	42 (5)	44 (5)	48 (5)	54 (5)	61 (6)	70 (8)	75 (10)
	4-8	177	68 (3)	41 (7) ^E	46 (7)	55 (5)	67 (3)	78 (5)	89 (8)	96 (11)
Male	9-13	111	90 (5)	53 (6)	58 (6)	70 (6)	85 (5)	103 (6)	123 (9)	137 (13)
	14-18	113	96 (9)	76 (12)	80 (11)	86 (11)	94 (14)	102 (19) ^E	110 (25) ^E	114 (30) ^E
	19-30	91	107 (9)	57 (16) ^E	66 (14) ^E	82 (11)	103 (9)	126 (13)	149 (20)	164 (26)
	31-50	101	100 (6)	69 (11)	75 (10)	86 (8)	100 (8)	115 (11)	131 (15)	141 (19)
	51-70	134	99 (7)	74 (10)	78 (9)	87 (7)	98 (7)	110 (9)	122 (13)	130 (17)
	>70	56	77 (6)	52 (10) ^E	57 (9) ^E	66 (8)	77 (7)	90 (9)	102 (13)	110 (17)
	19+	382	99 (4)	62 (5)	69 (5)	81 (4)	97 (5)	115 (6)	135 (9)	148 (12)
Female	9-13	105	72 (6)	50 (7)	54 (7)	62 (6)	72 (6)	83 (8)	95 (11)	103 (13)
	14-18	120	64 (6)	40 (9) ^E	45 (8) ^E	54 (7)	64 (6)	76 (8)	87 (11)	95 (14)
	19-30	91	73 (4)	54 (4)	58 (4)	65 (5)	73 (4)	81 (5)	88 (5)	92 (5)
	31-50	159	76 (4)	40 (6)	47 (6)	61 (6)	77 (5)	93 (6)	108 (7)	118 (8)
	51-70	174	70 (4)	49 (7)	53 (7)	59 (5)	68 (5)	77 (6)	86 (8)	92 (10)
	>70	80	62 (5)	37 (7) ^E	42 (7)	51 (6)	62 (6)	75 (8)	88 (12)	98 (15)
	19+	504	72 (2)	43 (3)	49 (3)	59 (3)	72 (3)	85 (4)	99 (5)	108 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.4 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	64 (4)	44 (7)	48 (6)	55 (5)	65 (5)	76 (6)	87 (9)	95 (12)
	4-8	140	73 (5)	52 (8)	56 (7)	64 (6)	73 (5)	84 (6)	95 (9)	103 (11)
Male	9-13	92	88 (6)	63 (10)	67 (9)	76 (8)	85 (7)	95 (8)	106 (11)	113 (14)
	14-18	107	114 (7)	73 (11)	81 (10)	96 (8)	117 (9)	140 (13)	165 (19)	182 (24)
	19-30	73	119 (11)	68 (13) ^E	78 (12)	96 (11)	118 (12)	149 (19)	186 (30)	213 (40) ^E
	31-50	134	99 (6)	64 (6)	70 (6)	81 (6)	95 (7)	112 (9)	131 (10)	144 (12)
	51-70	131	93 (6)	63 (6)	69 (6)	78 (6)	89 (7)	102 (7)	116 (8)	125 (9)
	>70	55	78 (5)	47 (9) ^E	53 (8)	63 (7)	75 (7)	87 (8)	100 (10)	108 (13)
	19+	393	100 (4)	63 (6)	69 (6)	81 (5)	98 (5)	117 (6)	137 (10)	151 (13)
Female	9-13	79	75 (7)	48 (9) ^E	53 (9) ^E	64 (8)	77 (9)	92 (11)	107 (15)	118 (20) ^E
	14-18	104	70 (3)	53 (4)	57 (4)	64 (4)	72 (4)	79 (5)	87 (5)	92 (6)
	19-30	101	70 (5)	47 (8)	52 (7)	61 (7)	74 (7)	89 (8)	106 (11)	116 (14)
	31-50	143	66 (4)	39 (8) ^E	45 (7)	54 (5)	65 (5)	75 (6)	84 (9)	90 (12)
	51-70	193	67 (4)	45 (8) ^E	48 (7)	55 (6)	64 (5)	74 (6)	85 (10)	92 (13)
	>70	94	61 (4)	46 (7)	48 (7)	53 (6)	58 (6)	64 (7)	70 (9)	73 (10)
	19+	531	66 (2)	43 (4)	48 (4)	55 (3)	65 (3)	76 (3)	87 (5)	94 (7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.5 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	311	62 (3)	44 (7)	48 (6)	55 (5)	63 (4)	72 (4)	80 (5)	86 (6)
	4-8	485	75 (3)	53 (6)	57 (5)	64 (4)	74 (3)	84 (4)	93 (6)	99 (8)
Male	9-13	277	103 (6)	76 (11)	82 (10)	94 (9)	108 (8)	124 (10)	140 (15)	151 (18)
	14-18	339	117 (6)	72 (8)	81 (7)	98 (7)	119 (8)	147 (10)	179 (15)	201 (19)
	19-30	237	103 (5)	89 (12)	93 (11)	99 (8)	106 (7)	114 (8)	121 (12)	125 (15)
	31-50	423	105 (5)	67 (8)	74 (8)	87 (6)	104 (6)	124 (7)	144 (11)	158 (14)
	51-70	387	91 (3)	58 (7)	65 (6)	76 (5)	90 (4)	105 (5)	122 (7)	133 (9)
	>70	132	78 (5)	45 (8) ^E	51 (8)	61 (8)	75 (8)	92 (9)	111 (10)	124 (12)
	19+	1179	98 (3)	62 (4)	68 (4)	81 (3)	98 (3)	116 (4)	135 (5)	148 (7)
Female	9-13	281	77 (4)	48 (6)	53 (6)	62 (5)	74 (4)	87 (5)	101 (7)	110 (9)
	14-18	321	75 (3)	50 (4)	55 (3)	64 (3)	76 (4)	90 (5)	104 (6)	114 (8)
	19-30	249	79 (4)	58 (8)	62 (7)	70 (5)	79 (5)	88 (6)	97 (9)	103 (11)
	31-50	364	85 (5)	55 (9)	61 (8)	71 (7)	85 (6)	101 (7)	118 (10)	130 (13)
	51-70	467	76 (3)	54 (6)	59 (5)	67 (4)	76 (3)	87 (4)	97 (7)	104 (9)
	>70	215	62 (2)	45 (4)	48 (4)	54 (4)	62 (4)	70 (4)	78 (6)	83 (7)
	19+	1295	78 (2)	51 (3)	57 (3)	66 (3)	78 (3)	92 (3)	106 (4)	115 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.6 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	644	55 (1)	34 (3)	38 (2)	45 (2)	55 (2)	66 (2)	76 (4)	83 (5)
	4-8	956	66 (1)	44 (3)	48 (2)	55 (2)	65 (1)	76 (2)	87 (3)	94 (4)
Male										
	9-13	589	85 (2)	58 (7)	64 (6)	73 (4)	84 (3)	96 (4)	109 (7)	118 (10)
	14-18	639	106 (3)	64 (7)	72 (6)	87 (5)	105 (4)	126 (5)	149 (7)	164 (10)
	19-30	481	106 (4)	59 (13) ^E	67 (11) ^E	81 (8)	101 (5)	124 (8)	149 (14)	165 (20)
	31-50	709	102 (3)	68 (11)	75 (9)	87 (7)	102 (4)	119 (5)	136 (11)	148 (15)
	51-70	758	90 (2)	59 (7)	64 (6)	74 (4)	87 (3)	102 (4)	118 (8)	129 (12)
	>70	734	72 (2)	46 (5)	51 (4)	60 (3)	71 (2)	84 (3)	97 (5)	104 (7)
	19+	2682	97 (2)	57 (4)	64 (4)	77 (3)	94 (2)	115 (3)	136 (5)	151 (7)
Female										
	9-13	585	70 (2)	46 (5)	51 (4)	58 (3)	68 (2)	78 (3)	89 (5)	96 (7)
	14-18	645	73 (2)	43 (4)	49 (4)	59 (3)	72 (2)	87 (3)	101 (5)	111 (7)
	19-30	514	69 (2)	38 (5)	43 (4)	53 (3)	66 (3)	80 (3)	95 (6)	105 (8)
	31-50	758	71 (2)	42 (3)	47 (3)	57 (2)	70 (2)	84 (3)	98 (4)	107 (5)
	51-70	955	71 (2)	47 (6)	52 (5)	60 (4)	70 (2)	81 (3)	92 (6)	99 (8)
	>70	1345	62 (1)	36 (2)	40 (2)	48 (2)	59 (2)	72 (2)	85 (3)	95 (4)
	19+	3572	69 (1)	40 (2)	46 (2)	55 (1)	67 (1)	81 (2)	96 (2)	106 (3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.7 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	324	51	(2)	31	(3)	34	(3)	41	(3)	50	(2)	61	(3)	72	(5)	80	(7)
	4-8	425	64	(3)	52	(8)	54	(7)	59	(5)	64	(4)	69	(5)	75	(8)	78	(10)
	9-13	274	84	(2)	57	(7)	62	(6)	72	(4)	84	(3)	97	(5)	110	(8)	118	(10)
Male	14-18	297	111	(4)	56	(7)	66	(6)	85	(6)	108	(6)	136	(7)	168	(11)	192	(14)
	19-30	249	114	(8)	66	(5)	73	(5)	88	(6)	107	(7)	127	(9)	148	(11)	163	(12)
	31-50	309	96	(6)	55	(10) ^E	62	(9)	74	(7)	90	(6)	109	(9)	132	(14)	149	(19)
	51-70	277	91	(4)	53	(6)	60	(5)	72	(4)	87	(4)	106	(5)	126	(7)	139	(9)
	>70	136	79	(5)	39	(9) ^E	46	(8) ^E	58	(7)	75	(6)	95	(8)	115	(12)	129	(15)
	19+	971	97	(3)	55	(6)	62	(5)	75	(4)	92	(3)	112	(4)	134	(7)	149	(9)
	9-13	265	70	(4)	43	(7) ^E	48	(7)	57	(5)	68	(4)	80	(5)	94	(7)	103	(10)
	14-18	290	69	(3)	52	(6)	55	(5)	61	(4)	68	(3)	75	(5)	82	(7)	86	(9)
Female	19-30	197	76	(5)	50	(9) ^E	55	(8)	64	(7)	75	(6)	87	(7)	99	(10)	106	(12)
	31-50	312	79	(4)	44	(7)	50	(7)	61	(5)	75	(5)	92	(6)	109	(10)	121	(13)
	51-70	312	73	(4)	50	(8)	55	(7)	62	(5)	72	(4)	83	(5)	94	(8)	101	(11)
	>70	239	57	(3)	35	(5)	39	(4)	47	(3)	56	(3)	67	(4)	78	(6)	86	(8)
	19+	1060	74	(2)	41	(3)	46	(3)	57	(3)	71	(2)	87	(3)	104	(5)	116	(7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.8 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	54 (3)	34 (4)	38 (4)	45 (4)	54 (4)	64 (5)	74 (6)	81 (7)
	4-8	213	65 (2)	48 (5)	51 (4)	57 (3)	65 (3)	73 (4)	81 (6)	86 (7)
	9-13	122	88 (6)	68 (6)	73 (6)	81 (7)	91 (8)	102 (9)	114 (11)	122 (13)
Male	14-18	150	114 (6)	67 (8)	75 (8)	91 (7)	111 (8)	135 (9)	159 (12)	175 (15)
	19-30	106	108 (7)	47 (10) ^E	58 (10) ^E	78 (8)	104 (8)	133 (11)	164 (16)	183 (20)
	31-50	155	97 (5)	50 (8) ^E	59 (8)	75 (7)	95 (6)	117 (7)	139 (9)	153 (11)
	51-70	122	92 (5)	60 (11) ^E	67 (10)	79 (8)	94 (7)	111 (9)	127 (13)	137 (16)
	>70	88	85 (5)	50 (7)	56 (7)	67 (6)	83 (6)	102 (8)	123 (13)	137 (17)
	19+	471	97 (3)	48 (4)	57 (4)	73 (3)	95 (4)	119 (5)	144 (6)	161 (8)
	9-13	103	72 (4)	52 (7)	56 (6)	63 (6)	71 (5)	81 (6)	91 (8)	98 (10)
	14-18	142	72 (4)	44 (4)	49 (4)	59 (4)	71 (5)	84 (6)	98 (7)	107 (9)
Female	19-30	111	67 (3)	38 (6)	43 (5)	53 (5)	64 (5)	78 (6)	91 (7)	99 (9)
	31-50	146	79 (4)	36 (7) ^E	42 (7) ^E	59 (6)	75 (5)	89 (5)	102 (8)	115 (9)
	51-70	184	73 (3)	51 (3)	55 (3)	63 (4)	72 (4)	82 (5)	93 (7)	100 (7)
	>70	143	68 (4)	38 (5)	43 (5)	53 (5)	68 (5)	86 (7)	108 (10)	123 (13)
	19+	584	73 (2)	40 (3)	46 (3)	57 (3)	70 (3)	87 (3)	102 (5)	112 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.9 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	169	50 (2)	26 (3)	30 (3)	39 (3)	49 (3)	60 (3)	72 (5)	81 (6)
	4-8	281	66 (2)	44 (6)	48 (5)	55 (4)	64 (3)	74 (4)	85 (7)	92 (9)
Male	9-13	183	92 (6)	69 (10)	74 (10)	82 (8)	94 (7)	107 (8)	122 (13)	132 (17)
	14-18	187	110 (5)	77 (12)	84 (11)	96 (8)	112 (7)	130 (9)	147 (14)	159 (19)
	19-30	223	105 (5)	67 (5)	74 (5)	87 (5)	103 (6)	119 (6)	135 (7)	145 (7)
	31-50	229	101 (6)	65 (12) ^E	72 (10)	84 (8)	100 (7)	119 (9)	140 (15)	154 (19)
	51-70	197	90 (5)	51 (11) ^E	59 (9)	73 (7)	89 (6)	106 (7)	122 (10)	132 (13)
	>70	72	82 (6)	69 (10)	71 (9)	76 (7)	81 (7)	86 (8)	90 (11)	93 (12)
	19+	721	98 (3)	63 (6)	70 (5)	83 (4)	97 (3)	114 (5)	132 (8)	144 (10)
Female	9-13	165	69 (3)	50 (7)	54 (6)	61 (5)	69 (4)	79 (5)	90 (9)	97 (12)
	14-18	206	69 (3)	49 (7)	52 (6)	58 (4)	66 (4)	76 (6)	85 (8)	92 (10)
	19-30	191	72 (5)	49 (9) ^E	53 (8)	62 (7)	72 (6)	84 (7)	96 (11)	104 (14)
	31-50	258	73 (4)	59 (10) ^E	63 (9)	68 (7)	75 (5)	82 (6)	89 (8)	94 (11)
	51-70	249	68 (3)	42 (6)	47 (5)	56 (4)	67 (3)	79 (4)	93 (7)	102 (10)
	>70	128	73 (6)	45 (5)	51 (5)	61 (6)	72 (6)	85 (7)	99 (9)	109 (11)
	19+	826	71 (2)	45 (3)	51 (3)	60 (3)	72 (3)	85 (3)	98 (4)	107 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.10 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	192	58 (2)	44 (4)	47 (4)	52 (3)	58 (3)	64 (4)	69 (6)	73 (7)
	4-8	321	66 (2)	42 (4)	46 (3)	54 (3)	64 (2)	76 (4)	87 (5)	95 (6)
Male										
	9-13	226	85 (3)	50 (5)	57 (4)	68 (4)	83 (4)	101 (5)	121 (7)	135 (9)
	14-18	262	116 (9)	69 (11)	77 (10)	93 (9)	113 (8)	137 (11)	162 (17)	179 (21)
	19-30	197	113 (6)	70 (8)	77 (8)	91 (6)	109 (7)	132 (10)	157 (15)	174 (20)
	31-50	282	121 (6)	66 (10)	76 (9)	95 (8)	119 (8)	147 (9)	175 (13)	194 (17)
	51-70	234	102 (5)	61 (6)	68 (6)	80 (5)	96 (6)	115 (7)	136 (11)	150 (14)
	>70	119	80 (5)	47 (8)	53 (7)	63 (7)	77 (7)	92 (8)	108 (11)	118 (14)
	19+	832	110 (3)	61 (4)	69 (3)	85 (3)	106 (4)	132 (5)	160 (7)	178 (8)
Female										
	9-13	226	74 (4)	46 (5)	51 (5)	60 (5)	71 (5)	85 (7)	101 (9)	112 (11)
	14-18	242	73 (3)	37 (5)	43 (4)	55 (4)	69 (4)	86 (5)	104 (6)	115 (8)
	19-30	208	75 (4)	54 (8)	59 (7)	67 (6)	76 (5)	87 (6)	97 (8)	103 (10)
	31-50	263	81 (4)	46 (5)	52 (4)	63 (4)	78 (5)	96 (7)	116 (10)	130 (13)
	51-70	322	73 (5)	49 (7)	54 (7)	62 (5)	73 (5)	85 (7)	97 (10)	105 (13)
	>70	198	68 (4)	41 (8) ^E	46 (7)	55 (6)	67 (6)	80 (7)	93 (9)	102 (12)
	19+	991	76 (2)	47 (3)	52 (3)	62 (2)	75 (3)	90 (3)	107 (5)	118 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.11 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	348	59	(2)	45	48	53	59	66	73	77	(4)	(4)	(3)	(3)	(3)	(5)	(6)
	4-8	554	70	(2)	50	54	61	71	81	91	98	(4)	(4)	(3)	(3)	(3)	(5)	(6)
Male																		
	9-13	409	90	(3)	62	67	76	88	100	114	123	(5)	(5)	(4)	(4)	(5)	(7)	(8)
	14-18	414	103	(4)	64	71	85	102	122	142	156	(6)	(6)	(6)	(7)	(9)	(10)	
	19-30	311	111	(5)	65	74	90	110	133	158	176	(6)	(6)	(6)	(8)	(12)	(15)	
	31-50	489	102	(4)	64	71	83	99	118	138	152	(6)	(5)	(5)	(7)	(10)	(12)	
	51-70	575	93	(3)	64	69	79	91	104	118	127	(6)	(5)	(4)	(5)	(7)	(9)	
	>70	239	82	(3)	58	63	72	83	95	108	116	(6)	(5)	(5)	(6)	(8)	(10)	
	19+	1614	100	(2)	62	68	81	97	117	137	151	(3)	(2)	(3)	(3)	(5)	(6)	
Female																		
	9-13	355	73	(3)	48	52	61	73	86	100	109	(5)	(5)	(4)	(5)	(6)	(8)	
	14-18	410	69	(4)	44	50	59	70	82	94	103	(5)	(5)	(4)	(5)	(8)	(10)	
	19-30	384	71	(3)	49	54	63	73	84	94	101	(4)	(3)	(3)	(4)	(5)	(6)	
	31-50	585	72	(2)	40	46	58	72	86	101	110	(4)	(4)	(3)	(4)	(5)	(6)	
	51-70	711	70	(2)	44	49	58	68	81	94	103	(4)	(3)	(3)	(4)	(6)	(7)	
	>70	342	63	(3)	40	44	52	62	73	85	93	(4)	(4)	(4)	(4)	(6)	(7)	
	19+	2022	70	(1)	44	49	58	70	82	96	104	(2)	(2)	(2)	(2)	(3)	(3)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.12 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	622	51	(1)	30	(2)	34	(2)	41	(2)	50	(2)	60	(2)	71	(3)	78	(3)
	4-8	919	65	(2)	46	(4)	49	(3)	56	(2)	64	(2)	73	(3)	82	(4)	88	(6)
Male																		
	9-13	579	90	(4)	68	(8)	72	(7)	80	(5)	91	(4)	103	(5)	116	(8)	124	(11)
	14-18	634	111	(4)	63	(5)	72	(5)	88	(5)	110	(5)	135	(6)	163	(8)	182	(11)
	19-30	578	107	(4)	66	(9)	74	(7)	88	(5)	104	(4)	123	(6)	141	(10)	153	(13)
	31-50	693	99	(4)	59	(7)	66	(6)	79	(5)	96	(5)	117	(6)	141	(8)	157	(11)
	51-70	596	90	(3)	50	(5)	58	(5)	72	(4)	90	(4)	108	(4)	127	(6)	139	(7)
	>70	296	82	(3)	48	(5)	54	(5)	65	(5)	79	(4)	95	(5)	112	(7)	123	(8)
	19+	2163	98	(2)	55	(3)	63	(3)	77	(2)	95	(2)	116	(3)	139	(4)	155	(5)
Female																		
	9-13	533	70	(2)	51	(5)	55	(5)	61	(4)	70	(3)	79	(4)	88	(6)	94	(7)
	14-18	638	70	(2)	46	(4)	51	(3)	58	(3)	67	(3)	79	(3)	90	(5)	96	(6)
	19-30	499	72	(3)	45	(6)	50	(6)	59	(5)	70	(4)	83	(4)	96	(6)	105	(8)
	31-50	716	75	(3)	46	(6)	51	(5)	61	(5)	75	(4)	89	(4)	104	(5)	114	(7)
	51-70	745	70	(2)	49	(5)	53	(4)	60	(3)	69	(2)	80	(3)	90	(5)	97	(7)
	>70	510	67	(3)	37	(2)	42	(2)	53	(3)	66	(3)	82	(5)	100	(6)	113	(7)
	19+	2470	72	(1)	42	(2)	48	(2)	58	(2)	71	(2)	86	(2)	102	(3)	112	(3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 34.13 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	56 (1)	37 (2)	41 (2)	48 (1)	56 (1)	66 (1)	76 (2)	83 (2)
	4-8	3235	68 (1)	45 (1)	49 (1)	57 (1)	67 (1)	78 (1)	90 (2)	98 (3)
	9-13	2080	90 (2)	62 (3)	67 (3)	77 (2)	90 (2)	104 (3)	119 (4)	128 (4)
Male	14-18	2288	110 (2)	65 (3)	74 (3)	90 (3)	110 (3)	134 (3)	160 (5)	178 (6)
	19-30	1804	107 (2)	65 (4)	73 (4)	87 (3)	104 (3)	124 (4)	146 (6)	160 (7)
	31-50	2596	105 (2)	63 (3)	70 (3)	85 (3)	103 (2)	125 (3)	148 (5)	163 (6)
	51-70	2550	92 (2)	56 (3)	63 (2)	74 (2)	89 (2)	107 (2)	125 (3)	137 (4)
	>70	1520	77 (2)	47 (2)	53 (2)	62 (2)	75 (2)	90 (3)	104 (3)	114 (4)
	19+	8470	99 (1)	58 (1)	65 (1)	79 (1)	97 (1)	118 (2)	140 (2)	155 (3)
	9-13	1980	72 (1)	46 (2)	51 (2)	59 (1)	70 (1)	82 (2)	96 (3)	104 (4)
	14-18	2256	73 (1)	44 (2)	49 (1)	59 (1)	72 (1)	86 (2)	100 (2)	110 (3)
Female	19-30	1854	73 (2)	46 (2)	51 (2)	60 (2)	71 (2)	84 (2)	96 (3)	105 (4)
	31-50	2686	76 (2)	44 (2)	50 (2)	61 (2)	75 (2)	92 (2)	109 (3)	120 (4)
	51-70	3200	72 (1)	49 (2)	54 (2)	62 (2)	72 (1)	83 (2)	94 (3)	102 (3)
	>70	2610	64 (1)	39 (2)	44 (2)	52 (1)	62 (2)	74 (2)	87 (2)	96 (3)
	19+	10350	73 (1)	44 (1)	50 (1)	59 (1)	72 (1)	86 (1)	101 (2)	111 (2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

35. Total carbohydrates (g/d): Usual intakes from food

Table 35.1 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						EAR ²	% <EAR (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	79	218 (14)	151 (16)	163 (16)	185 (17)	211 (18)	238 (19)	264 (21)	280 (22)	100	<3
	4-8	127	295 (15)	208 (29)	228 (26)	260 (21)	296 (19)	340 (23)	393 (35)	431 (47)	100	<3
Male	9-13	111	357 (24)	235 (24)	258 (23)	296 (23)	340 (25)	392 (32)	455 (39)	500 (45)	100	0.0 (0.0)
	14-18	107	395 (26)	240 (43) ^E	268 (40)	324 (34)	400 (31)	496 (42)	603 (67)	678 (90)	100	0.0 (0.0)
	19-30	77	316 (22)	221 (37) ^E	241 (33)	276 (29)	318 (28)	364 (33)	409 (43)	437 (51)	100	<3
	31-50	145	281 (17)	182 (27)	202 (24)	238 (21)	278 (19)	322 (21)	365 (26)	395 (32)	100	<3
	51-70	182	255 (13)	161 (28) ^E	179 (25)	210 (21)	250 (16)	293 (29)	335 (55)	362 (76) ^E	100	^F
	>70	63	267 (21)	178 (22)	196 (21)	229 (22)	271 (27)	318 (37)	366 (52)	398 (65)	100	<3
	19+	467	279 (10)	173 (13)	195 (12)	233 (11)	279 (12)	330 (14)	380 (18)	413 (21)	100	<3
Female	9-13	96	288 (12)	193 (16)	211 (16)	245 (16)	290 (16)	344 (20)	402 (28)	441 (36)	100	<3
	14-18	105	287 (20)	176 (24)	195 (24)	231 (25)	282 (28)	348 (36)	419 (49)	465 (58)	100	<3
	19-30	91	230 (13)	172 (14)	184 (15)	205 (16)	229 (17)	254 (18)	277 (20)	291 (21)	100	<3
	31-50	167	195 (10)	96 (15)	118 (14)	157 (12)	198 (12)	241 (14)	288 (20)	320 (25)	100	^F
	51-70	198	195 (11)	97 (19) ^E	116 (18)	152 (15)	195 (13)	241 (16)	281 (20)	304 (23)	100	^F
	>70	74	228 (21)	141 (24) ^E	157 (22)	185 (18)	219 (18)	257 (22)	293 (28)	315 (33)	100	<3
	19+	530	205 (7)	112 (9)	132 (8)	167 (8)	207 (8)	248 (9)	288 (12)	314 (15)	100	^F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.2 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	58	208 (14)	144 (19)	159 (18)	182 (17)	209 (18)	242 (20)	279 (24)	302 (27)	100	<3
	4-8	110	256 (9)	179 (21)	195 (18)	224 (13)	256 (11)	287 (15)	315 (22)	333 (27)	100	<3
Male	9-13	95	310 (16)	230 (31)	246 (28)	274 (25)	307 (23)	344 (26)	381 (33)	405 (40)	100	<3
	14-18	87	349 (25)	210 (37) ^E	239 (35)	295 (32)	366 (33)	449 (44)	534 (60)	590 (71)	100	<3
	19-30	70	354 (22)	206 (41) ^E	237 (37)	293 (31)	360 (27)	431 (31)	501 (41)	545 (50)	100	<3
	31-50	109	280 (17)	166 (17)	186 (17)	223 (18)	271 (22)	330 (29)	395 (37)	441 (45)	100	<3
	51-70	128	254 (12)	184 (22)	201 (19)	230 (15)	262 (16)	295 (22)	329 (30)	351 (36)	100	<3
	>70	65	223 (10)	127 (18)	147 (16)	183 (13)	225 (13)	269 (16)	311 (21)	338 (26)	100	^F
	19+	372	282 (9)	168 (12)	190 (11)	227 (11)	275 (13)	339 (17)	409 (21)	455 (25)	100	<3
Female	9-13	75	266 (15)	171 (21)	191 (23)	226 (29)	270 (42)	320 (67) ^E	374 (95) ^E	411 (111) ^E	100	<3
	14-18	81	273 (16)	168 (28)	193 (24)	235 (19)	280 (18)	324 (21)	368 (29)	399 (36)	100	<3
	19-30	101	260 (13)	199 (22)	217 (20)	246 (17)	279 (17)	313 (20)	347 (26)	370 (30)	100	<3
	31-50	116	223 (10)	131 (14)	150 (13)	184 (11)	224 (11)	265 (13)	303 (16)	326 (18)	100	<3
	51-70	146	207 (8)	139 (20)	154 (18)	178 (14)	208 (11)	239 (12)	271 (17)	292 (22)	100	<3
	>70	94	176 (12)	97 (14)	109 (14)	133 (13)	166 (14)	205 (16)	247 (22)	276 (28)	100	^F
	19+	457	219 (5)	137 (8)	155 (7)	186 (7)	223 (7)	261 (8)	297 (10)	321 (11)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.3 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	112	203 (8)	166 (17)	174 (15)	187 (12)	202 (11)	218 (14)	233 (18)	242 (21)	100	<3
	4-8	177	266 (11)	216 (21)	228 (19)	249 (16)	273 (14)	297 (16)	319 (20)	332 (23)	100	0.0 (0.0)
Male	9-13	111	338 (17)	239 (32)	260 (28)	296 (23)	339 (21)	385 (28)	428 (39)	455 (47)	100	<3
	14-18	113	345 (26)	223 (34)	244 (32)	282 (31)	331 (35)	394 (44)	463 (58)	510 (69)	100	<3
	19-30	91	344 (22)	202 (51) ^E	229 (45) ^E	276 (35)	329 (28)	384 (30)	438 (42)	473 (53)	100	<3
	31-50	101	318 (20)	189 (26)	213 (25)	257 (23)	314 (24)	378 (29)	443 (39)	486 (48)	100	<3
	51-70	134	262 (10)	189 (21)	203 (18)	228 (14)	261 (13)	297 (18)	328 (26)	347 (31)	100	<3
	>70	56	232 (13)	178 (22)	188 (20)	207 (16)	229 (15)	252 (17)	274 (24)	288 (29)	100	<3
	19+	382	298 (10)	179 (14)	200 (13)	238 (12)	289 (12)	346 (14)	402 (19)	437 (24)	100	<3
Female	9-13	105	280 (16)	182 (18)	201 (17)	233 (17)	272 (18)	315 (20)	355 (25)	381 (28)	100	<3
	14-18	120	276 (21)	132 (24) ^E	155 (25)	204 (26)	269 (25)	337 (25)	404 (32)	451 (40)	100	<3
	19-30	91	260 (18)	182 (27)	200 (26)	232 (26)	271 (26)	310 (34)	345 (45)	366 (55)	100	F
	31-50	159	230 (12)	127 (17)	150 (16)	191 (15)	236 (15)	277 (17)	315 (20)	340 (24)	100	F
	51-70	174	207 (9)	125 (21)	141 (18)	170 (13)	203 (10)	240 (12)	277 (20)	301 (26)	100	F
	>70	80	206 (11)	128 (21)	144 (20)	173 (19)	211 (19)	253 (20)	297 (25)	325 (30)	100	F
	19+	504	226 (6)	128 (9)	148 (9)	184 (8)	228 (9)	274 (10)	317 (12)	345 (14)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.4 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	99	201 (7)	150 (15)	161 (13)	180 (11)	203 (10)	228 (11)	251 (14)	266 (17)	100	<3
	4-8	140	283 (13)	221 (10)	233 (11)	255 (13)	282 (14)	311 (16)	340 (18)	357 (18)	100	0.0 (0.0)
Male	9-13	92	311 (23)	201 (31)	217 (31)	251 (29)	297 (29)	353 (34)	410 (43)	448 (50)	100	0.0 (0.0)
	14-18	107	395 (24)	272 (37)	297 (34)	342 (30)	402 (27)	477 (32)	555 (44)	606 (55)	100	<3
	19-30	73	384 (43)	214 (34)	246 (33)	305 (34)	380 (42)	473 (58)	578 (81)	644 (94)	100	<3
	31-50	134	311 (23)	171 (26)	189 (26)	225 (26)	280 (25)	353 (33)	428 (46)	476 (57)	100	<3
	51-70	131	254 (12)	184 (18)	197 (16)	220 (14)	248 (14)	280 (18)	315 (28)	338 (36)	100	<3
	>70	55	267 (22)	143 (30) ^E	165 (29) ^E	207 (30)	263 (30)	322 (30)	371 (31)	402 (34)	100	^F
	19+	393	305 (13)	165 (11)	186 (11)	227 (12)	288 (13)	371 (20)	464 (31)	530 (42)	100	<3
Female	9-13	79	285 (11)	212 (22)	230 (19)	260 (16)	297 (15)	336 (19)	374 (25)	397 (29)	100	<3
	14-18	104	265 (13)	183 (26)	201 (22)	230 (17)	264 (16)	301 (22)	340 (33)	367 (41)	100	<3
	19-30	101	228 (16)	127 (26) ^E	144 (26) ^E	180 (25)	227 (23)	281 (21)	336 (26)	374 (33)	100	^F
	31-50	143	222 (11)	147 (21)	162 (18)	189 (14)	220 (13)	254 (17)	289 (25)	311 (31)	100	<3
	51-70	193	200 (7)	129 (17)	141 (15)	162 (12)	189 (10)	220 (11)	254 (17)	277 (23)	100	<3
	>70	94	187 (9)	115 (14)	125 (13)	143 (12)	167 (11)	196 (14)	227 (21)	249 (27)	100	^F
	19+	531	212 (6)	128 (9)	144 (8)	172 (7)	205 (7)	245 (9)	287 (12)	314 (14)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.5 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004¹

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						EAR ²	% <EAR ²	(SE)
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	311	211	(8)	121 (11)	140 (11)	172 (11)	211 (11)	256 (13)	304 (18)	335 (21)	100	F
	4-8	485	270	(7)	203 (16)	218 (13)	243 (9)	274 (8)	308 (13)	341 (20)	362 (25)	100	<3
Male	9-13	277	369	(15)	239 (22)	265 (22)	317 (22)	383 (24)	454 (28)	526 (34)	575 (39)	100	<3
	14-18	339	421	(17)	259 (23)	293 (22)	355 (21)	438 (21)	529 (25)	619 (35)	688 (46)	100	0.0 (0.0)
	19-30	237	360	(14)	250 (33)	272 (29)	311 (23)	360 (20)	413 (27)	463 (40)	494 (49)	100	0.0 (0.0)
	31-50	423	319	(15)	191 (25)	215 (23)	262 (20)	320 (18)	382 (21)	441 (32)	480 (39)	100	<3
	51-70	387	269	(8)	175 (15)	194 (13)	227 (11)	266 (10)	313 (13)	358 (18)	386 (22)	100	<3
	>70	132	240	(14)	124 (28) ^E	146 (26) ^E	188 (22)	240 (20)	299 (21)	358 (25)	395 (29)	100	F
	19+	1179	307	(8)	178 (10)	201 (10)	247 (10)	305 (10)	369 (11)	433 (15)	477 (19)	100	<3
Female	9-13	281	296	(12)	182 (17)	203 (16)	242 (14)	289 (13)	341 (15)	390 (19)	422 (23)	100	<3
	14-18	321	282	(9)	190 (12)	210 (11)	246 (11)	290 (12)	339 (15)	386 (18)	416 (21)	100	<3
	19-30	249	274	(14)	178 (19)	196 (17)	229 (16)	269 (16)	313 (21)	356 (27)	383 (31)	100	<3
	31-50	364	238	(10)	156 (11)	171 (10)	196 (10)	232 (12)	281 (16)	334 (24)	375 (34)	100	<3
	51-70	467	227	(6)	132 (10)	150 (9)	181 (8)	221 (7)	269 (10)	321 (16)	357 (21)	100	<3
	>70	215	193	(7)	132 (12)	145 (12)	167 (11)	191 (10)	216 (11)	242 (13)	260 (15)	100	<3
	19+	1295	236	(5)	144 (6)	162 (5)	191 (5)	229 (6)	278 (8)	331 (12)	369 (15)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.6 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR	(SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	644	205 (6)	119 (11)	136 (9)	166 (7)	202 (6)	243 (7)	287 (12)	317 (17)	100	F	
	4-8	956	266 (4)	191 (11)	206 (9)	234 (6)	266 (5)	302 (7)	338 (11)	361 (15)	100	<3	
Male	9-13	589	326 (8)	207 (9)	228 (9)	267 (8)	317 (9)	378 (11)	449 (17)	501 (24)	100	0.0 (0.0)	
	14-18	639	363 (9)	245 (25)	270 (22)	316 (16)	371 (12)	431 (15)	491 (24)	530 (31)	100	<3	
	19-30	481	320 (10)	245 (29)	261 (24)	289 (16)	321 (12)	354 (18)	387 (30)	407 (38)	100	<3	
	31-50	709	277 (8)	152 (8)	177 (8)	220 (8)	270 (8)	328 (10)	390 (13)	432 (15)	100	<3	
	51-70	758	252 (6)	148 (12)	167 (10)	201 (8)	245 (7)	295 (9)	346 (14)	380 (19)	100	<3	
	>70	734	230 (5)	127 (9)	146 (9)	180 (7)	224 (6)	272 (7)	320 (10)	351 (13)	100	<3	
	19+	2682	276 (4)	163 (8)	184 (8)	223 (6)	271 (5)	323 (6)	378 (9)	414 (12)	100	<3	
Female	9-13	585	275 (6)	185 (14)	203 (12)	235 (9)	273 (7)	313 (10)	352 (16)	378 (20)	100	<3	
	14-18	645	287 (8)	178 (19)	200 (16)	239 (12)	287 (9)	342 (12)	399 (21)	437 (29)	100	<3	
	19-30	514	231 (7)	130 (12)	150 (11)	184 (9)	224 (8)	269 (9)	314 (13)	342 (17)	100	<3	
	31-50	758	220 (6)	113 (9)	132 (8)	167 (7)	213 (7)	264 (9)	317 (13)	351 (16)	100	F	
	51-70	955	206 (4)	124 (7)	139 (7)	167 (6)	201 (5)	239 (6)	278 (8)	304 (10)	100	<3	
	>70	1345	199 (3)	118 (5)	131 (5)	158 (4)	193 (4)	231 (5)	271 (8)	298 (10)	100	<3	
	19+	3572	216 (3)	118 (4)	135 (4)	168 (4)	209 (4)	255 (4)	302 (6)	333 (7)	100	2.0 (0.5) ^E	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.7 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	324	199 (6)	129 (16)	142 (14)	167 (10)	197 (7)	229 (10)	261 (16)	281 (21)	100	<3
	4-8	425	256 (8)	176 (15)	190 (14)	216 (11)	249 (10)	287 (13)	328 (20)	354 (26)	100	<3
Male	9-13	274	326 (11)	193 (17)	218 (15)	262 (13)	318 (13)	384 (17)	454 (26)	499 (33)	100	<3
	14-18	297	398 (18)	231 (29)	262 (26)	318 (22)	390 (21)	477 (31)	573 (50)	641 (65)	100	<3
	19-30	249	339 (19)	188 (43) ^E	216 (37) ^E	268 (27)	333 (22)	406 (31)	479 (48)	525 (62)	100	<3
	31-50	309	269 (10)	168 (21)	188 (18)	225 (14)	269 (13)	317 (16)	364 (23)	396 (30)	100	<3
	51-70	277	249 (10)	129 (18)	148 (16)	186 (14)	236 (12)	297 (14)	361 (23)	405 (31)	100	^F
	>70	136	225 (12)	129 (21)	148 (18)	181 (14)	218 (12)	262 (17)	314 (28)	350 (36)	100	^F
	19+	971	275 (7)	149 (10)	171 (9)	211 (8)	265 (8)	329 (10)	396 (14)	442 (18)	100	<3
Female	9-13	265	287 (12)	174 (17)	192 (14)	227 (11)	270 (11)	319 (16)	367 (25)	399 (32)	100	<3
	14-18	290	280 (18)	163 (24)	184 (22)	220 (20)	267 (20)	325 (24)	393 (35)	443 (45)	100	<3
	19-30	197	275 (24)	180 (29)	197 (26)	227 (22)	265 (21)	307 (27)	349 (37)	377 (46)	100	<3
	31-50	312	213 (8)	132 (15)	147 (13)	176 (11)	211 (9)	251 (12)	292 (18)	321 (24)	100	<3
	51-70	312	206 (8)	134 (17)	148 (15)	172 (12)	201 (10)	235 (12)	270 (18)	294 (23)	100	<3
	>70	239	187 (7)	111 (13)	124 (12)	150 (10)	183 (8)	222 (10)	262 (16)	288 (21)	100	^F
	19+	1060	220 (7)	127 (7)	143 (7)	173 (6)	212 (6)	258 (9)	307 (15)	341 (20)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.8 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	129	198 (9)	122 (12)	137 (11)	164 (10)	197 (11)	233 (13)	269 (16)	293 (18)	100	F
	4-8	213	265 (9)	197 (18)	212 (15)	236 (10)	264 (8)	294 (14)	325 (23)	347 (29)	100	<3
Male	9-13	122	332 (19)	241 (26)	261 (25)	298 (22)	343 (22)	394 (27)	444 (36)	476 (45)	100	0.0 (0.0)
	14-18	150	394 (29)	281 (40)	305 (36)	346 (30)	395 (29)	450 (37)	505 (52)	541 (64)	100	<3
	19-30	106	333 (26)	167 (32) ^E	195 (31)	249 (28)	317 (28)	396 (34)	475 (46)	527 (56)	100	<3
	31-50	155	303 (22)	150 (31) ^E	179 (28)	229 (24)	289 (23)	362 (29)	449 (47)	516 (66)	100	<3
	51-70	122	232 (12)	137 (23) ^E	157 (20)	193 (16)	236 (16)	281 (22)	320 (30)	343 (36)	100	F
	>70	88	240 (12)	152 (15)	167 (14)	196 (14)	235 (15)	280 (19)	329 (27)	361 (34)	100	<3
	19+	471	283 (11)	139 (11)	166 (10)	214 (10)	273 (11)	346 (15)	430 (24)	492 (34)	100	<3
Female	9-13	103	289 (16)	202 (21)	219 (20)	251 (18)	289 (19)	333 (24)	378 (32)	407 (38)	100	<3
	14-18	142	298 (15)	202 (26)	226 (25)	267 (24)	314 (24)	366 (26)	421 (30)	459 (36)	100	<3
	19-30	111	242 (12)	149 (15)	165 (14)	194 (12)	233 (13)	278 (18)	325 (28)	356 (36)	100	<3
	31-50	146	221 (11)	158 (23)	169 (20)	189 (17)	214 (14)	243 (15)	272 (21)	291 (27)	100	<3
	51-70	184	217 (14)	151 (24)	165 (22)	189 (18)	217 (15)	248 (16)	278 (21)	298 (26)	100	<3
	>70	143	204 (8)	136 (10)	151 (9)	175 (9)	204 (11)	239 (14)	276 (18)	300 (21)	100	<3
	19+	584	221 (6)	141 (9)	156 (8)	182 (8)	216 (8)	257 (8)	299 (11)	328 (13)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.9 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	169	188 (7)	113 (15)	128 (13)	155 (10)	187 (9)	220 (10)	255 (16)	280 (22)	100	F
	4-8	281	255 (8)	226 (32)	231 (27)	241 (17)	253 (9)	264 (17)	275 (32)	282 (42)	100	<3
Male	9-13	183	339 (16)	214 (25)	236 (23)	278 (20)	333 (18)	399 (45)	470 (93) ^E	518 (141) ^E	100	<3
	14-18	187	372 (14)	234 (33)	263 (28)	316 (21)	378 (17)	446 (24)	516 (37)	561 (48)	100	<3
	19-30	223	322 (13)	197 (26)	220 (24)	263 (19)	316 (16)	377 (21)	439 (32)	480 (41)	100	<3
	31-50	229	299 (19)	171 (32) ^E	195 (29)	241 (26)	301 (24)	378 (43)	460 (81) ^E	511 (114) ^E	100	F
	51-70	197	245 (9)	161 (26)	179 (22)	208 (14)	243 (10)	282 (16)	321 (28)	347 (36)	100	<3
	>70	72	222 (13)	157 (17)	170 (16)	196 (14)	227 (13)	262 (17)	297 (26)	321 (34)	100	<3
	19+	721	286 (9)	170 (15)	192 (13)	232 (11)	285 (11)	349 (13)	416 (19)	460 (24)	100	<3
Female	9-13	165	286 (13)	259 (32)	269 (29)	285 (23)	304 (19)	324 (24)	343 (35)	354 (44)	100	0.0 (0.0)
	14-18	206	259 (11)	187 (23)	200 (20)	223 (16)	252 (15)	289 (18)	328 (25)	351 (31)	100	<3
	19-30	191	245 (11)	121 (16)	144 (15)	186 (13)	238 (13)	299 (17)	366 (24)	415 (31)	100	F
	31-50	258	204 (9)	126 (20)	141 (18)	169 (15)	204 (12)	244 (18)	285 (24)	312 (29)	100	F
	51-70	249	199 (7)	143 (15)	155 (13)	176 (10)	200 (8)	227 (10)	252 (15)	268 (20)	100	<3
	>70	128	179 (9)	103 (11)	115 (11)	140 (11)	173 (12)	214 (14)	254 (18)	282 (22)	100	F
	19+	826	209 (5)	118 (6)	135 (6)	167 (6)	207 (6)	254 (8)	304 (11)	338 (14)	100	F

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.10 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004¹

Sex	Age (years)	n Mean (SE)		Percentiles (and SE) of usual intake						EAR ²	% <EAR (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both	1-3	192	197 (8)	134 (15)	145 (14)	166 (12)	192 (12)	220 (14)	247 (18)	264 (22)	100	<3
	4-8	321	257 (7)	171 (12)	187 (11)	217 (9)	253 (9)	295 (12)	340 (19)	371 (25)	100	<3
Male	9-13	226	319 (11)	213 (15)	234 (14)	272 (13)	321 (13)	379 (17)	443 (26)	489 (34)	100	0.0 (0.0)
	14-18	262	383 (19)	241 (29)	269 (27)	319 (24)	383 (23)	458 (28)	537 (39)	591 (48)	100	0.0 (0.0)
	19-30	197	368 (18)	228 (23)	255 (22)	304 (20)	366 (22)	437 (28)	509 (39)	555 (46)	100	<3
	31-50	282	347 (16)	187 (24)	220 (22)	278 (18)	345 (17)	423 (24)	494 (35)	535 (44)	100	<3
	51-70	234	276 (12)	147 (15)	171 (14)	214 (12)	268 (14)	331 (19)	398 (26)	444 (32)	100	<3
	>70	119	251 (13)	156 (19)	174 (17)	207 (15)	247 (16)	295 (23)	340 (30)	367 (33)	100	<3
	19+	832	322 (8)	175 (8)	201 (8)	251 (8)	317 (9)	393 (12)	467 (16)	516 (20)	100	<3
Female	9-13	226	288 (13)	188 (14)	206 (14)	240 (14)	288 (15)	347 (20)	413 (27)	459 (33)	100	<3
	14-18	242	273 (12)	141 (19)	167 (17)	213 (15)	268 (15)	329 (18)	391 (22)	433 (25)	100	<3
	19-30	208	242 (10)	159 (18)	177 (17)	207 (14)	242 (13)	283 (16)	326 (21)	355 (26)	100	<3
	31-50	263	239 (10)	133 (17)	153 (15)	188 (12)	231 (12)	279 (16)	325 (22)	352 (26)	100	<3
	51-70	322	209 (12)	110 (15)	130 (14)	166 (13)	206 (13)	254 (15)	310 (20)	348 (25)	100	^F
	>70	198	204 (9)	127 (11)	142 (11)	169 (11)	202 (11)	237 (14)	271 (17)	292 (19)	100	<3
	19+	991	226 (5)	126 (7)	145 (7)	178 (6)	221 (6)	270 (8)	320 (10)	352 (12)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.11 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR (SE)
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	348	206 (5)	150 (9)	162 (8)	182 (7)	205 (7)	232 (8)	257 (10)	273 (11)	100	<3
	4-8	554	277 (7)	212 (13)	226 (12)	251 (10)	281 (9)	315 (11)	349 (15)	371 (18)	100	0.0 (0.0)
Male	9-13	409	331 (11)	223 (18)	245 (16)	282 (14)	324 (13)	376 (16)	428 (22)	461 (26)	100	0.0 (0.0)
	14-18	414	372 (14)	229 (17)	256 (17)	306 (17)	373 (18)	457 (22)	549 (28)	610 (34)	100	<3
	19-30	311	351 (17)	203 (22)	232 (20)	285 (18)	348 (19)	415 (22)	488 (30)	539 (38)	100	<3
	31-50	489	305 (11)	172 (11)	194 (11)	236 (12)	291 (13)	358 (16)	429 (21)	475 (25)	100	<3
	51-70	575	257 (7)	180 (13)	194 (11)	219 (9)	254 (9)	294 (11)	331 (16)	354 (20)	100	<3
	>70	239	250 (10)	145 (12)	165 (12)	201 (12)	248 (12)	299 (14)	350 (17)	381 (19)	100	<3
	19+	1614	295 (6)	168 (6)	190 (6)	231 (6)	285 (7)	351 (9)	418 (12)	464 (15)	100	<3
Female	9-13	355	283 (7)	201 (11)	219 (11)	251 (10)	289 (10)	332 (11)	376 (15)	406 (18)	100	0.0 (0.0)
	14-18	410	275 (10)	152 (14)	175 (13)	218 (12)	271 (12)	330 (13)	391 (18)	433 (23)	100	<3
	19-30	384	243 (9)	169 (14)	185 (14)	215 (13)	251 (13)	288 (14)	323 (16)	345 (18)	100	<3
	31-50	585	219 (7)	117 (9)	139 (9)	179 (8)	225 (9)	270 (9)	313 (11)	341 (12)	100	F
	51-70	711	202 (5)	120 (8)	137 (7)	165 (6)	197 (6)	233 (7)	269 (9)	292 (11)	100	F
	>70	342	202 (7)	118 (9)	132 (9)	159 (9)	195 (10)	237 (12)	283 (16)	314 (21)	100	F
	19+	2022	216 (4)	126 (5)	145 (5)	178 (5)	217 (5)	260 (6)	301 (6)	328 (7)	100	1.3 (0.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.12 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004¹

Sex	Age (years)	Percentiles (and SE) of usual intake									EAR ²	% <EAR	(SE)							
		n	Mean	(SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)		95th (SE)								
Both	1-3	622	192	(5)	116	(8)	131	(7)	158	(7)	191	(6)	226	(7)	262	(9)	287	(12)	100	F
	4-8	919	257	(5)	188	(14)	201	(11)	225	(8)	253	(6)	283	(9)	314	(15)	334	(19)	100	0.0 (0.0)
Male	9-13	579	335	(10)	224	(13)	245	(12)	285	(11)	335	(12)	393	(16)	453	(22)	493	(27)	100	0.0 (0.0)
	14-18	634	381	(11)	247	(19)	275	(17)	325	(14)	386	(13)	453	(16)	522	(23)	569	(29)	100	<3
	19-30	578	327	(11)	178	(16)	205	(15)	254	(14)	318	(13)	391	(15)	465	(21)	514	(27)	100	<3
	31-50	693	293	(13)	174	(21)	196	(19)	239	(16)	294	(16)	359	(20)	430	(28)	477	(35)	100	<3
	51-70	596	243	(6)	142	(11)	163	(9)	199	(7)	241	(7)	288	(10)	339	(14)	374	(18)	100	<3
	>70	296	227	(8)	139	(10)	155	(9)	187	(9)	226	(8)	268	(10)	313	(15)	345	(19)	100	<3
	19+	2163	283	(6)	156	(7)	179	(7)	223	(6)	279	(7)	345	(9)	418	(11)	468	(14)	100	<3
Female	9-13	533	287	(9)	196	(14)	215	(14)	249	(13)	292	(12)	337	(14)	382	(17)	410	(20)	100	0.0 (0.0)
	14-18	638	271	(8)	168	(11)	189	(10)	223	(10)	266	(10)	320	(12)	377	(16)	415	(20)	100	<3
	19-30	499	251	(9)	134	(11)	155	(10)	193	(9)	241	(9)	297	(12)	358	(17)	400	(22)	100	<3
	31-50	716	209	(6)	135	(12)	149	(11)	176	(9)	209	(8)	246	(9)	284	(13)	309	(16)	100	<3
	51-70	745	204	(5)	134	(9)	149	(8)	173	(7)	203	(6)	237	(7)	270	(9)	292	(11)	100	<3
	>70	510	187	(5)	106	(6)	121	(6)	148	(6)	183	(7)	226	(9)	270	(10)	299	(12)	100	F
	19+	2470	214	(4)	123	(4)	140	(4)	171	(4)	210	(4)	256	(5)	305	(7)	338	(9)	100	1.4 (0.4) ^E

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 35.13 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004¹

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							EAR ²	% <EAR (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	2117	203 (3)	124 (5)	139 (4)	167 (4)	201 (4)	240 (5)	280 (7)	307 (9)	100	1.2 (0.4) ^E
	4-8	3235	265 (3)	186 (5)	202 (4)	230 (3)	265 (3)	304 (5)	342 (7)	367 (9)	100	0.0 (0.0)
Male	9-13	2080	337 (5)	225 (7)	246 (7)	286 (7)	337 (7)	395 (8)	456 (11)	497 (13)	100	0.0 (0.0)
	14-18	2288	382 (6)	238 (9)	267 (8)	321 (8)	389 (7)	465 (9)	542 (12)	596 (16)	100	0.0 (0.0)
	19-30	1804	339 (6)	205 (9)	231 (8)	277 (8)	335 (8)	399 (10)	464 (13)	507 (15)	100	<3
	31-50	2596	301 (6)	164 (7)	190 (7)	236 (6)	294 (7)	363 (8)	434 (11)	481 (13)	100	<3
	51-70	2550	259 (4)	157 (5)	176 (5)	210 (4)	254 (4)	303 (5)	353 (8)	385 (10)	100	<3
	>70	1520	236 (5)	138 (6)	156 (6)	191 (5)	232 (6)	280 (6)	330 (9)	363 (10)	100	<3
	19+	8470	292 (3)	162 (4)	186 (3)	229 (3)	285 (4)	350 (4)	418 (6)	464 (8)	100	0.4 (0.1) ^E
Female	9-13	1980	284 (4)	182 (6)	202 (5)	238 (5)	282 (5)	330 (6)	379 (7)	411 (9)	100	<3
	14-18	2256	280 (4)	163 (5)	187 (5)	230 (5)	281 (5)	341 (6)	402 (8)	443 (10)	100	<3
	19-30	1854	247 (5)	146 (6)	166 (5)	200 (5)	241 (5)	287 (7)	335 (9)	366 (11)	100	<3
	31-50	2686	225 (4)	125 (4)	143 (4)	177 (4)	218 (5)	269 (5)	320 (8)	357 (11)	100	1.5 (0.4) ^E
	51-70	3200	211 (3)	125 (4)	142 (4)	171 (3)	208 (4)	249 (4)	293 (6)	322 (7)	100	1.3 (0.4) ^E
	>70	2610	197 (3)	119 (4)	133 (4)	160 (4)	192 (4)	230 (4)	268 (5)	293 (6)	100	1.5 (0.4) ^E
	19+	10350	222 (2)	126 (2)	144 (2)	176 (2)	216 (2)	263 (3)	312 (4)	345 (6)	100	1.3 (0.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

36. Totat fats (g/d): Usual intakes from food

Table 36.1 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	79	57 (5)	36 (6) ^E	40 (6)	47 (6)	55 (6)	64 (7)	72 (9)	77 (10)
	4-8	127	78 (5)	55 (8)	60 (7)	68 (7)	79 (7)	91 (9)	103 (11)	111 (14)
Male										
	9-13	111	97 (6)	66 (9)	72 (9)	83 (8)	97 (8)	113 (10)	131 (15)	142 (18)
	14-18	107	94 (6)	65 (11) ^E	70 (10)	80 (9)	93 (7)	109 (8)	127 (13)	139 (17)
	19-30	77	94 (7)	60 (8)	67 (8)	80 (8)	94 (8)	109 (9)	123 (10)	132 (11)
	31-50	145	91 (8)	64 (9)	70 (8)	79 (7)	89 (7)	99 (9)	111 (11)	118 (13)
	51-70	182	70 (4)	40 (8) ^E	44 (8) ^E	53 (6)	64 (5)	77 (6)	91 (9)	101 (12)
	>70	63	71 (5)	49 (10) ^E	55 (9)	65 (7)	77 (7)	89 (8)	101 (10)	107 (11)
	19+	467	84 (4)	51 (5)	58 (5)	70 (4)	83 (4)	97 (5)	112 (7)	121 (8)
Female										
	9-13	96	72 (3)	45 (6)	50 (6)	59 (5)	71 (4)	84 (5)	99 (8)	109 (11)
	14-18	105	77 (6)	49 (10) ^E	54 (10) ^E	62 (9)	73 (8)	87 (10)	101 (14)	111 (17)
	19-30	91	70 (8)	45 (6)	50 (6)	57 (7)	67 (8)	77 (9)	89 (11)	97 (12)
	31-50	167	65 (4)	33 (6) ^E	38 (6)	48 (6)	62 (6)	78 (7)	95 (8)	106 (10)
	51-70	198	54 (4)	26 (6) ^E	31 (6) ^E	41 (5)	54 (4)	69 (7)	84 (11)	94 (13)
	>70	74	58 (7)	38 (7) ^E	42 (7)	49 (7)	58 (8)	69 (9)	80 (11)	86 (13)
	19+	530	62 (3)	35 (3)	39 (3)	48 (3)	60 (3)	74 (4)	88 (5)	98 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.2 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	50 (3)	37 (5)	39 (5)	44 (4)	49 (4)	54 (5)	59 (6)	63 (7)
	4-8	110	66 (4)	50 (4)	54 (5)	59 (5)	66 (5)	73 (6)	79 (6)	82 (6)
	9-13	95	80 (5)	57 (6)	62 (6)	71 (7)	83 (7)	95 (7)	108 (8)	117 (8)
Male	14-18	87	105 (7)	70 (13) ^E	77 (12)	91 (10)	108 (9)	127 (12)	147 (18)	160 (23)
	19-30	70	129 (15)	103 (19) ^E	111 (18)	126 (17)	144 (18)	165 (23)	186 (31)	198 (36) ^E
	31-50	109	93 (5)	63 (11) ^E	69 (9)	81 (7)	96 (6)	113 (8)	129 (12)	140 (15)
	51-70	128	76 (3)	49 (9) ^E	55 (9)	66 (7)	79 (5)	92 (5)	103 (7)	111 (8)
	>70	65	65 (4)	41 (7) ^E	45 (7)	52 (6)	62 (5)	73 (7)	86 (11)	94 (14)
	19+	372	92 (4)	57 (5)	64 (5)	77 (5)	95 (5)	117 (7)	139 (10)	154 (13)
	9-13	75	71 (6)	45 (9) ^E	50 (8)	60 (8)	72 (8)	87 (10)	102 (13)	113 (16)
	14-18	81	73 (5)	40 (7) ^E	47 (7)	60 (7)	75 (7)	91 (8)	107 (10)	117 (12)
Female	19-30	101	67 (5)	42 (6)	47 (5)	57 (5)	71 (6)	86 (9)	102 (12)	112 (15)
	31-50	116	64 (4)	41 (8) ^E	45 (7)	54 (6)	64 (5)	75 (6)	86 (8)	92 (10)
	51-70	146	61 (3)	38 (6)	43 (5)	50 (4)	60 (3)	70 (4)	80 (6)	87 (8)
	>70	94	54 (4)	36 (5)	39 (5)	45 (4)	53 (4)	62 (6)	72 (9)	79 (11)
	19+	457	62 (2)	38 (3)	42 (3)	51 (3)	63 (3)	76 (4)	89 (5)	97 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.3 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	50 (3)	35 (5)	38 (5)	43 (4)	49 (3)	56 (4)	64 (6)	69 (8)
	4-8	177	64 (3)	46 (3)	50 (3)	57 (3)	64 (4)	72 (4)	80 (4)	84 (4)
Male	9-13	111	86 (5)	61 (8)	65 (7)	74 (6)	84 (6)	95 (7)	106 (9)	113 (11)
	14-18	113	101 (8)	66 (12) ^E	72 (12)	83 (12)	97 (13)	116 (14)	138 (17)	152 (20)
	19-30	91	101 (9)	57 (8)	64 (8)	78 (9)	98 (11)	121 (14)	146 (17)	163 (19)
	31-50	101	101 (6)	60 (10) ^E	67 (10)	81 (8)	98 (8)	116 (10)	134 (15)	146 (18)
	51-70	134	82 (7)	46 (12) ^E	52 (11) ^E	64 (9)	78 (8)	96 (11)	116 (18)	127 (22) ^E
	>70	56	63 (5)	32 (6) ^E	37 (7) ^E	48 (6)	62 (6)	75 (9)	87 (12)	96 (15)
	19+	382	92 (4)	50 (7)	58 (6)	71 (5)	87 (5)	107 (6)	129 (10)	143 (12)
Female	9-13	105	71 (5)	49 (5)	53 (5)	61 (5)	71 (6)	82 (7)	94 (8)	102 (9)
	14-18	120	65 (7)	36 (12) ^E	42 (11) ^E	52 (9) ^E	64 (8)	78 (8)	91 (11)	99 (14)
	19-30	91	77 (7)	39 (7) ^E	47 (6)	61 (6)	76 (6)	94 (9)	119 (15)	139 (19)
	31-50	159	78 (6)	42 (8) ^E	49 (8)	63 (7)	79 (7)	95 (9)	111 (12)	121 (14)
	51-70	174	67 (6)	42 (8) ^E	46 (7)	54 (6)	63 (6)	74 (7)	84 (10)	91 (13)
	>70	80	58 (8)	41 (10) ^E	45 (9) ^E	51 (9) ^E	60 (9)	70 (10)	79 (12)	84 (13)
	19+	504	72 (4)	37 (4)	43 (4)	56 (4)	72 (4)	91 (6)	111 (8)	125 (9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.4 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	57 (3)	42 (6)	45 (6)	51 (5)	57 (5)	64 (5)	70 (6)	74 (7)
	4-8	140	69 (3)	52 (6)	56 (5)	63 (4)	71 (3)	79 (4)	87 (6)	92 (8)
Male	9-13	92	103 (16)	53 (16) ^E	63 (15) ^E	75 (14) ^E	95 (15)	123 (19)	148 (24)	162 (27)
	14-18	107	113 (9)	64 (12) ^E	73 (12)	90 (11)	114 (11)	142 (13)	172 (18)	193 (22)
	19-30	73	106 (8)	60 (11) ^E	70 (10)	87 (9)	107 (11)	134 (16)	167 (25)	193 (34) ^E
	31-50	134	94 (6)	54 (10) ^E	61 (9)	74 (8)	91 (7)	110 (9)	130 (13)	142 (16)
	51-70	131	84 (6)	57 (4)	62 (5)	71 (5)	81 (5)	92 (6)	105 (8)	113 (9)
	>70	55	74 (7)	43 (9) ^E	48 (9) ^E	58 (8)	71 (8)	87 (9)	106 (14)	119 (18)
	19+	393	92 (3)	55 (5)	62 (4)	74 (4)	91 (4)	111 (5)	133 (8)	148 (10)
Female	9-13	79	72 (8)	42 (12) ^E	48 (11) ^E	59 (10) ^E	73 (10)	88 (10)	104 (12)	114 (14)
	14-18	104	66 (4)	46 (5)	50 (4)	58 (5)	67 (5)	78 (7)	88 (8)	95 (9)
	19-30	101	63 (5)	38 (10) ^E	43 (9) ^E	54 (8)	66 (6)	80 (6)	94 (8)	103 (10)
	31-50	143	58 (3)	34 (3)	38 (3)	46 (3)	56 (4)	68 (5)	81 (6)	89 (6)
	51-70	193	58 (3)	41 (7)	44 (6)	50 (5)	56 (4)	63 (4)	70 (6)	74 (7)
	>70	94	50 (3)	29 (3)	32 (3)	37 (3)	45 (4)	55 (5)	65 (8)	72 (10)
	19+	531	58 (2)	35 (3)	39 (3)	47 (3)	57 (2)	68 (3)	79 (4)	87 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.5 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	311	57	(2)	34	(4)	39	(3)	48	(3)	58	(3)	70	(4)	82	(5)	89	(6)
	4-8	485	74	(3)	45	(4)	51	(4)	60	(3)	72	(3)	87	(4)	101	(6)	111	(7)
Male	9-13	277	101	(5)	65	(7)	72	(7)	86	(7)	104	(7)	125	(10)	146	(13)	160	(15)
	14-18	339	117	(5)	76	(9)	84	(8)	100	(7)	120	(6)	146	(8)	174	(13)	194	(18)
	19-30	237	101	(4)	74	(9)	80	(8)	89	(7)	101	(7)	114	(8)	127	(10)	135	(12)
	31-50	423	100	(6)	53	(8)	61	(8)	76	(7)	97	(7)	121	(9)	145	(13)	162	(16)
	51-70	387	88	(4)	46	(6)	53	(6)	66	(5)	84	(5)	104	(6)	126	(8)	142	(10)
	>70	132	73	(5)	35	(8) ^E	42	(8) ^E	56	(8)	73	(8)	93	(9)	112	(11)	124	(13)
	19+	1179	94	(3)	51	(3)	59	(3)	73	(3)	92	(4)	114	(5)	137	(6)	153	(7)
	Female	9-13	281	80	(4)	49	(5)	55	(5)	65	(5)	78	(5)	93	(6)	107	(7)	117
14-18	321	81	(4)	54	(7)	60	(7)	71	(6)	85	(6)	99	(8)	113	(10)	122	(12)	
19-30	249	76	(5)	42	(4)	49	(4)	59	(4)	73	(5)	88	(7)	104	(8)	114	(9)	
31-50	364	75	(4)	49	(7)	54	(6)	63	(6)	76	(5)	92	(6)	108	(9)	119	(12)	
51-70	467	68	(3)	39	(5)	45	(5)	55	(4)	68	(3)	82	(4)	96	(6)	105	(7)	
>70	215	54	(3)	37	(5)	40	(4)	46	(4)	53	(4)	62	(5)	71	(6)	76	(7)	
19+	1295	71	(2)	44	(3)	49	(3)	58	(2)	70	(2)	85	(3)	100	(4)	110	(6)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.6 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	644	49 (1)	26 (2)	31 (2)	38 (2)	48 (2)	59 (2)	71 (3)	78 (4)
	4-8	956	62 (1)	45 (4)	48 (4)	55 (3)	63 (2)	71 (2)	79 (4)	85 (5)
Male										
	9-13	589	80 (2)	53 (5)	58 (5)	68 (3)	80 (2)	93 (3)	107 (5)	116 (7)
	14-18	639	96 (3)	56 (6)	64 (5)	78 (4)	96 (3)	116 (5)	137 (7)	151 (9)
	19-30	481	93 (4)	50 (7)	57 (7)	70 (5)	88 (5)	110 (7)	135 (11)	152 (15)
	31-50	709	83 (3)	48 (9) ^E	54 (7)	66 (5)	82 (4)	100 (6)	120 (10)	133 (14)
	51-70	758	78 (3)	39 (6)	45 (5)	56 (4)	72 (4)	93 (4)	116 (6)	132 (9)
	>70	734	61 (2)	29 (2)	34 (2)	45 (2)	58 (2)	74 (3)	92 (4)	104 (5)
	19+	2682	82 (2)	40 (3)	46 (3)	60 (2)	78 (2)	100 (3)	126 (5)	143 (6)
Female										
	9-13	585	69 (2)	41 (4)	46 (4)	55 (3)	67 (2)	81 (3)	95 (5)	105 (7)
	14-18	645	72 (2)	49 (7)	54 (6)	62 (4)	73 (3)	85 (4)	97 (8)	105 (10)
	19-30	514	60 (2)	31 (4)	36 (4)	45 (3)	57 (3)	70 (4)	84 (5)	93 (7)
	31-50	758	66 (2)	38 (5)	43 (4)	52 (3)	64 (3)	78 (4)	92 (6)	101 (8)
	51-70	955	60 (2)	32 (4)	37 (4)	46 (3)	58 (2)	72 (2)	86 (4)	96 (5)
	>70	1345	54 (1)	31 (3)	35 (3)	42 (2)	52 (1)	64 (2)	76 (4)	85 (6)
	19+	3572	62 (1)	34 (2)	39 (2)	48 (2)	60 (1)	73 (2)	87 (3)	97 (3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.7 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	324	47	(2)	29	(3)	32	(3)	39	(2)	46	(2)	55	(3)	63	(4)	69	(5)
	4-8	425	62	(2)	41	(6)	45	(5)	52	(4)	61	(2)	71	(4)	82	(7)	89	(9)
Male	9-13	274	81	(3)	47	(7)	54	(6)	66	(4)	81	(3)	98	(5)	115	(8)	127	(10)
	14-18	297	113	(6)	54	(8)	64	(8)	85	(7)	111	(7)	142	(9)	177	(13)	202	(18)
	19-30	249	104	(6)	60	(5)	68	(5)	83	(6)	103	(7)	124	(9)	146	(10)	160	(12)
	31-50	309	83	(4)	44	(9) ^E	49	(9) ^E	62	(8)	81	(5)	101	(8)	123	(13)	135	(17)
	51-70	277	76	(4)	43	(9) ^E	49	(9) ^E	61	(7)	77	(6)	93	(6)	109	(10)	119	(13)
	>70	136	60	(4)	30	(5)	35	(5)	43	(4)	55	(4)	69	(5)	84	(7)	94	(10)
	19+	971	83	(3)	46	(7)	52	(7)	65	(5)	82	(3)	100	(5)	120	(8)	132	(11)
	Female	9-13	265	66	(2)	43	(6)	47	(5)	54	(4)	63	(3)	72	(3)	81	(5)	86
14-18	290	72	(5)	41	(3)	46	(3)	57	(4)	72	(5)	88	(6)	106	(7)	118	(9)	
19-30	197	69	(6)	41	(9) ^E	46	(8) ^E	55	(7)	67	(6)	81	(7)	94	(11)	103	(14)	
31-50	312	67	(3)	50	(7)	53	(6)	60	(5)	67	(4)	76	(5)	84	(8)	89	(11)	
51-70	312	59	(2)	41	(6)	45	(5)	51	(3)	59	(3)	67	(4)	75	(7)	80	(9)	
>70	239	47	(2)	31	(5)	34	(4)	39	(3)	46	(3)	54	(3)	62	(5)	67	(7)	
19+	1060	62	(2)	39	(4)	43	(3)	51	(3)	61	(2)	73	(3)	86	(6)	94	(7)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.8 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	53 (3)	33 (5)	37 (5)	45 (4)	54 (4)	65 (5)	76 (6)	83 (8)
	4-8	213	71 (4)	52 (6)	56 (6)	62 (5)	71 (4)	80 (5)	89 (7)	95 (9)
	9-13	122	90 (9)	65 (9)	70 (9)	79 (9)	91 (9)	104 (12)	117 (15)	126 (18)
Male	14-18	150	105 (6)	79 (12)	85 (10)	94 (8)	105 (8)	117 (10)	129 (14)	137 (17)
	19-30	106	106 (8)	53 (14) ^E	63 (13) ^E	82 (11)	105 (10)	132 (12)	160 (18)	178 (24)
	31-50	155	98 (7)	53 (11) ^E	62 (10)	78 (8)	97 (8)	120 (11)	144 (18)	161 (23)
	51-70	122	87 (6)	48 (11) ^E	56 (9)	71 (8)	89 (7)	113 (12)	137 (19)	152 (24)
	>70	88	77 (5)	43 (8) ^E	49 (7)	61 (6)	75 (6)	91 (7)	107 (10)	118 (12)
	19+	471	95 (4)	46 (5)	56 (5)	72 (4)	94 (5)	119 (6)	147 (9)	166 (11)
	9-13	103	74 (5)	53 (9) ^E	57 (9)	65 (8)	76 (7)	87 (8)	98 (9)	105 (11)
	14-18	142	71 (4)	43 (4)	49 (4)	59 (4)	70 (5)	84 (7)	97 (9)	107 (11)
Female	19-30	111	68 (4)	43 (9) ^E	48 (8) ^E	56 (7)	67 (6)	79 (8)	90 (11)	98 (14)
	31-50	146	65 (3)	37 (7) ^E	42 (6)	50 (5)	60 (4)	72 (5)	84 (7)	91 (9)
	51-70	184	67 (3)	45 (7)	49 (6)	57 (5)	67 (4)	79 (6)	90 (9)	97 (12)
	>70	143	55 (4)	31 (7) ^E	35 (7) ^E	43 (6)	55 (5)	70 (7)	84 (10)	94 (12)
	19+	584	65 (2)	37 (3)	42 (3)	51 (3)	63 (3)	77 (3)	90 (4)	99 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.9 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	169	46 (2)	29 (4)	32 (4)	38 (3)	45 (3)	53 (3)	61 (5)	66 (6)
	4-8	281	63 (2)	41 (2)	45 (2)	52 (2)	61 (3)	71 (3)	81 (4)	88 (4)
Male	9-13	183	87 (5)	59 (4)	64 (4)	73 (5)	86 (5)	101 (7)	117 (8)	128 (9)
	14-18	187	101 (5)	65 (13) ^E	72 (12)	85 (9)	101 (6)	119 (8)	138 (14)	150 (18)
	19-30	223	89 (4)	49 (9) ^E	56 (8)	69 (6)	86 (5)	106 (7)	127 (11)	140 (14)
	31-50	229	90 (6)	46 (5)	54 (5)	68 (5)	89 (6)	112 (8)	133 (10)	147 (12)
	51-70	197	75 (5)	48 (12) ^E	54 (11) ^E	64 (8)	76 (6)	89 (8)	101 (14)	109 (19) ^E
	>70	72	67 (6)	39 (10) ^E	44 (10) ^E	54 (8)	66 (7)	79 (8)	90 (11)	97 (13)
	19+	721	84 (3)	44 (5)	52 (5)	66 (4)	83 (3)	104 (4)	125 (6)	138 (8)
Female	9-13	165	66 (4)	40 (8) ^E	45 (7)	55 (6)	66 (5)	79 (6)	92 (9)	101 (12)
	14-18	206	67 (3)	39 (8) ^E	44 (7)	53 (5)	64 (4)	77 (6)	92 (10)	102 (13)
	19-30	191	69 (5)	36 (8) ^E	42 (7) ^E	52 (7)	67 (7)	85 (9)	103 (11)	115 (14)
	31-50	258	67 (4)	49 (4)	53 (4)	60 (5)	68 (6)	78 (7)	88 (8)	94 (8)
	51-70	249	63 (3)	36 (3)	41 (3)	50 (3)	62 (4)	75 (5)	90 (6)	100 (7)
	>70	128	48 (3)	24 (4)	28 (4)	37 (4)	46 (4)	58 (5)	72 (6)	81 (8)
	19+	826	65 (2)	36 (4)	41 (3)	51 (3)	64 (3)	79 (3)	95 (5)	106 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.10 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	192	52 (3)	35 (5)	38 (5)	44 (4)	51 (3)	58 (4)	66 (6)	71 (7)
	4-8	321	63 (2)	43 (5)	47 (5)	54 (4)	63 (3)	73 (4)	83 (6)	89 (7)
Male										
	9-13	226	88 (4)	50 (6)	58 (6)	72 (5)	89 (5)	108 (5)	129 (8)	144 (11)
	14-18	262	108 (6)	64 (5)	73 (5)	89 (6)	108 (7)	130 (9)	153 (11)	169 (13)
	19-30	197	105 (5)	72 (10)	78 (9)	90 (7)	105 (6)	122 (9)	140 (14)	152 (17)
	31-50	282	105 (5)	60 (9)	69 (8)	82 (6)	98 (7)	124 (10)	152 (15)	168 (18)
	51-70	234	89 (5)	49 (7)	56 (7)	69 (6)	86 (6)	106 (7)	127 (11)	142 (14)
	>70	119	70 (5)	35 (5)	40 (5)	52 (5)	66 (6)	84 (7)	104 (9)	117 (11)
	19+	832	97 (3)	54 (3)	62 (3)	76 (3)	95 (3)	117 (4)	141 (6)	157 (8)
Female										
	9-13	226	70 (4)	44 (4)	49 (4)	58 (4)	70 (4)	84 (5)	98 (7)	108 (8)
	14-18	242	68 (3)	32 (5)	39 (4)	50 (4)	65 (4)	82 (5)	101 (6)	114 (8)
	19-30	208	68 (4)	38 (5)	44 (5)	54 (5)	67 (5)	81 (6)	96 (7)	107 (9)
	31-50	263	72 (4)	42 (5)	48 (5)	58 (4)	72 (5)	89 (7)	107 (10)	119 (13)
	51-70	322	58 (4)	29 (4)	34 (4)	43 (4)	56 (4)	71 (5)	88 (8)	100 (10)
	>70	198	54 (3)	28 (4)	32 (4)	40 (4)	50 (4)	63 (5)	75 (7)	83 (8)
	19+	991	65 (2)	33 (2)	38 (2)	49 (2)	64 (2)	81 (3)	100 (4)	112 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.11 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	348	54 (2)	35 (3)	38 (3)	45 (3)	53 (3)	63 (3)	72 (4)	78 (4)
	4-8	554	69 (2)	52 (5)	56 (4)	63 (3)	71 (3)	79 (3)	87 (5)	92 (6)
Male	9-13	409	93 (6)	60 (6)	67 (6)	78 (6)	92 (6)	109 (8)	125 (10)	135 (12)
	14-18	414	103 (4)	64 (5)	72 (5)	84 (6)	103 (6)	126 (7)	149 (9)	164 (10)
	19-30	311	103 (5)	63 (6)	70 (6)	85 (5)	103 (6)	125 (8)	148 (12)	164 (15)
	31-50	489	96 (4)	57 (4)	64 (4)	77 (4)	93 (4)	111 (5)	130 (7)	143 (9)
	51-70	575	79 (4)	50 (6)	55 (5)	65 (4)	77 (4)	91 (6)	104 (8)	112 (10)
	>70	239	68 (3)	43 (5)	49 (5)	58 (4)	69 (4)	81 (5)	93 (7)	101 (8)
	19+	1614	90 (2)	51 (2)	58 (2)	71 (2)	88 (2)	107 (3)	128 (4)	142 (5)
Female	9-13	355	71 (3)	46 (6)	51 (5)	60 (4)	71 (4)	84 (5)	98 (6)	107 (8)
	14-18	410	69 (3)	39 (5)	45 (5)	55 (4)	67 (4)	82 (5)	97 (7)	107 (8)
	19-30	384	71 (4)	45 (5)	51 (4)	61 (4)	73 (4)	86 (5)	100 (7)	110 (9)
	31-50	585	68 (3)	37 (4)	42 (4)	53 (4)	67 (4)	83 (4)	98 (5)	108 (6)
	51-70	711	61 (3)	35 (3)	40 (3)	48 (3)	59 (3)	71 (4)	84 (5)	92 (5)
	>70	342	55 (4)	31 (4)	35 (4)	43 (4)	54 (5)	67 (6)	81 (7)	90 (8)
	19+	2022	65 (2)	36 (2)	41 (2)	52 (2)	65 (2)	79 (3)	94 (3)	105 (4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.12 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	622	48	(1)	30	(2)	33	(2)	40	(2)	47	(2)	55	(2)	63	(3)	68	(3)
	4-8	919	64	(2)	42	(4)	46	(3)	53	(2)	62	(2)	73	(3)	84	(5)	91	(6)
Male																		
	9-13	579	86	(3)	58	(5)	63	(5)	74	(4)	87	(4)	102	(5)	117	(7)	127	(9)
	14-18	634	104	(4)	66	(7)	74	(6)	87	(5)	104	(4)	123	(5)	143	(8)	157	(11)
	19-30	578	95	(3)	55	(8)	62	(7)	75	(6)	93	(5)	113	(6)	133	(10)	145	(13)
	31-50	693	90	(4)	47	(6)	54	(6)	68	(5)	89	(4)	110	(6)	132	(9)	146	(11)
	51-70	596	78	(3)	38	(6)	46	(6)	60	(5)	77	(4)	97	(5)	117	(7)	131	(9)
	>70	296	68	(3)	33	(4)	39	(4)	50	(4)	64	(4)	81	(4)	97	(6)	108	(7)
	19+	2163	86	(2)	43	(3)	51	(3)	65	(3)	84	(2)	107	(3)	130	(4)	145	(5)
Female																		
	9-13	533	67	(2)	44	(5)	49	(4)	57	(4)	67	(3)	78	(4)	89	(5)	97	(6)
	14-18	638	69	(2)	40	(3)	45	(3)	54	(3)	66	(3)	81	(4)	96	(5)	106	(6)
	19-30	499	69	(3)	39	(6)	44	(6)	54	(5)	67	(4)	82	(5)	97	(7)	107	(9)
	31-50	716	67	(2)	48	(7)	52	(6)	59	(5)	67	(4)	77	(4)	86	(6)	92	(8)
	51-70	745	63	(2)	37	(3)	42	(3)	51	(3)	62	(2)	75	(3)	88	(5)	97	(7)
	>70	510	49	(2)	26	(2)	30	(2)	38	(2)	48	(3)	60	(3)	73	(4)	82	(5)
	19+	2470	64	(1)	37	(2)	41	(2)	51	(2)	63	(2)	77	(2)	92	(3)	102	(4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 36.13 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	2117	51	(1)	31	(1)	35	(1)	42	(1)	51	(1)	61	(1)	71	(2)	77	(2)
	4-8	3235	66	(1)	43	(1)	47	(1)	55	(1)	65	(1)	76	(1)	88	(2)	95	(3)
Male																		
	9-13	2080	88	(2)	56	(2)	62	(2)	73	(2)	87	(2)	104	(3)	121	(3)	133	(4)
	14-18	2288	104	(2)	63	(3)	72	(3)	87	(3)	105	(2)	127	(3)	149	(4)	165	(6)
	19-30	1804	97	(2)	60	(3)	67	(3)	79	(3)	95	(3)	114	(4)	133	(5)	146	(7)
	31-50	2596	92	(2)	48	(3)	55	(2)	70	(2)	90	(2)	112	(3)	136	(5)	152	(6)
	51-70	2550	82	(2)	42	(2)	48	(2)	61	(2)	78	(2)	99	(2)	121	(3)	136	(4)
	>70	1520	67	(2)	34	(2)	40	(2)	50	(2)	64	(2)	81	(3)	98	(3)	109	(4)
	19+	8470	88	(1)	45	(1)	52	(1)	66	(1)	85	(1)	107	(2)	131	(2)	147	(3)
Female																		
	9-13	1980	72	(1)	44	(2)	49	(2)	58	(2)	70	(2)	84	(2)	98	(3)	107	(3)
	14-18	2256	73	(1)	43	(2)	49	(2)	59	(2)	72	(2)	88	(2)	104	(3)	114	(4)
	19-30	1854	67	(2)	37	(2)	42	(2)	52	(2)	64	(2)	78	(2)	92	(3)	102	(4)
	31-50	2686	69	(2)	40	(2)	46	(2)	56	(2)	69	(2)	84	(2)	99	(3)	110	(4)
	51-70	3200	62	(1)	35	(2)	40	(2)	49	(1)	61	(1)	75	(2)	89	(2)	98	(3)
	>70	2610	53	(1)	31	(1)	35	(1)	42	(1)	52	(1)	63	(2)	76	(2)	84	(3)
	19+	10350	65	(1)	36	(1)	41	(1)	51	(1)	64	(1)	78	(1)	94	(2)	104	(2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

37. Total monounsaturated fats (g/d): Usual intakes from food

Table 37.1 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	20.5 (1.7)	12.6 (2.5) ^E	14.1 (2.4) ^E	16.9 (2.2)	20.2 (2.1)	23.8 (2.4)	27.0 (2.9)	29.0 (3.4)
	4-8	127	31.1 (2.3)	21.5 (3.6)	23.5 (3.3)	27.0 (3.1)	31.3 (3.1)	36.4 (3.8)	41.8 (5.1)	45.3 (6.1)
Male	9-13	111	39.9 (2.6)	26.1 (4.3) ^E	28.7 (4.0)	33.6 (3.5)	39.9 (3.4)	47.2 (4.5)	54.9 (6.4)	60.0 (7.9)
	14-18	107	39.3 (2.6)	24.6 (5.3) ^E	27.4 (4.8) ^E	32.5 (3.9)	39.1 (3.3)	47.3 (4.0)	56.3 (6.3)	62.3 (8.4)
	19-30	77	39.9 (3.2)	23.6 (3.6)	26.9 (3.6)	32.7 (3.6)	39.3 (3.7)	45.9 (4.2)	52.2 (5.1)	56.3 (5.8)
	31-50	145	37.8 (3.0)	27.1 (3.8)	29.4 (3.4)	33.2 (3.0)	37.3 (3.0)	41.6 (3.5)	45.9 (4.4)	48.8 (5.2)
	51-70	182	28.9 (2.0)	15.7 (3.7) ^E	17.6 (3.4) ^E	21.2 (2.9)	26.1 (2.4)	31.8 (2.6)	37.8 (4.1)	41.7 (5.4)
	>70	63	29.3 (2.0)	19.5 (4.5) ^E	22.2 (4.0) ^E	27.0 (3.3)	32.4 (2.8)	37.6 (3.1)	42.1 (3.9)	44.8 (4.6)
	19+	467	34.8 (1.6)	20.1 (2.1)	23.0 (1.9)	28.3 (1.8)	34.6 (1.9)	41.3 (2.2)	47.8 (2.9)	52.0 (3.5)
Female	9-13	96	29.3 (1.4)	19.7 (3.0)	21.6 (2.6)	24.9 (2.1)	29.2 (1.8)	34.2 (2.4)	39.6 (3.8)	43.4 (5.0)
	14-18	105	31.3 (2.6)	19.1 (3.9) ^E	20.9 (3.7) ^E	24.3 (3.5)	28.9 (3.5)	34.8 (4.1)	41.3 (5.6)	45.8 (7.0)
	19-30	91	28.0 (3.0)	20.9 (3.8) ^E	22.2 (3.5)	24.3 (3.1)	26.9 (3.1)	29.6 (3.5)	32.3 (4.4)	34.1 (5.1)
	31-50	167	27.3 (1.9)	13.0 (2.7) ^E	15.5 (2.6) ^E	20.1 (2.5)	26.0 (2.6)	33.0 (3.0)	40.5 (3.9)	45.6 (4.7)
	51-70	198	21.5 (1.6)	8.9 (2.1) ^E	11.0 (2.0) ^E	15.3 (1.8)	21.1 (1.8)	28.3 (2.6)	35.9 (3.9)	40.8 (4.8)
	>70	74	22.5 (2.5)	11.9 (1.4)	13.7 (1.7)	17.2 (2.2)	22.1 (2.9)	27.8 (3.7)	33.4 (4.4)	36.9 (4.8)
	19+	530	25.2 (1.0)	12.4 (1.1)	14.5 (1.1)	18.5 (1.1)	24.0 (1.2)	30.5 (1.5)	37.1 (2.0)	41.4 (2.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.2 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	17.2 (1.3)	11.7 (1.9)	12.7 (1.8)	14.4 (1.6)	16.5 (1.6)	18.6 (1.9)	20.7 (2.4)	22.0 (2.7)
	4-8	110	25.0 (1.5)	18.1 (1.7)	19.5 (1.8)	22.1 (2.0)	25.3 (2.1)	28.5 (2.2)	31.2 (2.2)	32.7 (2.2)
Male	9-13	95	31.2 (2.4)	24.1 (3.7)	25.8 (3.5)	28.9 (3.2)	32.8 (3.0)	37.3 (3.3)	41.9 (4.2)	44.9 (5.1)
	14-18	87	42.7 (3.2)	28.7 (5.2) ^E	31.5 (4.8)	36.6 (4.2)	43.0 (4.0)	50.6 (5.3)	58.4 (8.0)	63.6 (10.3)
	19-30	70	50.9 (5.2)	41.2 (6.9) ^E	44.1 (6.5)	49.6 (6.0)	56.4 (6.1)	63.8 (7.6)	70.9 (10.0)	75.3 (11.8)
	31-50	109	37.7 (2.0)	29.4 (4.6)	31.5 (4.0)	35.2 (3.0)	39.7 (2.3)	44.3 (2.9)	48.7 (4.3)	51.4 (5.5)
	51-70	128	30.6 (1.4)	20.0 (4.0) ^E	22.3 (3.6)	26.4 (2.9)	31.4 (2.3)	36.8 (2.3)	42.0 (3.2)	45.2 (4.0)
	>70	65	24.4 (1.5)	15.3 (3.2) ^E	16.7 (2.9) ^E	19.5 (2.4)	23.0 (2.1)	27.1 (2.6)	31.4 (3.8)	34.3 (4.8)
	19+	372	37.0 (1.5)	23.0 (2.5)	25.9 (2.3)	31.3 (1.9)	38.1 (1.9)	46.3 (2.5)	54.6 (3.5)	60.0 (4.3)
Female	9-13	75	28.9 (3.0)	18.4 (3.4) ^E	20.4 (3.4) ^E	24.4 (3.4)	29.6 (3.9)	35.9 (4.8)	42.7 (6.4)	47.3 (7.6)
	14-18	81	29.6 (2.4)	17.0 (3.1) ^E	19.7 (3.0)	24.7 (2.8)	30.8 (3.1)	37.6 (3.8)	44.2 (4.9)	48.4 (5.7)
	19-30	101	26.7 (2.0)	17.2 (2.8)	19.2 (2.6)	23.1 (2.3)	28.2 (2.7)	34.1 (3.9)	40.0 (5.6)	43.9 (6.9)
	31-50	116	25.4 (1.6)	15.3 (3.2) ^E	17.3 (2.9)	20.8 (2.4)	25.1 (2.1)	29.8 (2.4)	34.4 (3.4)	37.3 (4.1)
	51-70	146	23.6 (1.1)	15.4 (2.7) ^E	17.0 (2.5)	19.8 (2.1)	23.3 (1.5)	27.1 (2.5)	30.8 (3.1)	33.2 (3.6)
	>70	94	21.7 (1.9)	14.7 (2.3)	16.0 (2.2)	18.4 (1.9)	21.5 (1.9)	25.2 (2.4)	29.0 (3.5)	31.6 (4.4)
	19+	457	24.7 (0.8)	15.2 (1.4)	17.0 (1.3)	20.5 (1.2)	24.9 (1.2)	30.0 (1.4)	35.0 (2.0)	38.2 (2.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.3 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	17.7 (1.0)	10.9 (1.8) ^E	12.2 (1.6)	14.5 (1.4)	17.2 (1.3)	20.4 (1.6)	23.6 (2.3)	25.8 (2.8)
	4-8	177	24.5 (1.3)	18.5 (3.5) ^E	19.8 (3.0)	22.1 (2.2)	24.8 (1.5)	27.6 (1.9)	30.3 (3.2)	31.9 (4.2)
Male	9-13	111	33.6 (2.1)	23.9 (3.2)	25.7 (2.9)	29.0 (2.5)	33.0 (2.3)	37.4 (2.9)	41.8 (4.0)	44.6 (4.9)
	14-18	113	42.4 (3.1)	28.6 (4.9) ^E	30.8 (4.5)	34.8 (4.5)	40.3 (5.2)	48.2 (6.2)	57.2 (7.5)	63.4 (8.6)
	19-30	91	40.2 (3.2)	22.6 (5.0) ^E	25.4 (4.7) ^E	30.9 (4.2)	38.3 (4.0)	47.4 (5.2)	57.4 (8.1)	64.3 (10.7) ^E
	31-50	101	41.5 (2.8)	20.4 (4.1) ^E	24.2 (3.9)	31.3 (3.7)	40.2 (3.9)	50.1 (4.6)	59.8 (5.6)	65.9 (6.5)
	51-70	134	33.0 (3.2)	20.6 (2.9)	22.7 (3.1)	26.8 (3.4)	31.7 (3.9)	37.2 (4.3)	43.0 (4.7)	46.7 (5.0)
	>70	56	24.4 (2.1)	15.2 (2.7) ^E	17.2 (2.7)	21.0 (2.6)	25.2 (2.8)	28.8 (3.3)	31.8 (3.6)	33.6 (3.9)
	19+	382	37.0 (1.6)	18.5 (2.7)	21.6 (2.4)	27.1 (2.1)	34.8 (2.0)	44.7 (2.9)	54.7 (4.5)	60.7 (5.4)
Female	9-13	105	28.1 (2.0)	22.1 (4.1) ^E	23.4 (3.7)	25.7 (3.0)	28.6 (2.5)	31.7 (2.9)	34.8 (4.4)	36.8 (5.8)
	14-18	120	25.2 (2.9)	14.4 (4.3) ^E	16.4 (3.9) ^E	20.1 (3.4) ^E	24.7 (3.0)	29.7 (3.4)	34.8 (4.6)	38.1 (5.7)
	19-30	91	32.8 (3.4)	15.4 (3.3) ^E	18.5 (3.2) ^E	24.6 (3.1)	32.3 (3.3)	40.8 (4.5)	52.7 (7.3)	62.9 (10.1)
	31-50	159	31.6 (2.7)	17.8 (3.6) ^E	20.6 (3.4)	25.7 (3.2)	31.7 (3.2)	37.9 (3.6)	43.7 (4.3)	47.3 (4.9)
	51-70	174	27.5 (2.7)	14.6 (3.1) ^E	16.7 (2.9) ^E	20.6 (2.5)	25.6 (2.5)	31.5 (3.3)	37.7 (4.9)	41.7 (6.2)
	>70	80	21.5 (2.3)	15.2 (3.2) ^E	16.8 (3.2) ^E	19.7 (3.2)	23.0 (3.3)	26.4 (3.6)	29.5 (3.9)	31.5 (4.3)
	19+	504	29.3 (1.6)	14.3 (1.5)	17.1 (1.6)	22.2 (1.6)	29.1 (1.9)	37.3 (2.5)	46.0 (3.3)	51.9 (3.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.4 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	19.7 (1.3)	12.4 (2.0)	13.8 (1.8)	16.3 (1.6)	19.4 (1.6)	22.8 (1.9)	26.1 (2.4)	28.2 (2.9)
	4-8	140	25.5 (1.1)	19.1 (1.2)	20.5 (1.2)	23.1 (1.3)	26.1 (1.3)	29.1 (1.4)	31.8 (1.5)	33.6 (1.6)
Male	9-13	92	41.6 (8.4) ^E	20.4 (6.4) ^E	24.1 (6.0) ^E	28.9 (6.2) ^E	35.9 (6.7) ^E	46.4 (8.9) ^E	57.7 (12.4) ^E	63.9 (14.5) ^E
	14-18	107	44.3 (3.3)	23.6 (4.3) ^E	27.4 (4.3)	34.7 (4.2)	44.3 (4.3)	56.2 (5.2)	69.4 (7.4)	78.8 (9.5)
	19-30	73	44.1 (3.6)	24.0 (5.2) ^E	28.3 (4.7)	35.6 (4.3)	44.7 (5.1)	56.2 (7.3)	70.4 (11.0)	81.3 (14.3) ^E
	31-50	134	36.0 (2.3)	24.2 (4.1) ^E	26.2 (3.7)	29.8 (2.9)	34.3 (2.5)	39.3 (3.3)	44.3 (4.9)	47.5 (6.2)
	51-70	131	34.8 (2.4)	23.1 (2.1)	25.4 (2.2)	29.2 (2.3)	33.4 (2.5)	37.9 (2.8)	42.5 (3.4)	45.6 (3.9)
	>70	55	30.8 (3.0)	19.1 (4.0) ^E	21.0 (3.8) ^E	24.8 (3.4)	29.7 (3.4)	35.5 (4.1)	41.7 (5.6)	45.9 (6.8)
	19+	393	36.9 (1.5)	22.2 (2.3)	24.9 (2.2)	29.9 (2.0)	36.3 (1.8)	43.7 (2.2)	51.6 (3.2)	57.2 (4.1)
Female	9-13	79	27.4 (3.3)	13.8 (4.3) ^E	16.5 (4.0) ^E	21.4 (3.6) ^E	27.2 (3.6)	34.0 (4.5)	41.4 (6.4)	46.7 (8.1) ^E
	14-18	104	25.6 (1.6)	17.0 (1.6)	18.8 (1.7)	22.0 (1.9)	25.7 (2.2)	29.9 (2.7)	34.2 (3.3)	37.0 (3.7)
	19-30	101	25.6 (2.2)	15.7 (4.5) ^E	18.0 (4.2) ^E	22.3 (3.6)	27.3 (3.0)	32.7 (2.7)	37.9 (3.4)	41.4 (4.3)
	31-50	143	22.9 (1.3)	13.5 (1.2)	15.2 (1.2)	18.3 (1.4)	22.1 (1.7)	26.7 (2.0)	31.8 (2.2)	35.2 (2.3)
	51-70	193	24.0 (1.3)	17.3 (2.9)	18.5 (2.5)	20.6 (2.0)	23.1 (1.7)	25.8 (2.0)	28.4 (2.8)	30.0 (3.6)
	>70	94	19.0 (1.2)	11.1 (1.5)	12.2 (1.5)	14.3 (1.4)	17.0 (1.6)	20.2 (2.1)	23.8 (3.1)	26.2 (3.9)
	19+	531	23.2 (0.8)	14.0 (1.5)	15.8 (1.4)	19.1 (1.2)	23.1 (1.1)	27.5 (1.3)	31.7 (1.7)	34.5 (2.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.5 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	311	20.1 (0.9)	11.2 (1.3)	13.0 (1.3)	16.3 (1.2)	20.3 (1.3)	24.8 (1.6)	29.4 (2.1)	32.5 (2.5)
	4-8	485	28.1 (1.0)	16.8 (1.8)	19.2 (1.6)	23.2 (1.3)	27.5 (1.3)	33.6 (1.6)	39.1 (2.0)	42.1 (2.3)
Male	9-13	277	39.0 (2.3)	24.5 (3.1)	27.4 (3.0)	33.0 (2.9)	40.2 (3.1)	48.5 (3.9)	57.3 (5.2)	63.1 (6.2)
	14-18	339	46.1 (2.1)	29.5 (4.0)	32.7 (3.8)	39.0 (3.3)	47.5 (3.0)	57.4 (3.7)	67.7 (5.6)	74.9 (7.4)
	19-30	237	40.9 (1.7)	27.5 (2.3)	30.2 (2.4)	35.0 (2.7)	40.8 (3.1)	47.3 (3.5)	54.0 (3.9)	58.4 (4.2)
	31-50	423	39.7 (2.5)	20.2 (3.4) ^E	23.6 (3.2)	30.2 (3.0)	38.8 (3.1)	48.9 (4.1)	59.5 (5.8)	66.5 (7.2)
	51-70	387	35.3 (1.7)	17.4 (2.3)	20.2 (2.2)	25.6 (2.1)	33.0 (2.0)	42.3 (2.4)	52.4 (3.4)	59.3 (4.4)
	>70	132	28.9 (2.2)	15.1 (3.8) ^E	17.8 (3.6) ^E	22.9 (3.3)	29.1 (3.2)	35.9 (3.6)	42.8 (4.5)	47.4 (5.3)
	19+	1179	37.8 (1.2)	19.8 (1.4)	23.0 (1.4)	29.1 (1.4)	37.0 (1.5)	46.3 (1.9)	56.0 (2.5)	62.7 (3.1)
Female	9-13	281	30.5 (1.6)	18.8 (2.1)	20.9 (2.0)	24.7 (1.8)	29.4 (1.8)	34.7 (2.2)	40.3 (3.0)	44.0 (3.6)
	14-18	321	31.7 (1.7)	21.3 (3.0)	23.6 (2.8)	27.8 (2.5)	33.0 (2.6)	38.6 (3.3)	44.0 (4.4)	47.5 (5.3)
	19-30	249	30.0 (1.9)	22.4 (3.4)	23.8 (3.1)	26.3 (2.5)	29.3 (2.3)	32.6 (2.8)	35.7 (3.9)	37.7 (4.7)
	31-50	364	29.8 (1.4)	18.4 (2.6)	20.5 (2.4)	24.4 (2.1)	29.8 (1.9)	36.3 (2.3)	43.1 (3.5)	47.7 (4.6)
	51-70	467	27.0 (1.2)	14.8 (2.0)	17.1 (1.8)	21.4 (1.5)	26.9 (1.3)	33.0 (1.7)	39.1 (2.5)	43.0 (3.1)
	>70	215	20.5 (1.0)	14.3 (2.1)	15.5 (2.0)	17.7 (1.8)	20.4 (1.7)	23.6 (1.9)	26.8 (2.7)	28.9 (3.4)
	19+	1295	27.8 (0.7)	17.1 (1.2)	19.2 (1.1)	23.0 (1.0)	27.8 (0.9)	33.3 (1.1)	39.1 (1.6)	42.9 (2.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.6 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	17.1 (0.6)	8.5 (1.0)	10.1 (0.9)	13.0 (0.7)	16.6 (0.6)	21.0 (0.8)	25.6 (1.3)	28.6 (1.7)
	4-8	956	23.4 (0.6)	15.7 (1.7)	17.2 (1.5)	19.9 (1.1)	23.3 (0.7)	27.2 (1.0)	31.0 (1.8)	33.4 (2.4)
Male	9-13	589	31.4 (0.8)	21.6 (2.5)	23.5 (2.2)	27.0 (1.6)	31.3 (1.0)	36.2 (1.4)	41.1 (2.6)	44.4 (3.5)
	14-18	639	38.5 (1.2)	22.8 (2.7)	25.9 (2.4)	31.6 (1.9)	38.6 (1.5)	46.6 (2.0)	54.9 (3.2)	60.4 (4.2)
	19-30	481	38.6 (2.0)	21.0 (4.0) ^E	23.7 (3.6)	30.0 (2.7)	37.5 (2.2)	46.1 (3.2)	56.4 (5.8)	63.5 (8.0)
	31-50	709	34.5 (1.4)	20.6 (3.9) ^E	23.1 (3.4)	27.8 (2.5)	34.1 (1.6)	41.5 (2.5)	49.1 (4.7)	54.3 (6.3)
	51-70	758	31.9 (1.6)	16.3 (2.9) ^E	18.7 (2.6)	23.3 (2.2)	29.6 (1.7)	37.7 (1.8)	47.1 (3.3)	53.8 (4.8)
	>70	734	24.3 (0.7)	10.8 (1.0)	13.0 (1.0)	17.3 (0.9)	22.9 (0.8)	29.7 (1.1)	37.1 (1.6)	42.3 (2.2)
	19+	2682	33.8 (0.9)	16.9 (1.3)	19.5 (1.2)	24.8 (1.1)	32.3 (1.0)	41.5 (1.3)	51.6 (2.1)	58.6 (2.9)
Female	9-13	585	26.8 (0.9)	16.5 (2.3)	18.4 (2.1)	21.8 (1.6)	26.2 (1.1)	31.4 (1.5)	36.9 (2.6)	40.6 (3.5)
	14-18	645	28.6 (0.9)	18.1 (2.4)	20.1 (2.1)	23.8 (1.5)	28.6 (1.1)	34.1 (1.8)	39.7 (3.1)	43.2 (4.0)
	19-30	514	23.8 (1.0)	10.7 (1.5)	12.9 (1.4)	17.0 (1.2)	22.4 (1.2)	28.5 (1.4)	34.9 (2.1)	39.4 (2.8)
	31-50	758	26.4 (1.1)	14.3 (1.9)	16.4 (1.7)	20.4 (1.4)	25.5 (1.1)	31.5 (1.6)	37.8 (2.6)	42.1 (3.5)
	51-70	955	23.8 (0.8)	13.0 (1.9)	14.9 (1.7)	18.6 (1.4)	23.4 (1.0)	28.9 (1.1)	34.7 (1.8)	38.6 (2.6)
	>70	1345	21.1 (0.5)	12.4 (1.7)	13.9 (1.5)	16.8 (1.0)	20.5 (0.6)	24.8 (1.1)	29.3 (2.0)	32.3 (2.7)
	19+	3572	24.6 (0.5)	12.8 (0.8)	14.8 (0.8)	18.7 (0.7)	23.7 (0.6)	29.7 (0.7)	35.8 (1.1)	39.9 (1.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.7 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	16.7 (0.8)	8.1 (1.0)	9.7 (0.9)	12.5 (0.9)	16.0 (0.9)	20.0 (1.1)	24.2 (1.5)	27.1 (1.9)
	4-8	425	23.7 (0.8)	14.8 (3.0) ^E	16.5 (2.6)	19.5 (1.9)	23.2 (0.9)	27.5 (2.4)	31.8 (4.1)	34.7 (5.2)
Male	9-13	274	31.8 (1.4)	19.5 (3.4) ^E	22.0 (2.9)	26.4 (2.1)	31.8 (1.6)	38.0 (2.3)	44.6 (3.9)	49.2 (5.2)
	14-18	297	44.9 (2.2)	21.0 (3.6) ^E	25.3 (3.4)	33.4 (3.0)	43.8 (2.9)	56.4 (3.5)	70.5 (5.5)	80.5 (7.6)
	19-30	249	42.8 (2.7)	24.6 (2.0)	27.8 (2.2)	34.3 (2.6)	42.6 (3.1)	51.9 (3.7)	61.4 (4.6)	68.0 (5.5)
	31-50	309	33.9 (2.0)	16.1 (3.8) ^E	18.7 (3.6) ^E	23.9 (3.3)	32.1 (2.6)	43.0 (3.5)	54.9 (6.3)	64.0 (9.3)
	51-70	277	31.1 (1.6)	19.1 (3.9) ^E	21.4 (3.6) ^E	25.7 (3.0)	31.0 (2.3)	36.8 (2.6)	42.5 (4.2)	46.2 (5.6)
	>70	136	23.9 (1.7)	11.0 (2.0) ^E	12.9 (1.9)	16.6 (1.7)	21.6 (1.6)	27.6 (2.1)	34.2 (3.3)	38.6 (4.5)
	19+	971	34.2 (1.2)	18.3 (2.8)	20.8 (2.7)	25.8 (2.2)	33.2 (1.6)	42.2 (2.1)	51.5 (3.8)	57.8 (5.2)
Female	9-13	265	26.4 (1.2)	15.3 (2.2)	17.2 (1.9)	20.8 (1.5)	25.1 (1.3)	29.7 (1.6)	34.2 (2.3)	37.0 (2.8)
	14-18	290	28.9 (2.1)	15.8 (1.1)	18.2 (1.2)	22.7 (1.5)	28.5 (1.9)	35.4 (2.4)	42.7 (3.2)	47.6 (3.8)
	19-30	197	27.6 (2.4)	15.4 (1.5)	17.5 (1.7)	21.4 (2.0)	26.5 (2.4)	32.4 (2.8)	38.5 (3.2)	42.6 (3.5)
	31-50	312	27.2 (1.5)	19.2 (3.0)	20.8 (2.7)	23.7 (2.2)	27.4 (1.8)	31.6 (2.3)	35.9 (3.7)	38.7 (5.0)
	51-70	312	22.8 (0.9)	13.9 (2.1)	15.5 (1.9)	18.5 (1.4)	22.3 (1.1)	26.7 (1.7)	31.2 (2.9)	34.3 (3.8)
	>70	239	18.3 (1.0)	11.6 (2.1) ^E	12.7 (1.8)	14.8 (1.5)	17.4 (1.2)	20.7 (1.5)	24.1 (2.5)	26.4 (3.2)
	19+	1060	24.7 (0.8)	15.0 (1.8)	16.7 (1.6)	19.9 (1.2)	24.2 (0.9)	29.4 (1.4)	34.9 (2.5)	38.7 (3.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.8 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	19.9 (1.2)	11.0 (1.8)	12.7 (1.7)	15.9 (1.6)	20.1 (1.7)	24.9 (2.1)	29.7 (2.7)	32.9 (3.2)
	4-8	213	27.8 (1.9)	20.4 (2.7)	21.8 (2.4)	24.3 (2.0)	27.4 (1.8)	30.9 (2.3)	34.3 (3.5)	36.5 (4.4)
Male	9-13	122	34.1 (2.5)	23.1 (3.1)	25.1 (3.0)	29.1 (2.8)	34.4 (3.0)	40.9 (4.0)	47.6 (5.5)	52.0 (6.7)
	14-18	150	40.5 (2.6)	26.8 (2.3)	29.8 (2.4)	34.8 (2.8)	40.4 (3.4)	46.6 (4.4)	53.7 (5.2)	58.5 (5.7)
	19-30	106	45.4 (3.7)	19.1 (6.2) ^E	23.6 (5.8) ^E	32.4 (5.0)	44.1 (4.7)	57.8 (6.1)	71.8 (9.5)	81.0 (12.7)
	31-50	155	40.1 (3.1)	21.6 (5.0) ^E	25.1 (4.5) ^E	31.4 (3.8)	39.4 (3.7)	48.7 (5.0)	58.5 (7.4)	65.2 (9.3)
	51-70	122	36.0 (2.4)	21.8 (4.3) ^E	24.9 (3.8)	30.6 (3.1)	37.8 (3.2)	46.0 (4.8)	54.2 (7.3)	59.6 (9.3)
	>70	88	30.2 (1.8)	16.2 (3.2) ^E	18.6 (2.9)	23.0 (2.4)	28.9 (2.2)	35.6 (3.0)	42.6 (4.6)	47.2 (5.8)
	19+	471	39.0 (1.7)	19.1 (2.0)	22.7 (1.9)	29.4 (1.9)	38.3 (2.1)	49.1 (2.7)	60.7 (3.7)	68.6 (4.6)
Female	9-13	103	27.9 (1.9)	20.2 (3.7) ^E	21.9 (3.5)	24.8 (3.1)	28.4 (2.8)	32.3 (3.0)	36.1 (3.7)	38.5 (4.3)
	14-18	142	28.8 (1.5)	18.7 (2.2)	20.9 (2.0)	24.6 (1.8)	29.1 (2.1)	34.0 (2.8)	39.1 (3.9)	42.4 (4.7)
	19-30	111	27.9 (1.8)	17.8 (3.8) ^E	19.7 (3.5) ^E	23.2 (2.9)	27.7 (2.6)	32.7 (3.2)	37.9 (4.7)	41.2 (5.9)
	31-50	146	25.1 (1.3)	13.2 (2.7) ^E	15.1 (2.4)	18.5 (2.1)	22.8 (1.8)	28.1 (1.9)	33.4 (2.7)	36.9 (3.4)
	51-70	184	26.8 (1.3)	19.9 (3.2)	21.4 (2.8)	24.1 (2.1)	27.3 (1.7)	30.7 (2.3)	34.1 (3.6)	36.2 (4.7)
	>70	143	21.9 (1.7)	12.7 (3.1) ^E	14.5 (2.9) ^E	17.9 (2.5)	22.6 (2.3)	28.4 (3.0)	34.7 (4.6)	39.0 (6.0)
	19+	584	25.6 (0.8)	14.6 (1.3)	16.6 (1.2)	20.3 (1.1)	25.0 (1.1)	30.3 (1.2)	35.7 (1.6)	39.2 (2.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.9 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	169	14.9 (0.9)	7.0 (1.9) ^E	8.5 (1.7) ^E	11.3 (1.3)	14.8 (1.0)	18.8 (1.3)	23.0 (2.1)	25.8 (2.8)
	4-8	281	24.1 (1.0)	16.5 (2.7)	17.8 (2.4)	20.3 (1.8)	23.4 (1.3)	27.0 (1.8)	30.4 (3.1)	32.6 (4.1)
Male	9-13	183	34.8 (2.2)	21.4 (1.7)	23.7 (1.8)	28.2 (2.0)	34.2 (2.4)	41.4 (3.1)	49.1 (3.9)	54.3 (4.6)
	14-18	187	39.8 (2.2)	26.2 (4.9) ^E	29.0 (4.4)	34.0 (3.4)	40.1 (2.6)	47.0 (3.2)	54.1 (5.2)	58.8 (6.9)
	19-30	223	36.8 (2.1)	20.4 (4.0) ^E	23.3 (3.6)	28.9 (2.8)	36.0 (2.4)	44.3 (3.3)	52.8 (5.2)	58.4 (6.7)
	31-50	229	36.8 (2.6)	17.9 (2.1)	21.1 (2.2)	27.4 (2.4)	36.5 (2.9)	47.2 (3.9)	57.6 (5.4)	64.6 (6.7)
	51-70	197	30.5 (2.2)	14.6 (2.5) ^E	17.5 (2.5)	23.0 (2.5)	30.4 (2.7)	38.7 (3.1)	46.9 (3.9)	52.5 (5.0)
	>70	72	26.7 (2.5)	15.9 (4.0) ^E	17.9 (3.8) ^E	21.6 (3.3)	26.1 (3.0)	30.8 (3.3)	35.1 (4.1)	37.5 (4.7)
	19+	721	34.6 (1.4)	16.4 (1.2)	19.7 (1.2)	25.9 (1.3)	34.1 (1.5)	43.7 (2.0)	53.8 (2.6)	60.5 (3.2)
Female	9-13	165	24.6 (1.5)	15.8 (3.4) ^E	17.5 (3.1) ^E	20.7 (2.6)	24.7 (2.1)	29.1 (3.4)	33.7 (4.6)	36.8 (5.7)
	14-18	206	25.8 (1.4)	15.3 (3.1) ^E	17.1 (2.7)	20.5 (2.0)	24.7 (1.6)	29.6 (2.3)	34.8 (3.9)	38.4 (5.1)
	19-30	191	26.7 (1.8)	12.9 (2.9) ^E	15.2 (2.8) ^E	19.6 (2.6)	25.9 (2.7)	33.3 (3.3)	40.8 (4.4)	46.1 (5.6)
	31-50	258	26.7 (1.4)	19.6 (1.6)	21.2 (1.8)	23.9 (2.0)	27.3 (2.3)	31.0 (2.7)	34.7 (3.0)	37.0 (3.3)
	51-70	249	25.1 (1.4)	10.7 (2.0) ^E	13.1 (1.9)	17.9 (1.6)	24.1 (1.7)	31.9 (2.3)	40.5 (3.4)	46.7 (4.3)
	>70	128	18.4 (1.2)	8.4 (1.4) ^E	10.0 (1.4)	13.2 (1.5)	17.4 (1.7)	22.4 (1.8)	27.6 (2.3)	31.1 (2.8)
	19+	826	25.4 (0.8)	13.3 (1.3)	15.5 (1.3)	19.6 (1.2)	25.1 (1.3)	31.7 (1.4)	38.6 (1.8)	43.2 (2.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.10 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	192	18.4 (1.1)	9.7 (1.8) ^E	11.2 (1.6)	14.0 (1.4)	17.5 (1.4)	21.3 (1.7)	25.0 (2.3)	27.4 (2.8)
	4-8	321	24.0 (1.0)	16.8 (2.5)	18.2 (2.2)	20.8 (1.7)	24.0 (1.3)	27.5 (1.7)	31.0 (2.5)	33.3 (3.2)
Male	9-13	226	34.1 (1.7)	18.8 (2.6)	21.7 (2.5)	27.2 (2.3)	34.1 (2.2)	42.0 (2.5)	50.6 (3.5)	56.8 (4.7)
	14-18	262	42.9 (2.5)	31.4 (4.9)	33.8 (4.4)	38.2 (3.6)	43.4 (3.2)	49.1 (3.9)	54.7 (5.6)	58.2 (7.0)
	19-30	197	42.6 (2.2)	28.1 (3.4)	30.8 (3.1)	36.0 (2.6)	42.5 (2.6)	50.1 (3.8)	57.9 (5.9)	63.1 (7.5)
	31-50	282	41.4 (2.2)	22.7 (3.5)	25.8 (3.1)	31.1 (2.7)	38.5 (2.9)	49.2 (4.0)	61.3 (5.8)	69.1 (7.4)
	51-70	234	35.2 (2.0)	18.7 (3.1) ^E	21.4 (2.9)	26.5 (2.6)	33.6 (2.5)	42.1 (3.2)	51.4 (4.8)	57.7 (6.2)
	>70	119	27.1 (2.0)	12.3 (2.4) ^E	14.8 (2.4)	19.4 (2.4)	25.5 (2.4)	32.9 (2.7)	41.3 (3.5)	47.2 (4.3)
	19+	832	38.5 (1.1)	20.3 (1.4)	23.3 (1.3)	29.2 (1.2)	37.1 (1.4)	47.0 (1.9)	57.9 (2.7)	65.2 (3.4)
Female	9-13	226	26.9 (1.4)	17.7 (2.0)	19.6 (1.9)	23.0 (1.7)	27.3 (1.6)	32.3 (1.9)	37.4 (2.6)	40.8 (3.2)
	14-18	242	26.6 (1.4)	12.6 (2.1) ^E	15.0 (2.0)	19.5 (1.8)	25.5 (1.7)	32.5 (2.1)	39.7 (2.8)	44.4 (3.4)
	19-30	208	26.7 (1.5)	13.6 (2.2)	16.0 (2.1)	20.5 (2.0)	25.8 (2.0)	31.9 (2.3)	38.7 (3.2)	43.8 (4.1)
	31-50	263	29.8 (2.1)	16.0 (2.1)	18.4 (2.1)	23.1 (2.0)	29.5 (2.3)	37.3 (3.1)	45.6 (4.2)	51.1 (5.2)
	51-70	322	23.5 (1.6)	10.3 (1.7) ^E	12.4 (1.7)	16.3 (1.7)	21.8 (1.8)	29.4 (2.4)	38.1 (3.6)	44.0 (4.6)
	>70	198	21.1 (1.1)	12.2 (1.8)	13.8 (1.7)	16.7 (1.5)	20.3 (1.5)	24.6 (1.9)	28.9 (2.7)	31.7 (3.4)
	19+	991	26.3 (1.0)	12.5 (0.9)	14.8 (0.9)	19.3 (0.9)	25.6 (1.1)	33.2 (1.4)	41.4 (1.9)	47.0 (2.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.11 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	348	18.9 (0.7)	11.1 (1.0)	12.5 (0.9)	15.0 (0.9)	18.5 (1.0)	22.5 (1.1)	26.5 (1.3)	29.0 (1.5)
	4-8	554	26.3 (0.8)	18.5 (1.9)	20.3 (1.7)	23.3 (1.3)	26.9 (1.1)	30.8 (1.3)	34.4 (1.9)	36.7 (2.4)
Male	9-13	409	37.3 (2.9)	24.2 (2.6)	26.5 (2.5)	31.0 (2.4)	36.6 (2.8)	43.1 (3.6)	49.8 (4.8)	54.1 (5.7)
	14-18	414	42.3 (1.8)	25.7 (2.3)	28.8 (2.3)	34.4 (2.3)	42.0 (2.5)	51.8 (3.0)	62.0 (3.7)	68.3 (4.3)
	19-30	311	42.0 (1.9)	24.4 (2.8)	27.8 (2.7)	34.1 (2.4)	42.0 (2.5)	51.0 (3.3)	60.7 (4.7)	67.4 (6.0)
	31-50	489	38.7 (1.5)	22.9 (2.0)	25.8 (1.9)	31.0 (1.7)	37.5 (1.8)	44.7 (2.3)	52.0 (3.0)	56.7 (3.5)
	51-70	575	32.5 (1.6)	20.3 (2.8)	22.4 (2.5)	26.3 (2.0)	31.4 (1.7)	37.1 (2.4)	42.4 (3.7)	45.9 (4.7)
	>70	239	27.5 (1.4)	18.4 (2.2)	20.5 (2.1)	24.1 (1.9)	28.2 (1.9)	32.5 (2.1)	36.8 (2.8)	39.8 (3.3)
	19+	1614	36.5 (0.9)	20.0 (1.1)	22.9 (1.0)	28.4 (1.0)	35.4 (1.1)	43.6 (1.3)	52.2 (1.8)	58.0 (2.2)
Female	9-13	355	28.2 (1.4)	17.0 (2.5)	19.2 (2.2)	23.3 (1.9)	28.3 (1.7)	33.8 (2.0)	39.7 (2.8)	43.6 (3.6)
	14-18	410	27.0 (1.4)	14.8 (2.1)	17.0 (1.9)	21.0 (1.7)	26.0 (1.7)	31.9 (2.0)	38.2 (2.8)	42.4 (3.5)
	19-30	384	29.1 (1.7)	17.7 (1.4)	20.0 (1.5)	24.5 (1.6)	30.5 (1.9)	36.4 (2.3)	42.6 (2.9)	46.8 (3.4)
	31-50	585	27.6 (1.3)	14.8 (1.6)	17.1 (1.6)	21.4 (1.6)	27.0 (1.6)	33.4 (1.8)	39.5 (2.2)	43.3 (2.5)
	51-70	711	24.8 (1.2)	13.7 (1.4)	15.6 (1.4)	19.2 (1.3)	24.0 (1.4)	29.2 (1.6)	34.5 (2.1)	38.0 (2.5)
	>70	342	20.9 (1.2)	11.8 (1.2)	13.3 (1.3)	16.2 (1.4)	20.3 (1.6)	25.3 (2.0)	30.4 (2.5)	33.8 (2.9)
	19+	2022	26.2 (0.8)	14.1 (0.8)	16.3 (0.8)	20.5 (0.9)	26.0 (1.0)	32.3 (1.2)	38.8 (1.5)	43.2 (1.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.12 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	622	16.2 (0.6)	7.7 (1.0)	9.2 (0.9)	12.1 (0.8)	15.8 (0.7)	20.1 (0.8)	24.5 (1.2)	27.5 (1.5)
	4-8	919	24.7 (0.7)	15.3 (1.6)	16.9 (1.4)	20.0 (1.1)	23.9 (0.9)	28.5 (1.2)	33.1 (2.0)	36.2 (2.6)
Male	9-13	579	34.0 (1.4)	21.4 (2.4)	23.7 (2.2)	28.1 (1.8)	33.9 (1.6)	40.7 (2.1)	48.0 (3.2)	52.9 (4.2)
	14-18	634	41.0 (1.5)	26.0 (3.2)	29.2 (2.8)	34.9 (2.2)	41.5 (1.8)	48.4 (2.1)	55.9 (3.3)	61.2 (4.4)
	19-30	578	39.4 (1.6)	22.0 (3.5)	25.1 (3.1)	31.0 (2.4)	38.6 (2.0)	47.4 (2.8)	56.4 (4.5)	62.3 (5.8)
	31-50	693	36.8 (1.8)	18.2 (2.8)	21.3 (2.6)	27.4 (2.2)	35.9 (2.0)	46.1 (2.8)	56.5 (4.6)	63.5 (6.1)
	51-70	596	31.7 (1.4)	16.1 (3.2) ^E	19.0 (2.9)	24.5 (2.3)	31.7 (1.8)	39.9 (2.2)	48.4 (3.5)	54.0 (4.7)
	>70	296	26.9 (1.4)	12.5 (1.7)	14.8 (1.8)	19.2 (1.8)	25.0 (1.7)	32.0 (1.9)	39.4 (2.5)	44.3 (3.1)
	19+	2163	35.3 (0.9)	17.0 (1.3)	20.2 (1.2)	26.4 (1.1)	34.6 (1.1)	44.5 (1.4)	54.5 (2.0)	61.3 (2.6)
Female	9-13	533	25.6 (1.0)	15.9 (2.0)	17.8 (1.8)	21.2 (1.5)	25.4 (1.4)	30.2 (1.7)	35.1 (2.4)	38.4 (3.0)
	14-18	638	27.0 (1.0)	15.0 (1.3)	17.2 (1.2)	21.0 (1.1)	25.8 (1.1)	31.7 (1.5)	38.0 (2.2)	42.2 (2.8)
	19-30	499	27.1 (1.3)	15.1 (2.5) ^E	17.2 (2.3)	21.2 (2.0)	26.3 (1.8)	32.2 (2.1)	38.3 (3.1)	42.2 (3.9)
	31-50	716	26.5 (1.0)	19.2 (2.8)	20.7 (2.5)	23.4 (2.0)	26.8 (1.6)	30.5 (1.7)	34.2 (2.5)	36.6 (3.2)
	51-70	745	24.9 (0.9)	14.0 (1.4)	16.1 (1.3)	19.8 (1.1)	24.6 (1.1)	30.1 (1.4)	35.8 (2.0)	39.8 (2.6)
	>70	510	19.2 (0.7)	9.7 (1.0)	11.1 (1.0)	14.2 (1.0)	18.4 (1.1)	23.9 (1.2)	29.0 (1.5)	32.2 (1.9)
	19+	2470	25.3 (0.5)	13.9 (0.8)	16.0 (0.8)	19.8 (0.8)	24.8 (0.8)	30.9 (0.9)	37.3 (1.2)	41.4 (1.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.13 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	17.9 (0.4)	9.0 (0.5)	10.7 (0.5)	13.6 (0.4)	17.4 (0.5)	22.0 (0.6)	26.5 (0.8)	29.6 (0.9)
	4-8	3235	24.9 (0.4)	15.7 (0.6)	17.5 (0.6)	20.7 (0.5)	24.7 (0.5)	29.3 (0.6)	33.9 (0.9)	36.9 (1.1)
Male	9-13	2080	34.3 (0.7)	21.7 (1.0)	24.1 (0.9)	28.5 (0.9)	34.2 (0.9)	40.8 (1.1)	47.8 (1.5)	52.6 (2.0)
	14-18	2288	41.5 (0.8)	25.0 (1.3)	28.3 (1.3)	34.3 (1.1)	41.9 (1.1)	50.6 (1.3)	59.7 (1.9)	65.8 (2.4)
	19-30	1804	40.1 (1.0)	23.8 (1.5)	26.7 (1.4)	32.3 (1.2)	39.4 (1.3)	47.5 (1.7)	56.0 (2.5)	61.7 (3.2)
	31-50	2596	37.3 (0.9)	18.8 (1.1)	21.9 (1.1)	28.2 (1.0)	36.3 (1.1)	46.0 (1.4)	56.3 (2.1)	63.2 (2.6)
	51-70	2550	33.3 (0.8)	16.8 (1.1)	19.5 (1.0)	24.6 (0.9)	31.6 (0.9)	40.2 (1.1)	49.6 (1.7)	56.0 (2.2)
	>70	1520	26.4 (0.7)	12.8 (0.9)	15.2 (0.9)	19.7 (0.9)	25.4 (0.9)	32.2 (1.1)	39.1 (1.4)	43.8 (1.7)
	19+	8470	35.8 (0.5)	17.9 (0.5)	20.9 (0.5)	26.7 (0.5)	34.5 (0.6)	44.0 (0.8)	54.2 (1.1)	61.1 (1.4)
Female	9-13	1980	27.6 (0.6)	16.5 (0.8)	18.5 (0.7)	22.3 (0.7)	27.1 (0.7)	32.5 (0.8)	38.2 (1.1)	42.0 (1.4)
	14-18	2256	28.6 (0.6)	16.5 (0.7)	18.7 (0.7)	23.0 (0.7)	28.4 (0.8)	34.7 (1.0)	41.0 (1.3)	45.1 (1.6)
	19-30	1854	26.6 (0.7)	14.2 (0.9)	16.4 (0.8)	20.4 (0.8)	25.5 (0.8)	31.2 (1.0)	37.1 (1.4)	41.1 (1.7)
	31-50	2686	27.7 (0.7)	15.4 (0.9)	17.6 (0.8)	21.8 (0.8)	27.3 (0.8)	33.8 (0.9)	40.6 (1.3)	45.1 (1.7)
	51-70	3200	24.9 (0.5)	13.3 (0.7)	15.4 (0.7)	19.3 (0.6)	24.5 (0.6)	30.3 (0.7)	36.4 (1.0)	40.6 (1.2)
	>70	2610	20.6 (0.4)	11.7 (0.6)	13.2 (0.6)	16.2 (0.5)	20.1 (0.5)	24.7 (0.7)	29.7 (1.0)	33.0 (1.3)
	19+	10350	25.8 (0.3)	13.7 (0.4)	15.9 (0.4)	19.9 (0.4)	25.2 (0.4)	31.5 (0.5)	38.0 (0.7)	42.5 (0.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

38. Total polyunsaturated fats (g/d): Usual intakes from food

Table 38.1 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	6.7 (0.6)	3.9 (0.8) ^E	4.4 (0.7)	5.3 (0.7)	6.4 (0.8)	7.6 (1.0)	8.9 (1.2)	9.8 (1.5)
	4-8	127	12.1 (0.7)	7.8 (1.3) ^E	8.6 (1.2)	10.2 (1.1)	12.5 (1.2)	14.9 (1.5)	16.9 (1.7)	18.2 (1.9)
Male	9-13	111	16.2 (1.0)	9.7 (1.7) ^E	10.9 (1.6)	13.2 (1.4)	16.2 (1.5)	19.8 (1.8)	23.7 (2.5)	26.3 (3.1)
	14-18	107	15.7 (1.1)	7.8 (1.9) ^E	9.1 (1.8) ^E	11.8 (1.5)	15.5 (1.4)	20.1 (1.7)	25.1 (2.5)	28.6 (3.4)
	19-30	77	14.9 (1.3)	11.8 (2.8) ^E	12.4 (2.5) ^E	13.3 (1.9)	14.4 (1.6)	15.5 (2.1)	16.5 (3.2) ^E	17.2 (4.0) ^E
	31-50	145	16.4 (1.2)	9.0 (1.4)	10.3 (1.3)	12.8 (1.3)	16.1 (1.5)	19.8 (1.9)	23.4 (2.4)	25.7 (2.8)
	51-70	182	12.7 (1.0)	7.2 (2.0) ^E	8.0 (1.8) ^E	9.4 (1.5)	11.3 (1.2)	13.5 (1.5)	15.7 (2.5)	17.2 (3.4) ^E
	>70	63	13.8 (0.9)	9.4 (2.0) ^E	10.6 (1.7)	12.8 (1.4)	15.3 (1.2)	17.9 (1.6)	20.3 (2.1)	21.8 (2.6)
	19+	467	14.8 (0.7)	7.9 (0.8)	9.0 (0.8)	11.2 (0.7)	14.1 (0.9)	17.9 (1.1)	21.4 (1.3)	23.2 (1.4)
Female	9-13	96	11.9 (0.7)	8.7 (1.3)	9.4 (1.2)	10.7 (1.1)	12.3 (1.0)	14.2 (1.3)	16.2 (1.9)	17.4 (2.4)
	14-18	105	13.2 (1.3)	8.0 (1.9) ^E	8.7 (1.9) ^E	10.3 (1.8) ^E	12.5 (1.7)	15.2 (2.0)	17.8 (2.7)	19.5 (3.2)
	19-30	91	11.7 (1.4)	8.3 (1.2)	8.9 (1.2)	10.0 (1.4)	11.2 (1.5)	12.7 (1.7)	14.2 (2.0)	15.1 (2.2)
	31-50	167	11.3 (0.8)	5.2 (1.0) ^E	6.1 (1.0)	8.0 (1.0)	10.8 (1.0)	14.3 (1.3)	18.3 (1.9)	21.0 (2.5)
	51-70	198	9.9 (0.7)	4.0 (1.0) ^E	4.9 (0.9) ^E	6.6 (0.9)	9.6 (0.9)	13.5 (1.3)	17.2 (1.8)	19.6 (2.2)
	>70	74	10.9 (1.1)	6.3 (1.2) ^E	7.2 (1.1)	8.9 (1.2)	11.2 (1.3)	14.0 (1.8)	17.1 (2.4)	19.3 (3.0)
	19+	530	10.9 (0.5)	5.4 (0.5)	6.3 (0.5)	8.0 (0.5)	10.5 (0.5)	13.6 (0.7)	16.9 (1.0)	19.1 (1.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.2 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	6.1 (0.5)	3.4 (0.8) ^E	3.9 (0.8) ^E	4.6 (0.7)	5.6 (0.6)	6.6 (0.7)	7.6 (0.8)	8.3 (0.9)
	4-8	110	9.4 (0.4)	7.0 (0.9)	7.5 (0.8)	8.3 (0.6)	9.3 (0.6)	10.4 (0.7)	11.4 (1.1)	11.9 (1.3)
Male	9-13	95	12.0 (1.0)	8.0 (1.4) ^E	8.7 (1.3)	10.0 (1.2)	11.6 (1.2)	13.5 (1.5)	15.5 (2.0)	16.9 (2.4)
	14-18	87	16.9 (1.4)	12.3 (2.5) ^E	13.3 (2.3) ^E	15.1 (2.0)	17.3 (1.8)	19.7 (2.1)	22.2 (2.9)	23.8 (3.7)
	19-30	70	21.1 (1.7)	15.1 (2.7) ^E	16.5 (2.6)	19.2 (2.3)	22.8 (2.3)	26.8 (2.7)	30.5 (3.4)	32.8 (4.0)
	31-50	109	17.2 (1.4)	13.2 (2.5) ^E	14.2 (2.3)	16.0 (2.1)	18.2 (1.9)	20.7 (2.2)	23.0 (2.8)	24.5 (3.4)
	51-70	128	14.1 (0.9)	9.0 (1.8) ^E	10.1 (1.7)	12.0 (1.4)	14.6 (1.3)	17.5 (1.6)	20.9 (2.4)	23.4 (3.2)
	>70	65	11.4 (0.8)	7.2 (1.3) ^E	8.0 (1.2)	9.2 (1.1)	10.6 (1.1)	12.4 (1.5)	14.5 (2.1)	16.0 (2.7) ^E
	19+	372	16.5 (0.7)	10.3 (1.0)	11.6 (1.0)	14.0 (1.0)	17.1 (1.1)	20.8 (1.2)	24.5 (1.5)	27.0 (1.8)
Female	9-13	75	12.9 (1.5)	5.7 (1.3) ^E	6.7 (1.3) ^E	8.9 (1.3)	12.1 (1.6)	16.5 (2.3)	21.7 (3.6)	25.6 (4.7) ^E
	14-18	81	12.6 (1.1)	5.9 (1.3) ^E	7.0 (1.3) ^E	9.1 (1.3)	12.0 (1.4)	15.8 (1.9)	20.5 (2.9)	24.1 (4.0) ^E
	19-30	101	11.5 (0.8)	8.2 (0.9)	8.9 (1.0)	10.3 (1.1)	12.1 (1.3)	14.0 (1.5)	16.0 (1.8)	17.3 (2.0)
	31-50	116	11.9 (1.2)	6.9 (1.4) ^E	7.8 (1.3) ^E	9.5 (1.2)	11.7 (1.2)	14.2 (1.6)	16.7 (2.3)	18.4 (2.8)
	51-70	146	10.7 (0.8)	6.1 (1.1) ^E	6.7 (1.0)	8.1 (0.9)	10.0 (0.7)	12.3 (1.1)	14.6 (1.7)	16.0 (2.3)
	>70	94	10.1 (1.0)	6.2 (1.2) ^E	6.8 (1.1)	8.0 (1.0)	9.6 (0.9)	11.4 (1.1)	13.3 (1.8)	14.6 (2.3)
	19+	457	11.3 (0.6)	6.7 (0.7)	7.5 (0.7)	9.1 (0.6)	11.1 (0.6)	13.6 (0.9)	16.2 (1.4)	17.9 (1.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.3 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	
Both											
	1-3	112	6.7 (0.3)	3.8 (0.7) ^E	4.4 (0.6)	5.3 (0.5)	6.5 (0.5)	7.8 (0.6)	9.2 (0.8)	10.1 (0.9)	
	4-8	177	9.8 (0.6)	7.0 (1.4) ^E	7.6 (1.2)	8.8 (1.0)	10.2 (0.8)	11.7 (1.1)	13.2 (1.7)	14.2 (2.2)	
Male	9-13	111	14.0 (0.8)	9.8 (1.5)	10.6 (1.4)	12.0 (1.1)	13.6 (1.0)	15.4 (1.3)	17.0 (1.8)	18.0 (2.2)	
	14-18	113	16.7 (1.7)	10.1 (2.7) ^E	11.2 (2.7) ^E	13.3 (2.5) ^E	16.2 (2.3)	19.8 (2.5)	23.5 (3.3)	25.9 (4.1)	
	19-30	91	15.0 (1.1)	7.8 (1.9) ^E	8.8 (1.7) ^E	10.9 (1.5)	13.7 (1.4)	17.0 (1.9)	20.7 (3.1)	23.2 (4.2) ^E	
	31-50	101	16.9 (1.2)	8.0 (1.8) ^E	9.4 (1.7) ^E	12.1 (1.6)	15.8 (1.7)	20.0 (2.1)	24.2 (2.6)	26.8 (3.1)	
	51-70	134	15.3 (1.4)	^F	9.6 (2.3) ^E	12.2 (1.9)	14.6 (1.7)	18.5 (2.4)	22.4 (3.8) ^E	24.3 (4.6) ^E	
	>70	56	10.2 (1.0)	7.3 (1.3) ^E	7.8 (1.2)	8.7 (1.0)	9.7 (1.1)	10.8 (1.5)	12.0 (2.2) ^E	12.8 (2.8) ^E	
	19+	382	15.4 (0.7)	7.4 (1.1)	8.7 (1.0)	11.2 (0.8)	14.4 (0.8)	18.4 (1.2)	22.2 (1.8)	24.5 (2.2)	
	Female	9-13	105	11.5 (0.8)	9.5 (1.9) ^E	10.0 (1.7) ^E	10.8 (1.3)	11.8 (1.0)	12.9 (1.1)	13.9 (1.7)	14.5 (2.2)
		14-18	120	11.2 (1.3)	^F	6.3 (1.8) ^E	8.2 (1.5) ^E	10.6 (1.4)	13.5 (1.6)	16.3 (2.4)	18.3 (3.1) ^E
19-30		91	13.9 (1.6)	^F	7.5 (2.1) ^E	9.7 (2.0) ^E	12.5 (2.1) ^E	16.1 (2.5)	20.6 (3.9) ^E	24.2 (5.4) ^E	
31-50		159	13.7 (1.2)	8.5 (1.8) ^E	9.5 (1.7) ^E	11.4 (1.5)	13.7 (1.5)	16.3 (1.8)	18.8 (2.4)	20.4 (2.8)	
51-70		174	13.1 (1.3)	8.6 (1.9) ^E	9.4 (1.8) ^E	10.7 (1.6)	12.5 (1.5)	14.4 (1.7)	16.2 (2.4)	17.4 (3.0) ^E	
>70		80	8.7 (0.6)	5.5 (1.0) ^E	6.2 (0.9)	7.4 (0.8)	8.7 (0.8)	10.0 (1.0)	11.5 (1.5)	12.8 (2.0)	
19+		504	12.9 (0.7)	6.7 (0.8)	7.8 (0.8)	9.8 (0.8)	12.7 (0.9)	16.1 (1.2)	19.9 (1.6)	22.4 (1.9)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.4 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	7.7 (0.8)	4.7 (0.8) ^E	5.3 (0.8)	6.3 (0.8)	7.6 (0.9)	9.1 (1.1)	10.5 (1.4)	11.4 (1.5)
	4-8	140	10.0 (0.4)	7.3 (0.4)	7.8 (0.4)	8.7 (0.4)	9.8 (0.5)	11.0 (0.6)	12.2 (0.7)	13.0 (0.8)
Male	9-13	92	19.6 (4.5) ^E	F	10.2 (2.9) ^E	13.1 (3.0) ^E	17.8 (3.7) ^E	23.2 (5.1) ^E	29.5 (7.1) ^E	33.7 (8.7) ^E
	14-18	107	17.7 (1.5)	11.1 (2.2) ^E	12.3 (2.2) ^E	14.5 (2.1)	17.3 (2.1)	20.6 (2.2)	24.1 (2.8)	26.3 (3.4)
	19-30	73	17.4 (1.2)	9.1 (1.7) ^E	10.8 (1.6)	13.9 (1.5)	18.1 (1.7)	23.0 (2.4)	28.0 (3.5)	31.3 (4.2)
	31-50	134	15.8 (1.0)	8.0 (1.9) ^E	9.2 (1.8) ^E	11.5 (1.5)	14.6 (1.3)	18.5 (1.6)	22.7 (2.5)	25.6 (3.3)
	51-70	131	16.2 (1.5)	10.5 (1.1)	11.5 (1.1)	13.3 (1.3)	15.8 (1.6)	18.6 (2.1)	21.6 (2.7)	23.6 (3.2)
	>70	55	13.5 (1.3)	7.1 (2.2) ^E	8.3 (2.1) ^E	10.5 (1.9) ^E	13.4 (1.8)	16.8 (2.0)	20.2 (2.5)	22.5 (3.1)
	19+	393	16.0 (0.7)	8.4 (0.8)	9.6 (0.8)	12.1 (0.8)	15.4 (0.8)	19.7 (1.0)	24.6 (1.6)	28.1 (2.2)
Female	9-13	79	12.6 (1.5)	7.8 (1.6) ^E	8.9 (1.5) ^E	10.7 (1.4)	13.0 (1.6)	15.6 (2.3)	18.2 (3.5) ^E	20.0 (4.4) ^E
	14-18	104	11.7 (1.2)	7.2 (0.9)	8.0 (1.0)	9.5 (1.2)	11.5 (1.4)	13.9 (1.7)	16.5 (2.1)	18.1 (2.4)
	19-30	101	12.5 (1.1)	8.8 (2.1) ^E	9.8 (1.9) ^E	11.5 (1.6)	13.5 (1.4)	15.7 (1.5)	18.0 (2.1)	19.5 (2.7)
	31-50	143	10.1 (0.5)	6.2 (0.5)	6.9 (0.6)	8.3 (0.6)	10.0 (0.7)	11.9 (0.7)	13.7 (0.8)	15.0 (0.9)
	51-70	193	11.3 (0.5)	7.6 (1.2)	8.3 (1.0)	9.5 (0.8)	11.0 (0.7)	12.6 (0.8)	14.2 (1.2)	15.3 (1.6)
	>70	94	8.7 (0.6)	4.6 (0.6)	5.2 (0.6)	6.3 (0.6)	7.8 (0.8)	9.7 (1.0)	11.9 (1.3)	13.4 (1.6)
	19+	531	10.7 (0.4)	6.7 (0.7)	7.4 (0.6)	8.9 (0.5)	10.7 (0.4)	12.7 (0.5)	14.7 (0.7)	16.0 (0.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.5 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake							
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	95th (SE)
Both	1-3	311	7.0 (0.3)	3.9 (0.5)	4.5 (0.5)	5.6 (0.5)	7.0 (0.5)	8.5 (0.6)	10.1 (0.8)
	4-8	485	10.6 (0.4)	6.2 (0.6)	7.0 (0.6)	8.4 (0.5)	10.6 (0.5)	13.1 (0.7)	15.7 (1.1)
Male	9-13	277	16.1 (1.0)	9.9 (1.3)	11.2 (1.3)	13.6 (1.3)	16.8 (1.4)	20.5 (1.8)	27.6 (2.9)
	14-18	339	17.6 (0.8)	11.2 (1.6)	12.4 (1.5)	14.8 (1.3)	17.8 (1.2)	21.4 (1.4)	25.3 (2.2)
	19-30	237	17.4 (0.9)	13.4 (1.9)	14.2 (1.7)	15.7 (1.5)	17.6 (1.4)	19.7 (1.7)	21.8 (2.3)
	31-50	423	16.1 (1.2)	8.9 (1.5) ^E	10.2 (1.5)	12.6 (1.4)	15.8 (1.4)	19.5 (1.7)	23.5 (2.4)
	51-70	387	15.0 (0.6)	7.1 (0.8)	8.4 (0.8)	10.8 (0.7)	13.9 (0.8)	18.2 (1.0)	22.9 (1.3)
	>70	132	12.5 (0.9)	7.4 (1.6) ^E	8.5 (1.5) ^E	10.5 (1.4)	13.1 (1.3)	15.9 (1.4)	18.5 (1.7)
	19+	1179	15.8 (0.6)	8.6 (0.7)	9.9 (0.6)	12.3 (0.7)	15.4 (0.7)	19.2 (0.9)	23.3 (1.3)
Female	9-13	281	12.8 (0.7)	7.9 (1.0)	8.8 (1.0)	10.4 (0.8)	12.4 (0.8)	14.9 (1.1)	17.7 (1.6)
	14-18	321	13.9 (1.0)	10.3 (1.7)	11.2 (1.6)	12.8 (1.5)	14.9 (1.4)	17.2 (1.7)	19.5 (2.3)
	19-30	249	12.9 (0.9)	7.5 (0.7)	8.4 (0.7)	10.0 (0.8)	12.1 (0.9)	14.5 (1.2)	17.2 (1.4)
	31-50	364	12.8 (0.8)	9.2 (1.4)	9.9 (1.3)	11.3 (1.1)	13.1 (1.0)	15.4 (1.4)	17.9 (2.4)
	51-70	467	12.3 (0.6)	6.2 (0.8)	7.3 (0.8)	9.4 (0.7)	12.0 (0.6)	15.0 (0.8)	18.2 (1.2)
	>70	215	9.9 (0.6)	7.1 (1.2)	7.6 (1.1)	8.6 (1.0)	9.9 (1.0)	11.4 (1.2)	13.0 (1.6)
	19+	1295	12.3 (0.4)	7.9 (0.6)	8.7 (0.5)	10.2 (0.5)	12.2 (0.4)	14.7 (0.6)	17.3 (1.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.6 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	6.4 (0.2)	3.2 (0.6) ^E	3.8 (0.5)	4.8 (0.4)	6.2 (0.3)	7.9 (0.4)	9.6 (0.8)	10.9 (1.1)
	4-8	956	9.7 (0.3)	6.2 (0.7)	6.8 (0.6)	8.0 (0.5)	9.5 (0.3)	11.3 (0.4)	13.2 (0.8)	14.6 (1.2)
Male	9-13	589	13.0 (0.4)	7.9 (0.9)	8.8 (0.8)	10.4 (0.6)	12.6 (0.5)	15.2 (0.6)	18.1 (1.1)	20.0 (1.5)
	14-18	639	15.8 (0.5)	10.3 (1.6)	11.4 (1.3)	13.3 (0.9)	15.8 (0.6)	18.5 (1.0)	21.3 (1.7)	23.1 (2.3)
	19-30	481	16.8 (0.9)	7.6 (1.6) ^E	9.1 (1.5)	12.0 (1.2)	15.9 (1.0)	20.6 (1.4)	26.0 (2.5)	30.2 (3.6)
	31-50	709	14.7 (0.6)	10.2 (1.8) ^E	11.0 (1.6)	12.5 (1.1)	14.4 (0.7)	16.5 (1.1)	18.5 (2.2)	19.8 (3.0)
	51-70	758	14.9 (0.8)	7.0 (1.4) ^E	8.2 (1.3)	10.5 (1.1)	13.8 (0.8)	18.2 (0.9)	23.3 (1.9)	27.2 (2.8)
	>70	734	11.2 (0.4)	5.0 (0.7)	6.0 (0.6)	7.9 (0.5)	10.4 (0.5)	13.7 (0.6)	17.4 (1.0)	20.1 (1.5)
	19+	2682	14.9 (0.4)	7.2 (0.6)	8.3 (0.6)	10.7 (0.5)	14.1 (0.4)	18.2 (0.6)	22.9 (1.0)	26.2 (1.4)
Female	9-13	585	11.7 (0.5)	6.5 (0.9)	7.4 (0.8)	9.1 (0.7)	11.3 (0.5)	14.1 (0.7)	17.1 (1.2)	19.4 (1.8)
	14-18	645	12.4 (0.4)	6.9 (1.0)	7.9 (0.9)	9.7 (0.7)	12.1 (0.5)	15.0 (0.7)	18.0 (1.4)	20.0 (1.8)
	19-30	514	10.7 (0.5)	5.7 (1.1) ^E	6.5 (1.0)	8.1 (0.8)	10.1 (0.6)	12.6 (0.8)	15.4 (1.4)	17.4 (1.9)
	31-50	758	12.2 (0.5)	5.7 (0.7)	6.8 (0.6)	8.8 (0.6)	11.6 (0.5)	14.9 (0.7)	18.5 (1.1)	21.1 (1.5)
	51-70	955	11.0 (0.4)	5.8 (0.9)	6.6 (0.8)	8.3 (0.7)	10.6 (0.6)	13.4 (0.6)	16.6 (1.1)	18.8 (1.6)
	>70	1345	10.0 (0.3)	5.4 (0.7)	6.2 (0.6)	7.6 (0.5)	9.5 (0.3)	11.8 (0.5)	14.3 (0.9)	16.0 (1.3)
	19+	3572	11.3 (0.3)	5.8 (0.4)	6.7 (0.4)	8.4 (0.4)	10.8 (0.3)	13.7 (0.4)	16.9 (0.6)	19.1 (0.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.7 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	6.0	(0.3)	2.5 (0.6) ^E	3.1 (0.5)	4.3 (0.4)	5.6 (0.4)	7.2 (0.4)	9.0 (0.7)	10.2 (0.9)
	4-8	425	9.8	(0.4)	5.5 (1.1) ^E	6.3 (1.0)	7.6 (0.8)	9.4 (0.5)	11.6 (0.6)	14.0 (1.2)	15.8 (1.8)
Male	9-13	274	13.7	(0.7)	7.1 (1.7) ^E	8.2 (1.5) ^E	10.3 (1.2)	13.2 (0.9)	16.7 (1.1)	20.8 (2.0)	23.8 (2.9)
	14-18	297	18.0	(1.1)	9.7 (2.0) ^E	11.1 (1.8) ^E	13.9 (1.6)	17.6 (1.4)	22.0 (1.8)	26.8 (2.8)	30.1 (3.6)
	19-30	249	18.6	(1.4)	10.8 (0.8)	12.0 (0.9)	14.6 (1.3)	18.4 (1.6)	23.1 (2.0)	28.2 (2.6)	31.6 (3.2)
	31-50	309	13.9	(0.9)	6.8 (1.3) ^E	7.8 (1.3)	10.0 (1.1)	12.9 (1.0)	16.8 (1.4)	21.8 (2.6)	25.7 (3.7)
	51-70	277	13.1	(0.7)	9.1 (1.7) ^E	9.9 (1.5)	11.2 (1.2)	12.9 (0.9)	14.8 (1.2)	16.7 (2.0)	17.9 (2.6)
	>70	136	11.9	(1.2)	6.3 (1.5) ^E	7.1 (1.4) ^E	8.7 (1.3)	10.8 (1.2)	13.5 (2.3) ^E	16.6 (3.0) ^E	18.8 (3.6) ^E
	19+	971	14.6	(0.6)	9.0 (1.7) ^E	10.0 (1.5)	12.0 (1.1)	14.4 (0.7)	17.4 (1.0)	20.7 (2.1)	23.0 (2.9)
	Female	9-13	265	11.2	(0.5)	6.4 (1.1) ^E	7.2 (0.9)	8.7 (0.7)	10.5 (0.5)	12.6 (0.7)	14.7 (1.2)
14-18	290	11.7	(0.8)	6.1 (1.3) ^E	7.1 (1.1)	9.0 (0.9)	11.4 (0.8)	14.2 (1.1)	17.3 (1.8)	19.5 (2.5)	
19-30	197	11.3	(0.8)	7.1 (0.7)	7.9 (0.7)	9.3 (0.8)	11.1 (0.9)	13.1 (1.0)	15.1 (1.1)	16.3 (1.2)	
31-50	312	11.6	(0.6)	8.0 (1.3)	8.7 (1.2)	9.9 (0.9)	11.5 (0.8)	13.3 (1.0)	15.2 (1.7)	16.4 (2.3)	
51-70	312	10.3	(0.5)	5.6 (0.9)	6.3 (0.8)	7.8 (0.7)	9.8 (0.6)	12.3 (0.8)	15.1 (1.3)	16.9 (1.7)	
>70	239	8.6	(0.5)	4.6 (0.9) ^E	5.2 (0.8)	6.4 (0.7)	8.0 (0.5)	9.9 (0.7)	11.9 (1.2)	13.2 (1.6)	
19+	1060	10.7	(0.3)	6.2 (0.8)	7.0 (0.7)	8.4 (0.5)	10.4 (0.4)	12.7 (0.6)	15.1 (1.1)	16.8 (1.4)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.8 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	7.0 (0.5)	4.1 (0.9) ^E	4.6 (0.8) ^E	5.8 (0.7)	7.2 (0.6)	8.8 (0.8)	10.4 (1.1)	11.5 (1.4)
	4-8	213	11.5 (0.9)	6.5 (0.8)	7.3 (0.8)	8.8 (0.8)	10.8 (0.8)	13.4 (1.2)	16.2 (1.7)	18.1 (2.2)
Male	9-13	122	18.2 (5.2) ^E	8.6 (2.4) ^E	9.7 (2.5) ^E	13.2 (2.8) ^E	16.5 (3.5) ^E	21.7 (5.0) ^E	26.7 (6.7) ^E	31.1 (8.4) ^E
	14-18	150	19.1 (1.4)	12.6 (1.2)	13.9 (1.3)	16.3 (1.5)	19.4 (1.9)	22.9 (2.4)	26.5 (3.1)	28.9 (3.6)
	19-30	106	18.9 (1.8)	8.4 (2.7) ^E	10.1 (2.6) ^E	13.4 (2.3) ^E	18.0 (2.1)	23.5 (2.9)	29.5 (4.7)	33.5 (6.5) ^E
	31-50	155	17.5 (1.6)	9.8 (2.1) ^E	11.2 (2.0) ^E	13.9 (1.9)	17.2 (2.0)	21.3 (2.5)	25.7 (3.5)	28.7 (4.3)
	51-70	122	15.5 (1.3)	8.9 (2.0) ^E	10.2 (1.8) ^E	12.7 (1.6)	16.0 (1.6)	20.0 (2.2)	24.4 (3.6)	27.6 (4.9) ^E
	>70	88	14.3 (1.1)	7.1 (1.2) ^E	8.2 (1.2)	10.4 (1.1)	13.5 (1.1)	17.3 (1.8)	21.6 (2.9)	24.6 (3.8)
	19+	471	16.9 (0.8)	8.0 (1.0)	9.6 (0.9)	12.5 (0.9)	16.3 (1.0)	21.1 (1.3)	27.0 (1.9)	31.4 (2.6)
Female	9-13	103	11.4 (0.8)	8.7 (1.5) ^E	9.3 (1.4)	10.3 (1.2)	11.6 (1.1)	13.1 (1.2)	14.5 (1.5)	15.5 (1.8)
	14-18	142	12.1 (0.6)	7.5 (1.1)	8.4 (1.0)	10.0 (0.9)	12.1 (0.9)	14.5 (1.2)	17.1 (1.8)	18.8 (2.3)
	19-30	111	12.2 (0.7)	9.3 (1.8) ^E	9.9 (1.6) ^E	11.0 (1.3)	12.4 (1.1)	14.0 (1.6)	15.5 (2.6) ^E	16.6 (3.5) ^E
	31-50	146	10.8 (0.8)	5.5 (1.8) ^E	6.2 (1.6) ^E	7.7 (1.3) ^E	9.6 (1.1)	12.0 (1.1)	14.5 (1.7)	16.3 (2.3)
	51-70	184	12.8 (0.8)	6.9 (1.8) ^E	7.9 (1.6) ^E	9.8 (1.4)	12.4 (1.2)	15.6 (1.4)	19.2 (2.3)	21.7 (3.2)
	>70	143	10.5 (0.8)	6.9 (1.4) ^E	7.7 (1.3) ^E	9.1 (1.2)	10.9 (1.2)	13.1 (1.5)	15.5 (2.1)	17.0 (2.6)
	19+	584	11.6 (0.4)	6.4 (0.9)	7.2 (0.9)	8.9 (0.7)	11.1 (0.6)	13.8 (0.7)	16.7 (1.1)	18.7 (1.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.9 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	169	5.5	(0.4)	F	2.8	(0.8) ^E	3.9	(0.7) ^E	5.3	(0.5)	7.1	(0.7)	9.0	(1.3)	10.4	(1.8) ^E	
	4-8	281	9.5	(0.4)	6.0	(1.1) ^E	6.6	(0.9)	7.8	(0.7)	9.2	(0.5)	10.8	(0.7)	12.5	(1.2)	13.5	(1.7)
Male	9-13	183	14.4	(0.9)	8.8	(1.8) ^E	9.8	(1.6) ^E	11.6	(1.4)	14.2	(1.2)	17.5	(1.5)	21.0	(2.5)	23.5	(3.3)
	14-18	187	16.6	(1.2)	10.8	(1.9) ^E	11.9	(1.8)	13.9	(1.5)	16.5	(1.3)	19.6	(1.7)	22.8	(2.8)	25.0	(3.9)
	19-30	223	14.9	(0.9)	6.9	(1.3) ^E	8.2	(1.2)	10.9	(1.1)	14.5	(1.0)	18.9	(1.4)	23.6	(2.2)	27.0	(2.9)
	31-50	229	16.9	(1.6)	9.8	(1.1)	11.2	(1.1)	13.8	(1.3)	17.2	(1.6)	21.2	(2.2)	25.4	(3.1)	28.3	(3.7)
	51-70	197	13.6	(0.9)	7.6	(1.8) ^E	8.7	(1.7) ^E	10.7	(1.4)	13.4	(1.2)	16.6	(1.6)	20.1	(2.5)	22.5	(3.3)
	>70	72	12.8	(1.1)	7.2	(1.7) ^E	8.2	(1.5) ^E	10.1	(1.3)	12.7	(1.2)	15.6	(1.7)	18.7	(2.7)	20.9	(3.6) ^E
	19+	721	15.3	(0.8)	8.6	(1.6) ^E	9.9	(1.5)	12.3	(1.2)	15.4	(0.9)	19.2	(1.1)	23.2	(1.7)	25.9	(2.3)
	Female	9-13	165	11.1	(0.9)	7.2	(0.8)	7.9	(0.9)	9.3	(1.0)	11.2	(1.2)	13.4	(1.4)	15.8	(1.7)	17.4
14-18	206	11.9	(0.7)	7.4	(1.6) ^E	8.2	(1.5) ^E	9.7	(1.2)	11.6	(0.9)	13.7	(1.2)	15.9	(1.9)	17.4	(2.6)	
19-30	191	11.2	(0.7)	5.7	(1.2) ^E	6.7	(1.2) ^E	8.5	(1.0)	10.8	(1.0)	13.8	(1.4)	17.1	(2.2)	19.7	(3.1)	
31-50	258	12.1	(0.8)	8.6	(0.7)	9.2	(0.8)	10.5	(0.9)	12.1	(1.1)	13.8	(1.2)	15.6	(1.4)	16.7	(1.6)	
51-70	249	10.7	(0.6)	5.2	(0.9) ^E	6.1	(0.9)	7.9	(0.8)	10.3	(0.7)	13.2	(0.9)	16.3	(1.4)	18.4	(1.8)	
>70	128	8.8	(1.0)	3.6	(1.0) ^E	4.4	(1.0) ^E	6.1	(1.1) ^E	8.5	(1.3)	11.2	(1.6)	15.1	(2.3)	19.0	(3.3) ^E	
19+	826	11.2	(0.4)	5.9	(0.7)	6.8	(0.7)	8.6	(0.6)	11.0	(0.6)	13.9	(0.6)	17.0	(0.9)	19.1	(1.2)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.10 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	192	7.2 (0.5)	4.1 (0.7) ^E	4.6 (0.7)	5.5 (0.6)	6.7 (0.6)	8.0 (0.8)	9.3 (1.0)	10.2 (1.2)
	4-8	321	10.4 (0.6)	5.7 (0.4)	6.5 (0.4)	8.0 (0.5)	10.0 (0.7)	12.5 (1.0)	15.2 (1.3)	17.2 (1.6)
Male	9-13	226	14.6 (0.8)	7.7 (1.2)	8.8 (1.2)	11.1 (1.1)	14.3 (1.1)	18.2 (1.4)	22.7 (2.1)	26.1 (2.8)
	14-18	262	20.3 (1.9)	11.2 (1.2)	12.8 (1.3)	15.8 (1.5)	19.7 (1.8)	24.2 (2.4)	28.9 (3.1)	32.0 (3.6)
	19-30	197	18.9 (1.2)	12.2 (1.8)	13.5 (1.7)	15.9 (1.4)	19.1 (1.5)	23.0 (2.1)	27.1 (3.3)	29.9 (4.2)
	31-50	282	18.5 (1.1)	8.7 (1.6) ^E	10.2 (1.5)	13.3 (1.4)	17.6 (1.4)	23.1 (1.9)	29.3 (2.9)	33.5 (3.8)
	51-70	234	15.3 (0.9)	7.7 (1.5) ^E	8.9 (1.4)	11.2 (1.2)	14.3 (1.1)	18.3 (1.4)	22.7 (2.4)	26.0 (3.4)
	>70	119	13.1 (1.0)	5.8 (1.0) ^E	6.9 (1.0)	9.2 (1.2)	12.3 (1.3)	16.1 (1.7)	20.3 (2.1)	23.2 (2.5)
	19+	832	17.1 (0.6)	8.5 (0.6)	9.9 (0.6)	12.7 (0.6)	16.5 (0.7)	21.3 (0.9)	26.7 (1.4)	30.6 (1.8)
Female	9-13	226	11.4 (0.8)	7.9 (1.2)	8.6 (1.1)	9.9 (1.0)	11.6 (0.9)	13.4 (1.1)	15.3 (1.6)	16.5 (2.0)
	14-18	242	12.2 (0.7)	6.6 (1.3) ^E	7.6 (1.2)	9.4 (1.0)	11.9 (1.0)	14.7 (1.1)	17.7 (1.5)	19.8 (1.9)
	19-30	208	12.6 (0.8)	5.2 (0.9) ^E	6.3 (0.9)	8.7 (0.9)	11.8 (1.0)	15.3 (1.3)	19.2 (1.9)	22.3 (2.6)
	31-50	263	13.6 (0.9)	7.3 (1.2)	8.3 (1.1)	10.3 (0.9)	13.1 (1.0)	16.6 (1.4)	20.4 (2.1)	23.2 (2.7)
	51-70	322	11.1 (0.7)	4.9 (0.9) ^E	5.9 (0.9)	7.7 (0.8)	10.4 (0.8)	13.8 (1.1)	17.6 (1.7)	20.2 (2.1)
	>70	198	9.6 (0.5)	5.4 (0.9)	6.1 (0.8)	7.3 (0.7)	9.0 (0.7)	11.0 (0.9)	13.3 (1.3)	14.9 (1.7)
	19+	991	12.2 (0.4)	5.5 (0.4)	6.6 (0.4)	8.7 (0.4)	11.5 (0.5)	15.1 (0.7)	19.3 (1.0)	22.3 (1.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.11 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	348	7.0 (0.3)	3.9 (0.4)	4.5 (0.4)	5.5 (0.4)	6.7 (0.4)	8.3 (0.5)	9.9 (0.7)	10.9 (0.8)
	4-8	554	10.3 (0.3)	7.3 (0.8)	7.9 (0.7)	9.1 (0.5)	10.6 (0.5)	12.1 (0.6)	13.7 (0.9)	14.7 (1.1)
Male	9-13	409	16.1 (1.5)	9.7 (1.1)	10.9 (1.1)	13.2 (1.2)	16.1 (1.5)	19.3 (2.1)	22.7 (2.8)	24.9 (3.5)
	14-18	414	16.8 (0.9)	10.2 (1.1)	11.4 (1.1)	13.6 (1.1)	16.6 (1.1)	20.2 (1.4)	23.9 (1.8)	26.4 (2.1)
	19-30	311	16.1 (0.7)	9.1 (1.0)	10.3 (0.9)	12.6 (0.8)	15.5 (0.9)	19.1 (1.2)	22.9 (1.7)	25.5 (2.2)
	31-50	489	16.4 (0.6)	8.5 (0.7)	9.8 (0.7)	12.3 (0.7)	15.7 (0.8)	19.8 (1.1)	23.8 (1.3)	26.4 (1.5)
	51-70	575	14.9 (0.8)	9.4 (1.7) ^E	10.5 (1.5)	12.5 (1.1)	14.8 (0.9)	17.3 (1.3)	20.0 (2.0)	21.7 (2.6)
	>70	239	12.1 (0.6)	6.7 (0.7)	7.7 (0.7)	9.5 (0.7)	12.0 (0.8)	14.8 (1.0)	17.8 (1.3)	19.7 (1.6)
	19+	1614	15.5 (0.4)	7.9 (0.4)	9.2 (0.4)	11.6 (0.4)	14.9 (0.5)	18.9 (0.6)	23.0 (0.8)	25.8 (1.0)
Female	9-13	355	12.0 (0.6)	^F	6.3 (1.9) ^E	8.2 (1.6) ^E	10.6 (1.3)	13.4 (1.6)	16.3 (2.7)	18.3 (3.6) ^E
	14-18	410	11.9 (0.7)	6.3 (0.8)	7.2 (0.8)	9.0 (0.8)	11.5 (0.9)	14.4 (1.0)	17.5 (1.4)	19.6 (1.8)
	19-30	384	12.8 (0.8)	8.8 (1.3)	9.6 (1.2)	11.2 (1.0)	13.1 (1.0)	15.2 (1.3)	17.4 (1.7)	18.8 (2.1)
	31-50	585	11.9 (0.6)	6.8 (0.8)	7.7 (0.7)	9.5 (0.7)	11.8 (0.7)	14.4 (0.9)	17.0 (1.2)	18.7 (1.5)
	51-70	711	11.7 (0.6)	6.1 (0.6)	7.0 (0.6)	8.8 (0.6)	11.2 (0.7)	13.9 (0.8)	16.8 (1.1)	18.7 (1.3)
	>70	342	9.2 (0.4)	5.1 (0.5)	5.8 (0.5)	7.1 (0.5)	8.8 (0.5)	11.0 (0.6)	13.5 (0.9)	15.3 (1.2)
	19+	2022	11.7 (0.3)	6.4 (0.4)	7.3 (0.4)	9.1 (0.4)	11.5 (0.4)	14.4 (0.5)	17.4 (0.7)	19.4 (0.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.12 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	622	5.9	(0.3)	2.6 (0.4) ^E	3.1 (0.4)	4.2 (0.3)	5.6 (0.3)	7.4 (0.4)	9.3 (0.7)	10.6 (0.9)
	4-8	919	9.9	(0.3)	5.1 (0.5)	5.9 (0.4)	7.4 (0.4)	9.4 (0.3)	11.7 (0.5)	14.3 (0.8)	16.3 (1.1)
Male	9-13	579	14.9	(1.1)	8.1 (1.0)	9.1 (1.0)	11.2 (0.9)	14.2 (1.0)	18.3 (1.4)	23.2 (2.2)	26.9 (3.1)
	14-18	634	17.4	(0.8)	11.6 (1.7)	12.8 (1.5)	14.9 (1.2)	17.6 (1.0)	20.7 (1.2)	24.0 (2.0)	26.3 (2.7)
	19-30	578	16.3	(0.7)	9.1 (1.8) ^E	10.4 (1.6)	13.0 (1.2)	16.3 (0.9)	20.1 (1.3)	24.1 (2.2)	27.0 (3.1)
	31-50	693	16.4	(1.1)	9.5 (2.1) ^E	10.7 (1.9) ^E	13.2 (1.5)	16.4 (1.1)	20.2 (1.4)	24.2 (2.3)	27.0 (3.1)
	51-70	596	13.8	(0.6)	7.5 (1.3) ^E	8.7 (1.2)	10.9 (1.0)	13.7 (0.8)	17.1 (1.0)	20.7 (1.6)	23.2 (2.2)
	>70	296	12.9	(0.7)	6.7 (0.8)	7.7 (0.7)	9.6 (0.7)	12.3 (0.8)	15.5 (1.0)	19.0 (1.6)	21.4 (2.0)
	19+	2163	15.4	(0.5)	8.2 (0.9)	9.5 (0.8)	11.9 (0.7)	15.3 (0.6)	19.3 (0.7)	23.7 (1.1)	26.8 (1.5)
Female	9-13	533	11.2	(0.6)	7.1 (1.2) ^E	7.9 (1.1)	9.3 (0.9)	11.1 (0.8)	13.2 (0.9)	15.4 (1.4)	16.8 (1.8)
	14-18	638	11.9	(0.5)	6.7 (0.8)	7.7 (0.8)	9.5 (0.6)	11.7 (0.6)	14.2 (0.8)	16.9 (1.2)	18.8 (1.5)
	19-30	499	11.4	(0.5)	7.1 (1.1)	7.9 (0.9)	9.4 (0.8)	11.3 (0.7)	13.5 (1.0)	15.9 (1.5)	17.4 (2.0)
	31-50	716	11.8	(0.5)	8.2 (1.3)	8.8 (1.2)	10.0 (0.9)	11.6 (0.7)	13.3 (0.8)	15.1 (1.2)	16.2 (1.6)
	51-70	745	11.0	(0.4)	5.7 (0.6)	6.6 (0.6)	8.3 (0.5)	10.6 (0.5)	13.5 (0.6)	16.5 (0.9)	18.5 (1.2)
	>70	510	9.1	(0.5)	4.5 (0.6)	5.2 (0.6)	6.8 (0.6)	8.7 (0.7)	11.4 (0.8)	14.3 (1.2)	16.7 (1.5)
	19+	2470	11.2	(0.3)	6.0 (0.4)	6.9 (0.4)	8.5 (0.4)	10.8 (0.4)	13.7 (0.4)	16.7 (0.6)	18.7 (0.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.13 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	6.6 (0.2)	3.2 (0.2)	3.8 (0.2)	4.9 (0.2)	6.3 (0.2)	8.0 (0.2)	9.8 (0.3)	11.1 (0.4)
	4-8	3235	10.1 (0.2)	6.0 (0.3)	6.7 (0.2)	8.1 (0.2)	9.9 (0.2)	12.0 (0.3)	14.3 (0.4)	15.9 (0.6)
Male	9-13	2080	14.5 (0.4)	8.4 (0.4)	9.4 (0.4)	11.4 (0.4)	14.1 (0.4)	17.5 (0.6)	21.4 (0.8)	24.2 (1.1)
	14-18	2288	17.1 (0.4)	10.6 (0.7)	11.9 (0.7)	14.2 (0.6)	17.3 (0.5)	20.8 (0.7)	24.5 (1.0)	27.0 (1.4)
	19-30	1804	17.1 (0.5)	9.9 (0.6)	11.2 (0.6)	13.6 (0.6)	16.8 (0.6)	20.6 (0.8)	24.7 (1.2)	27.6 (1.5)
	31-50	2596	15.9 (0.4)	8.0 (0.5)	9.3 (0.5)	11.8 (0.5)	15.4 (0.5)	19.5 (0.7)	24.0 (1.0)	27.1 (1.3)
	51-70	2550	14.8 (0.4)	7.4 (0.5)	8.6 (0.5)	10.9 (0.4)	14.0 (0.4)	18.1 (0.5)	22.8 (0.8)	26.2 (1.2)
	>70	1520	12.1 (0.3)	6.0 (0.3)	7.0 (0.3)	9.0 (0.4)	11.7 (0.4)	14.9 (0.5)	18.4 (0.6)	20.8 (0.8)
	19+	8470	15.5 (0.2)	7.8 (0.3)	9.1 (0.2)	11.5 (0.3)	14.9 (0.3)	19.1 (0.3)	23.8 (0.5)	27.1 (0.7)
Female	9-13	1980	11.9 (0.3)	7.0 (0.4)	7.9 (0.4)	9.5 (0.3)	11.6 (0.3)	14.2 (0.4)	17.0 (0.6)	19.0 (0.8)
	14-18	2256	12.6 (0.3)	7.4 (0.4)	8.4 (0.4)	10.2 (0.4)	12.5 (0.4)	15.3 (0.5)	18.2 (0.7)	20.1 (0.8)
	19-30	1854	11.7 (0.3)	6.7 (0.4)	7.5 (0.4)	9.1 (0.4)	11.2 (0.4)	13.7 (0.5)	16.3 (0.7)	18.1 (0.9)
	31-50	2686	12.5 (0.3)	6.9 (0.4)	7.9 (0.4)	9.7 (0.4)	12.1 (0.4)	15.1 (0.5)	18.4 (0.8)	20.6 (1.1)
	51-70	3200	11.4 (0.2)	5.6 (0.3)	6.6 (0.3)	8.4 (0.3)	10.9 (0.3)	14.0 (0.4)	17.4 (0.5)	19.7 (0.7)
	>70	2610	9.7 (0.2)	5.3 (0.3)	6.0 (0.3)	7.4 (0.3)	9.4 (0.3)	11.8 (0.4)	14.5 (0.6)	16.5 (0.7)
	19+	10350	11.7 (0.2)	6.2 (0.2)	7.1 (0.2)	8.9 (0.2)	11.3 (0.2)	14.2 (0.2)	17.5 (0.4)	19.7 (0.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

39. Total saturated fats (g/d): Usual intakes from food

Table 39.1 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	23.5 (2.4)	14.6 (3.2) ^E	16.2 (3.0) ^E	19.3 (2.7)	23.2 (2.7)	27.3 (3.5)	31.3 (5.0)	33.8 (6.2) ^E
	4-8	127	28.1 (2.0)	20.2 (3.2)	21.9 (2.9)	25.0 (2.6)	28.8 (2.6)	33.0 (3.2)	37.4 (4.4)	40.2 (5.4)
Male	9-13	111	32.3 (2.6)	20.8 (3.4)	22.9 (3.1)	26.7 (2.7)	31.6 (2.8)	37.6 (4.0)	44.3 (6.2)	49.0 (8.1)
	14-18	107	29.5 (1.9)	20.1 (3.6) ^E	21.9 (3.2)	25.2 (2.7)	29.4 (2.5)	34.1 (3.0)	38.9 (4.2)	42.0 (5.2)
	19-30	77	31.4 (2.9)	18.7 (3.3) ^E	21.1 (3.4)	25.6 (3.7)	31.5 (3.9)	37.9 (4.1)	43.6 (4.3)	47.0 (4.5)
	31-50	145	28.5 (3.9)	18.2 (3.5) ^E	19.9 (3.2)	23.0 (2.9)	27.0 (3.0)	31.5 (3.8)	36.1 (5.2)	39.2 (6.3)
	51-70	182	21.5 (1.2)	11.5 (0.9)	13.0 (1.0)	15.9 (1.1)	19.8 (1.3)	24.2 (1.6)	28.9 (1.8)	31.9 (2.1)
	>70	63	21.4 (2.1)	15.4 (3.1) ^E	16.8 (3.0) ^E	19.3 (2.9)	22.3 (3.0)	25.8 (3.5)	29.2 (4.2)	31.3 (4.8)
	19+	467	26.4 (1.7)	17.0 (1.8)	18.7 (1.7)	21.7 (1.5)	25.7 (1.6)	30.5 (2.0)	35.0 (2.7)	38.1 (3.3)
Female	9-13	96	23.7 (1.3)	12.4 (1.9)	14.4 (1.7)	17.8 (1.5)	22.0 (1.5)	27.4 (2.0)	34.0 (3.4)	39.0 (4.9)
	14-18	105	23.9 (1.7)	14.3 (3.1) ^E	15.8 (2.9) ^E	18.5 (2.6)	22.3 (2.3)	27.0 (2.7)	32.0 (4.0)	35.2 (5.0)
	19-30	91	22.9 (2.8)	13.1 (1.8)	14.7 (2.0)	17.5 (2.3)	21.0 (2.8)	25.4 (3.5)	30.4 (4.6)	34.1 (5.5)
	31-50	167	20.7 (1.9)	10.6 (2.3) ^E	12.1 (2.2) ^E	14.9 (2.2)	18.8 (2.3)	23.9 (2.6)	29.2 (3.2)	32.4 (3.7)
	51-70	198	16.6 (2.1)	8.1 (2.4) ^E	9.5 (2.1) ^E	12.2 (1.8)	15.7 (1.8)	20.2 (2.8)	25.2 (4.8) ^E	28.9 (6.5) ^E
	>70	74	19.2 (3.7) ^E	11.5 (3.1) ^E	12.8 (3.1) ^E	15.3 (3.1) ^E	18.7 (3.4) ^E	22.9 (4.9) ^E	27.7 (6.4) ^E	31.0 (7.7) ^E
	19+	530	19.7 (1.2)	11.1 (1.3)	12.4 (1.2)	15.0 (1.2)	18.5 (1.4)	22.8 (1.8)	27.6 (2.5)	31.0 (3.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.2 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)	
Both											
	1-3	58	21.0 (1.9)	12.1 (2.4) ^E	13.8 (2.3)	16.9 (2.2)	20.8 (2.4)	24.9 (3.0)	28.8 (3.6)	31.0 (4.1)	
	4-8	110	25.4 (2.0)	18.4 (2.1)	19.9 (2.2)	22.4 (2.4)	25.5 (2.6)	28.7 (2.8)	31.7 (2.9)	33.6 (3.0)	
Male	9-13	95	29.2 (2.2)	19.6 (3.9) ^E	21.7 (3.7) ^E	25.4 (3.6)	30.3 (3.6)	35.9 (4.1)	41.7 (5.1)	45.6 (5.9)	
	14-18	87	36.2 (2.6)	20.0 (4.0) ^E	23.5 (3.8)	30.3 (3.5)	38.6 (3.4)	47.4 (4.2)	56.0 (6.1)	61.9 (7.8)	
	19-30	70	45.6 (7.4)	25.5 (6.7) ^E	29.9 (6.5) ^E	39.3 (6.8) ^E	50.2 (8.6) ^E	63.5 (12.8) ^E	78.8 (18.2) ^E	89.6 (22.1) ^E	
	31-50	109	28.8 (1.9)	17.7 (3.7) ^E	20.0 (3.4) ^E	24.2 (2.8)	29.3 (2.4)	35.1 (2.8)	41.1 (4.0)	45.2 (5.1)	
	51-70	128	23.9 (1.4)	14.3 (3.3) ^E	16.2 (3.1) ^E	19.8 (2.5)	24.2 (2.1)	28.9 (2.2)	33.6 (3.0)	36.6 (3.9)	
	>70	65	21.3 (1.6)	11.1 (2.5) ^E	12.7 (2.4) ^E	15.7 (2.2)	20.1 (2.1)	25.8 (2.9)	32.4 (4.7)	37.1 (6.3) ^E	
	19+	372	30.1 (1.9)	15.5 (2.0)	18.0 (2.0)	23.0 (2.0)	30.3 (2.3)	39.3 (3.3)	49.4 (5.1)	56.9 (6.9)	
	Female	9-13	75	23.1 (1.5)	15.2 (1.6)	16.7 (1.8)	19.7 (2.0)	23.6 (2.3)	28.1 (2.7)	32.6 (3.2)	35.5 (3.6)
		14-18	81	23.9 (2.0)	11.5 (2.7) ^E	13.9 (2.6) ^E	18.5 (2.5)	24.1 (2.7)	30.5 (3.3)	36.7 (4.2)	40.6 (5.0)
19-30		101	21.7 (1.7)	12.8 (2.0)	14.8 (2.0)	18.7 (2.0)	23.6 (2.4)	29.2 (3.2)	34.7 (4.2)	38.3 (5.1)	
31-50		116	20.5 (1.6)	13.6 (2.8) ^E	14.9 (2.6) ^E	17.4 (2.3)	20.4 (2.1)	23.8 (2.5)	27.2 (3.4)	29.4 (4.2)	
51-70		146	20.5 (1.2)	10.6 (1.7)	12.3 (1.6)	15.7 (1.5)	20.2 (1.5)	25.4 (1.9)	30.4 (2.6)	33.5 (3.1)	
>70		94	17.0 (1.3)	10.2 (1.2)	11.3 (1.1)	13.4 (1.1)	16.5 (1.3)	20.3 (1.9)	24.5 (2.8)	27.4 (3.5)	
19+		457	20.3 (0.8)	11.5 (1.0)	13.1 (1.0)	16.1 (1.0)	20.5 (1.2)	25.6 (1.6)	30.6 (2.0)	33.6 (2.3)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.3 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	112	19.7	(1.2)	13.4	(2.7) ^E	14.7	(2.4)	17.0	(1.9)	19.8	(1.6)	22.9	(1.9)	26.0	(3.1)	27.9	(4.1)
	4-8	177	23.7	(1.3)	16.3	(1.3)	17.8	(1.3)	20.5	(1.4)	23.7	(1.6)	27.1	(1.7)	30.5	(1.9)	32.6	(2.0)
Male																		
	9-13	111	29.9	(2.0)	19.9	(3.2)	21.7	(2.9)	24.9	(2.5)	29.0	(2.3)	33.5	(2.6)	37.9	(3.6)	40.8	(4.4)
	14-18	113	32.7	(2.9)	17.8	(4.7) ^E	20.0	(4.8) ^E	24.5	(4.8) ^E	31.2	(4.7)	39.4	(4.9)	47.7	(6.1)	53.2	(7.3)
	19-30	91	33.5	(3.3)	19.8	(5.1) ^E	22.3	(4.6) ^E	27.0	(3.9)	33.1	(3.6)	40.2	(5.2)	47.7	(8.2) ^E	52.7	(10.5) ^E
	31-50	101	33.3	(2.2)	19.3	(4.4) ^E	21.9	(3.8) ^E	26.3	(3.0)	31.5	(2.5)	37.8	(3.2)	44.5	(5.0)	48.9	(6.5)
	51-70	134	25.3	(2.3)	11.7	(3.0) ^E	13.6	(2.8) ^E	17.7	(2.5)	23.4	(2.6)	30.9	(3.8)	39.5	(6.1)	45.6	(8.1) ^E
	>70	56	21.3	(2.0)	11.1	(2.4) ^E	13.0	(2.5) ^E	16.8	(2.4)	21.4	(2.4)	25.6	(2.9)	29.4	(3.4)	32.0	(3.9)
	19+	382	29.8	(1.4)	15.4	(2.1)	18.0	(1.9)	22.7	(1.6)	28.5	(1.5)	35.1	(2.1)	42.3	(3.4)	47.4	(4.5)
Female																		
	9-13	105	24.4	(1.9)	15.2	(1.7)	16.8	(1.7)	19.8	(1.9)	23.7	(2.2)	28.3	(2.7)	33.2	(3.3)	36.5	(3.9)
	14-18	120	22.0	(2.3)	10.1	(2.7) ^E	12.2	(2.7) ^E	16.1	(2.6)	21.4	(2.7)	27.4	(3.2)	33.5	(4.2)	37.4	(5.2)
	19-30	91	24.3	(2.4)	11.7	(2.5) ^E	14.2	(2.2)	18.5	(2.0)	23.5	(2.3)	29.8	(3.4)	37.9	(5.1)	44.4	(6.7)
	31-50	159	25.9	(2.3)	11.3	(2.6) ^E	14.0	(2.5) ^E	19.0	(2.4)	25.5	(2.7)	32.9	(3.4)	40.2	(4.6)	44.9	(5.5)
	51-70	174	20.1	(1.7)	10.6	(1.1)	12.2	(1.2)	15.3	(1.4)	18.8	(1.5)	23.0	(1.8)	27.8	(2.2)	31.2	(2.5)
	>70	80	22.5	(4.9) ^E	F		F		18.8	(5.5) ^E	23.0	(5.5) ^E	27.8	(5.6) ^E	32.3	(5.8) ^E	35.0	(6.0) ^E
	19+	504	23.5	(1.4)	10.2	(1.2)	12.4	(1.3)	16.8	(1.3)	22.7	(1.6)	30.4	(2.2)	38.9	(3.0)	44.7	(3.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.4 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake														
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)								
Both																			
	1-3	99	22.9	(1.5)	13.4	(2.3) ^E	15.0	(2.1)	18.0	(1.7)	22.0	(1.8)	27.0	(2.6)	32.3	(3.8)	35.8	(4.7)	
	4-8	140	26.1	(1.6)	17.6	(2.7)	19.3	(2.4)	22.5	(2.0)	26.4	(1.8)	30.7	(2.2)	34.9	(3.1)	37.8	(3.9)	
Male																			
	9-13	92	31.7	(3.1)	16.3	(5.0) ^E	19.3	(4.5) ^E	24.6	(3.9)	30.9	(3.8)	37.5	(4.1)	43.9	(4.8)	47.9	(5.4)	
	14-18	107	39.5	(3.9)	25.1	(5.2) ^E	28.2	(5.1) ^E	33.8	(4.9)	40.9	(4.8)	49.2	(5.4)	58.3	(7.2)	64.4	(9.0)	
	19-30	73	35.8	(3.5)	19.5	(4.1) ^E	22.5	(3.8) ^E	28.0	(3.6)	35.4	(4.2)	45.6	(6.6)	59.1	(11.4) ^E	70.0	(16.1) ^E	
	31-50	134	31.5	(2.2)	16.7	(3.6) ^E	19.4	(3.3) ^E	24.5	(2.7)	30.6	(2.5)	37.7	(3.4)	45.7	(5.3)	51.5	(6.9)	
	51-70	131	25.9	(1.6)	17.1	(2.4)	18.6	(2.2)	21.5	(1.8)	25.1	(1.7)	29.1	(2.4)	33.1	(3.4)	35.7	(4.2)	
	>70	55	23.0	(2.2)	11.8	(2.4) ^E	13.3	(2.4) ^E	16.4	(2.3)	21.0	(2.5)	27.4	(3.4)	35.1	(5.2)	40.8	(7.0) ^E	
	19+	393	30.0	(1.3)	16.2	(1.3)	18.6	(1.3)	23.3	(1.4)	29.5	(1.6)	37.0	(2.1)	45.2	(3.0)	51.2	(3.9)	
Female																			
	9-13	79	23.7	(2.8)	F		15.2	(4.1) ^E	19.0	(3.6) ^E	23.5	(3.3)	28.7	(4.0)	34.3	(5.3)	38.2	(6.4) ^E	
	14-18	104	23.1	(1.6)		14.2	(1.6)	16.1	(1.7)	19.7	(1.9)	24.1	(2.3)	28.8	(2.6)	33.4	(3.0)	36.3	(3.4)
	19-30	101	18.8	(1.7)	F		10.8	(3.0) ^E	14.4	(2.6) ^E	19.1	(2.2)	24.9	(2.4)	31.2	(3.3)	35.5	(4.3)	
	31-50	143	19.4	(1.3)		10.6	(2.4) ^E	12.1	(2.1) ^E	14.9	(1.8)	18.4	(1.6)	22.6	(2.2)	26.9	(3.3)	29.8	(4.2)
	51-70	193	17.5	(0.9)		11.0	(2.3) ^E	12.2	(2.0) ^E	14.4	(1.7)	17.0	(1.3)	19.7	(1.4)	22.3	(1.9)	23.9	(2.4)
	>70	94	17.2	(1.4)		10.0	(1.9) ^E	10.9	(1.8)	12.7	(1.6)	15.0	(1.6)	17.9	(2.2)	21.0	(3.4)	23.1	(4.4) ^E
	19+	531	18.4	(0.7)		9.7	(1.1)	11.2	(1.0)	14.0	(1.0)	17.7	(1.0)	22.2	(1.2)	26.9	(1.6)	30.1	(2.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.5 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	311	23.9	(1.0)	11.2	(1.3)	13.7	(1.3)	18.3	(1.3)	24.1	(1.4)	30.6	(1.6)	37.3	(2.2)	41.8	(2.6)
	4-8	485	27.8	(1.3)	15.6	(1.6)	17.7	(1.5)	21.5	(1.4)	26.8	(1.4)	33.1	(2.3)	39.2	(3.3)	43.4	(4.0)
Male																		
	9-13	277	35.4	(1.8)	21.1	(2.5)	23.9	(2.4)	29.3	(2.3)	36.2	(2.6)	43.9	(3.4)	51.2	(4.3)	55.7	(4.9)
	14-18	339	41.8	(2.2)	25.8	(1.5)	28.7	(1.6)	34.2	(1.9)	42.0	(2.7)	52.4	(3.8)	64.5	(5.0)	73.0	(5.8)
	19-30	237	33.3	(1.8)	22.2	(3.5)	24.2	(3.2)	27.9	(2.7)	32.3	(2.5)	37.4	(3.1)	42.6	(4.3)	46.1	(5.2)
	31-50	423	35.2	(2.1)	18.8	(3.0)	21.5	(2.8)	26.5	(2.5)	33.3	(2.5)	41.8	(3.4)	50.9	(5.0)	57.1	(6.3)
	51-70	387	30.1	(1.9)	13.4	(2.5) ^E	15.8	(2.5)	20.8	(2.3)	27.8	(2.2)	36.2	(2.8)	45.9	(4.3)	52.8	(5.7)
	>70	132	25.7	(2.3)	9.5	(2.6) ^E	11.9	(2.7) ^E	16.8	(2.8)	23.7	(3.1)	32.5	(4.0)	42.0	(5.5)	48.6	(6.7)
	19+	1179	32.5	(1.1)	16.0	(1.3)	18.7	(1.3)	23.9	(1.2)	30.7	(1.3)	39.2	(1.7)	48.8	(2.3)	55.3	(2.9)
Female																		
	9-13	281	28.9	(1.8)	16.7	(2.4)	18.9	(2.3)	22.9	(2.1)	28.0	(2.0)	33.7	(2.4)	39.3	(3.1)	43.0	(3.6)
	14-18	321	26.7	(1.5)	17.0	(2.4)	18.9	(2.3)	22.4	(2.1)	27.1	(2.0)	32.6	(2.5)	38.2	(3.4)	41.9	(4.1)
	19-30	249	26.2	(1.9)	13.7	(2.2)	15.8	(2.1)	19.9	(1.9)	25.2	(2.1)	31.5	(3.0)	38.2	(4.5)	42.5	(5.6)
	31-50	364	25.9	(1.7)	17.3	(3.4) ^E	19.0	(3.2) ^E	22.3	(2.8)	26.6	(2.4)	31.5	(2.5)	36.6	(3.3)	39.9	(4.3)
	51-70	467	22.0	(0.9)	12.1	(1.9)	14.1	(1.7)	17.7	(1.3)	22.1	(1.1)	27.0	(1.4)	31.9	(2.1)	35.2	(2.8)
	>70	215	18.3	(1.0)	11.9	(1.7)	13.1	(1.6)	15.3	(1.5)	18.2	(1.5)	21.6	(1.7)	25.2	(2.2)	27.5	(2.7)
	19+	1295	23.9	(0.8)	13.3	(1.1)	15.3	(1.1)	18.9	(1.1)	23.6	(1.1)	29.3	(1.3)	35.4	(1.7)	39.6	(2.1)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.6 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	644	19.3	(0.6)	7.1	(1.0)	9.4	(0.9)	13.7	(0.7)	18.7	(0.6)	24.5	(0.8)	31.1	(1.3)	35.8	(1.7)
	4-8	956	22.9	(0.5)	15.8	(1.9)	17.2	(1.6)	19.8	(1.1)	22.9	(0.6)	26.4	(1.0)	29.8	(1.8)	32.0	(2.4)
Male																		
	9-13	589	28.5	(0.8)	15.9	(1.6)	18.1	(1.5)	22.3	(1.2)	27.7	(1.0)	34.3	(1.3)	41.2	(2.2)	45.8	(2.9)
	14-18	639	32.2	(1.0)	17.6	(2.0)	20.3	(1.8)	25.5	(1.5)	32.3	(1.2)	40.2	(1.6)	48.5	(2.6)	54.0	(3.4)
	19-30	481	28.7	(1.4)	15.7	(1.2)	17.9	(1.2)	22.1	(1.4)	27.5	(1.6)	33.9	(2.0)	40.5	(2.4)	44.9	(2.8)
	31-50	709	26.3	(1.0)	13.6	(2.3) ^E	15.7	(2.2)	20.0	(1.7)	25.9	(1.3)	33.2	(1.8)	41.3	(3.4)	46.9	(4.7)
	51-70	758	23.4	(1.0)	10.4	(1.3)	12.4	(1.3)	16.3	(1.1)	21.6	(1.0)	28.3	(1.3)	36.1	(2.1)	41.6	(2.9)
	>70	734	19.8	(0.7)	9.4	(0.5)	11.1	(0.5)	14.3	(0.6)	18.6	(0.7)	23.7	(0.9)	29.5	(1.2)	33.5	(1.6)
	19+	2682	25.5	(0.6)	11.0	(0.8)	13.3	(0.8)	17.8	(0.7)	24.0	(0.7)	31.9	(0.9)	40.9	(1.4)	47.3	(1.9)
Female																		
	9-13	585	23.5	(0.7)	13.2	(1.6)	15.0	(1.4)	18.3	(1.1)	22.6	(0.8)	27.7	(1.1)	33.2	(2.0)	37.0	(2.8)
	14-18	645	24.5	(0.9)	17.7	(2.6)	19.2	(2.2)	21.8	(1.6)	25.0	(1.1)	28.6	(1.6)	32.2	(2.8)	34.6	(3.8)
	19-30	514	19.3	(0.9)	9.9	(1.7) ^E	11.4	(1.6)	14.4	(1.3)	18.2	(1.0)	22.7	(1.3)	27.1	(2.1)	29.9	(2.7)
	31-50	758	21.0	(0.9)	11.2	(1.4)	12.8	(1.3)	16.0	(1.1)	20.2	(0.9)	25.2	(1.2)	30.7	(2.0)	34.4	(2.6)
	51-70	955	18.7	(0.6)	8.8	(0.9)	10.5	(0.9)	13.5	(0.8)	17.6	(0.7)	22.8	(0.9)	28.6	(1.4)	32.7	(1.9)
	>70	1345	17.4	(0.5)	8.4	(0.8)	9.8	(0.7)	12.7	(0.6)	16.7	(0.5)	21.6	(0.8)	26.8	(1.3)	30.4	(1.7)
	19+	3572	19.6	(0.4)	9.7	(0.6)	11.3	(0.5)	14.5	(0.5)	18.7	(0.5)	23.9	(0.6)	29.4	(0.9)	33.2	(1.2)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.7 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake						
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	18.7	(0.8)	10.0 (1.6)	11.5 (1.4)	14.6 (1.1)	18.5 (1.0)	22.8 (1.4)	27.1 (2.1)	29.9 (2.7)
	4-8	425	22.5	(1.1)	12.5 (2.5) ^E	14.3 (2.2)	17.6 (1.6)	21.7 (1.2)	26.6 (1.7)	31.8 (3.1)	35.4 (4.1)
Male	9-13	274	28.1	(1.2)	19.4 (2.5)	21.2 (2.1)	24.4 (1.6)	28.3 (1.4)	32.7 (2.1)	37.0 (3.4)	39.7 (4.4)
	14-18	297	40.0	(2.8)	17.0 (2.7)	20.8 (2.5)	27.8 (2.5)	37.5 (2.9)	50.4 (4.1)	66.5 (6.3)	79.5 (8.9)
	19-30	249	32.3	(2.1)	17.6 (1.5)	20.0 (1.7)	24.7 (2.0)	31.2 (2.4)	39.2 (3.0)	47.7 (4.0)	53.4 (4.8)
	31-50	309	26.8	(1.5)	14.2 (2.9) ^E	16.2 (2.6)	20.2 (2.1)	25.6 (1.7)	32.2 (2.7)	39.3 (4.6)	44.1 (6.1)
	51-70	277	24.3	(1.3)	10.5 (2.7) ^E	12.4 (2.7) ^E	17.1 (2.3)	23.9 (1.9)	30.8 (2.1)	37.5 (3.4)	42.5 (4.7)
	>70	136	18.3	(1.1)	7.9 (1.9) ^E	9.4 (1.8) ^E	12.5 (1.5)	16.7 (1.3)	22.0 (1.7)	27.8 (3.0)	31.8 (4.2)
	19+	971	26.5	(0.9)	12.9 (1.7)	15.0 (1.6)	19.2 (1.4)	25.2 (1.1)	32.8 (1.5)	40.6 (2.7)	46.0 (3.7)
	Female	9-13	265	22.5	(0.9)	14.0 (1.3)	15.5 (1.3)	18.2 (1.2)	21.4 (1.2)	24.8 (1.2)	28.2 (1.4)
14-18	290	24.7	(1.5)	17.6 (3.1) ^E	19.1 (2.7)	21.9 (2.2)	25.5 (1.7)	29.6 (2.1)	33.8 (3.4)	36.6 (4.4)	
19-30	197	23.6	(2.1)	13.5 (3.1) ^E	15.2 (2.9) ^E	18.4 (2.4)	22.6 (2.3)	27.6 (2.9)	32.8 (4.3)	36.4 (5.4)	
31-50	312	21.8	(1.3)	16.6 (2.7)	17.6 (2.4)	19.4 (1.9)	21.6 (1.5)	23.9 (1.7)	26.1 (2.6)	27.5 (3.4)	
51-70	312	19.8	(1.4)	12.6 (2.3) ^E	13.9 (2.1)	16.4 (1.7)	19.7 (1.5)	23.5 (2.1)	27.4 (3.1)	30.0 (3.9)	
>70	239	15.1	(0.8)	8.3 (1.6) ^E	9.5 (1.4)	11.7 (1.1)	14.7 (0.9)	18.3 (1.2)	22.1 (2.0)	24.7 (2.7)	
19+	1060	20.6	(0.7)	12.0 (1.1)	13.4 (1.0)	16.3 (0.9)	20.1 (0.8)	24.8 (1.1)	29.6 (1.8)	32.8 (2.3)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.8 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	21.4 (1.3)	10.4 (1.8) ^E	12.6 (1.7)	16.4 (1.6)	21.2 (1.6)	26.9 (2.0)	33.3 (2.9)	37.9 (3.7)
	4-8	213	24.3 (1.1)	16.2 (2.4)	17.7 (2.1)	20.6 (1.6)	24.3 (1.2)	28.4 (1.6)	32.5 (2.6)	35.3 (3.4)
Male	9-13	122	29.5 (2.1)	21.2 (3.0)	22.8 (2.8)	25.7 (2.5)	29.5 (2.5)	34.0 (3.3)	38.4 (4.4)	41.2 (5.3)
	14-18	150	35.0 (2.1)	25.5 (4.0)	27.5 (3.6)	31.0 (3.0)	35.1 (2.6)	39.7 (3.3)	44.2 (4.8)	47.1 (6.1)
	19-30	106	33.3 (2.3)	18.8 (2.4)	21.5 (2.4)	26.5 (2.7)	33.0 (3.0)	40.4 (3.6)	47.8 (4.4)	52.6 (5.1)
	31-50	155	31.5 (2.6)	18.0 (3.6) ^E	20.3 (3.3)	24.8 (2.9)	31.0 (2.9)	38.5 (4.0)	46.5 (6.4)	52.2 (8.7) ^E
	51-70	122	27.3 (2.0)	15.2 (3.0) ^E	17.7 (2.7)	22.2 (2.4)	28.0 (2.6)	35.0 (3.7)	42.0 (5.5)	46.4 (6.8)
	>70	88	24.3 (1.9)	12.5 (2.4) ^E	14.5 (2.3)	18.4 (2.1)	23.6 (2.2)	29.6 (2.8)	36.0 (4.0)	40.2 (4.9)
	19+	471	30.0 (1.3)	14.3 (1.5)	17.0 (1.4)	22.1 (1.4)	29.3 (1.6)	38.1 (2.0)	47.3 (3.1)	53.8 (4.3)
Female	9-13	103	26.6 (2.1)	19.3 (3.3) ^E	20.8 (3.0)	23.7 (2.8)	27.2 (2.8)	31.2 (3.3)	35.2 (4.2)	37.9 (5.0)
	14-18	142	22.6 (1.5)	12.0 (1.4)	14.1 (1.4)	17.7 (1.5)	22.1 (1.9)	27.4 (2.7)	33.3 (3.7)	37.6 (4.6)
	19-30	111	21.5 (1.6)	12.4 (1.2)	14.0 (1.3)	17.0 (1.6)	20.5 (2.0)	24.5 (2.4)	28.5 (2.9)	31.1 (3.1)
	31-50	146	21.7 (1.5)	12.6 (2.4) ^E	14.0 (2.2)	16.5 (1.9)	19.7 (1.8)	23.7 (2.0)	28.2 (2.9)	31.3 (3.8)
	51-70	184	21.0 (1.3)	13.3 (2.6) ^E	14.7 (2.3)	17.4 (1.9)	20.8 (1.6)	24.7 (2.3)	28.7 (3.6)	31.4 (4.7)
	>70	143	16.4 (1.4)	7.2 (2.2) ^E	8.6 (2.1) ^E	11.5 (2.0) ^E	15.6 (1.8)	20.8 (2.1)	26.8 (3.0)	31.0 (3.8)
	19+	584	20.6 (0.8)	10.8 (1.0)	12.5 (0.9)	15.7 (0.8)	19.6 (0.9)	24.3 (1.2)	29.5 (1.7)	33.0 (2.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.9 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	169	17.9	(1.0)	8.0	(2.3) ^E	10.0	(2.0) ^E	13.6	(1.5)	18.1	(1.1)	23.1	(1.6)	28.2	(2.6)	31.7	(3.5)
	4-8	281	23.0	(0.9)	14.9	(0.7)	16.3	(0.8)	19.0	(0.9)	22.4	(1.1)	26.4	(1.3)	30.7	(1.6)	33.5	(1.8)
Male																		
	9-13	183	29.2	(1.7)	19.2	(1.2)	21.0	(1.3)	24.3	(1.5)	28.8	(1.8)	34.2	(2.3)	40.1	(2.9)	44.1	(3.4)
	14-18	187	35.1	(2.2)	20.6	(1.9)	23.8	(2.0)	29.0	(2.2)	34.5	(2.5)	40.6	(3.3)	48.0	(3.8)	53.4	(4.1)
	19-30	223	29.2	(1.6)	15.3	(3.3) ^E	17.5	(3.0) ^E	21.9	(2.3)	27.6	(1.8)	34.5	(2.5)	41.9	(4.2)	47.0	(5.7)
	31-50	229	27.6	(1.5)	12.9	(3.3) ^E	15.5	(2.9) ^E	20.3	(2.1)	26.4	(1.7)	33.4	(2.8)	40.5	(4.7)	45.2	(6.1)
	51-70	197	23.6	(1.7)	9.3	(1.8) ^E	11.6	(1.9)	16.2	(2.0)	22.6	(2.0)	30.2	(2.2)	38.4	(2.9)	43.8	(3.7)
	>70	72	21.3	(2.0)	10.6	(2.0) ^E	12.4	(2.2) ^E	16.0	(2.5)	20.7	(2.5)	25.6	(2.4)	30.1	(2.5)	33.0	(2.7)
	19+	721	26.6	(0.9)	12.6	(1.5)	15.1	(1.4)	19.8	(1.2)	25.7	(1.0)	32.5	(1.3)	39.6	(1.8)	44.3	(2.4)
Female																		
	9-13	165	23.5	(1.3)	13.4	(3.0) ^E	15.3	(2.7) ^E	18.9	(2.3)	23.2	(2.1)	28.2	(2.5)	33.9	(3.8)	38.0	(5.1)
	14-18	206	22.5	(1.5)	11.8	(2.0) ^E	13.4	(1.8)	16.6	(1.6)	20.8	(1.7)	26.3	(2.3)	32.7	(3.8)	37.4	(5.1)
	19-30	191	24.8	(2.6)	15.7	(4.3) ^E	17.5	(4.2) ^E	20.7	(3.9) ^E	24.9	(3.7)	29.8	(4.0)	34.7	(5.1)	38.0	(6.3)
	31-50	258	21.6	(1.4)	15.1	(3.2) ^E	16.4	(3.0) ^E	18.8	(2.6)	21.9	(2.2)	25.4	(2.1)	29.0	(2.7)	31.4	(3.5)
	51-70	249	20.7	(1.5)	9.8	(2.0) ^E	11.5	(1.9)	14.7	(1.7)	19.4	(1.6)	25.4	(2.2)	32.3	(3.8)	37.5	(5.3)
	>70	128	15.8	(1.2)	7.6	(1.1)	9.1	(1.0)	11.8	(1.0)	14.9	(1.1)	18.6	(1.5)	23.0	(2.3)	26.6	(3.2)
	19+	826	21.5	(1.0)	11.3	(1.4)	13.0	(1.4)	16.4	(1.3)	21.0	(1.3)	26.5	(1.4)	32.5	(1.9)	36.8	(2.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.10 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	192	20.6	(1.3)	10.8	(1.9) ^E	12.6	(1.8)	15.9	(1.6)	20.0	(1.7)	24.6	(2.2)	29.2	(2.9)	32.1	(3.3)
	4-8	321	22.4	(0.8)	13.8	(1.6)	15.4	(1.4)	18.4	(1.1)	22.1	(1.0)	26.4	(1.4)	31.1	(2.3)	34.2	(3.0)
Male																		
	9-13	226	30.0	(1.4)	17.1	(2.3)	19.7	(2.1)	24.7	(1.8)	30.8	(1.7)	37.6	(2.1)	44.7	(3.1)	49.6	(4.0)
	14-18	262	34.8	(1.9)	22.9	(3.6)	25.2	(3.2)	29.6	(2.6)	35.1	(2.3)	41.3	(3.0)	47.8	(4.6)	52.2	(5.9)
	19-30	197	33.5	(1.8)	20.1	(2.6)	22.4	(2.4)	26.7	(2.0)	32.3	(2.1)	39.1	(3.1)	46.4	(4.9)	51.2	(6.2)
	31-50	282	33.9	(2.0)	17.8	(3.5) ^E	20.6	(3.1)	26.0	(2.4)	33.0	(2.2)	41.3	(3.4)	50.1	(5.5)	56.2	(7.0)
	51-70	234	29.1	(1.7)	14.9	(2.3)	17.2	(2.2)	21.8	(2.2)	27.9	(2.2)	35.4	(2.7)	43.4	(3.7)	48.8	(4.6)
	>70	119	21.9	(2.0)	9.5	(1.9) ^E	11.5	(1.9) ^E	15.3	(2.0)	20.5	(2.2)	26.8	(2.5)	33.4	(2.9)	37.9	(3.3)
	19+	832	31.3	(1.1)	15.9	(1.2)	18.6	(1.1)	23.6	(1.1)	30.3	(1.2)	38.4	(1.7)	47.3	(2.5)	53.4	(3.1)
Female																		
	9-13	226	25.0	(1.5)	14.9	(1.4)	16.7	(1.4)	19.9	(1.4)	24.1	(1.7)	29.5	(2.5)	35.5	(3.6)	39.8	(4.5)
	14-18	242	22.0	(1.0)	9.6	(1.2)	11.8	(1.2)	15.7	(1.1)	20.7	(1.2)	26.8	(1.6)	33.7	(2.2)	38.7	(2.7)
	19-30	208	22.4	(1.5)	12.6	(2.1)	14.4	(2.0)	17.8	(1.9)	22.2	(1.9)	27.1	(2.4)	32.1	(3.1)	35.4	(3.6)
	31-50	263	22.4	(1.4)	12.4	(1.0)	14.2	(1.1)	17.5	(1.4)	21.9	(1.7)	27.9	(2.1)	34.3	(2.6)	38.9	(3.2)
	51-70	322	18.1	(1.2)	9.4	(2.1) ^E	10.8	(2.0) ^E	13.5	(1.7)	17.4	(1.6)	22.3	(1.9)	27.3	(2.8)	30.7	(3.6)
	>70	198	17.9	(1.3)	7.4	(1.2)	8.9	(1.2)	11.7	(1.2)	15.8	(1.4)	21.4	(2.0)	27.8	(3.0)	32.1	(3.8)
	19+	991	20.6	(0.7)	9.7	(0.8)	11.4	(0.8)	14.9	(0.8)	19.9	(0.9)	26.1	(1.2)	32.6	(1.6)	36.9	(1.9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.11 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	348	21.7	(0.9)	11.6	(1.3)	13.3	(1.3)	16.7	(1.1)	21.1	(1.1)	26.3	(1.4)	32.1	(2.0)	36.0	(2.6)
	4-8	554	25.5	(0.9)	18.0	(1.6)	19.6	(1.5)	22.5	(1.2)	26.0	(1.1)	29.9	(1.3)	33.6	(1.8)	36.1	(2.2)
Male																		
	9-13	409	30.9	(1.5)	18.6	(2.0)	20.9	(1.8)	25.1	(1.7)	30.4	(1.7)	36.4	(2.1)	42.4	(2.8)	46.3	(3.4)
	14-18	414	34.3	(1.7)	18.7	(2.0)	21.4	(2.1)	26.9	(2.2)	34.1	(2.3)	42.6	(2.7)	51.8	(3.4)	58.1	(4.0)
	19-30	311	34.4	(1.8)	19.8	(2.4)	22.6	(2.2)	27.8	(2.1)	34.7	(2.3)	43.3	(3.2)	52.9	(4.9)	59.7	(6.5)
	31-50	489	31.4	(1.4)	17.5	(1.6)	19.9	(1.5)	24.3	(1.4)	30.0	(1.5)	36.6	(2.0)	43.7	(2.9)	48.6	(3.6)
	51-70	575	24.5	(1.1)	13.1	(1.2)	14.9	(1.2)	18.4	(1.2)	23.2	(1.2)	28.8	(1.6)	34.6	(2.3)	38.5	(2.9)
	>70	239	21.8	(1.2)	12.6	(1.8)	14.3	(1.8)	17.4	(1.6)	21.5	(1.5)	26.2	(1.9)	31.2	(2.7)	34.5	(3.2)
	19+	1614	29.1	(0.8)	15.2	(0.8)	17.6	(0.8)	22.1	(0.8)	28.0	(0.9)	35.2	(1.2)	43.0	(1.7)	48.6	(2.2)
Female																		
	9-13	355	23.9	(1.2)	14.2	(2.1)	16.0	(1.9)	19.3	(1.6)	23.3	(1.5)	28.0	(1.8)	33.1	(2.6)	36.5	(3.3)
	14-18	410	22.9	(1.1)	12.0	(1.4)	14.0	(1.3)	17.7	(1.3)	22.4	(1.4)	27.9	(1.5)	33.4	(1.7)	37.1	(1.9)
	19-30	384	22.1	(1.3)	13.0	(1.5)	14.8	(1.4)	18.1	(1.3)	22.3	(1.4)	27.3	(1.8)	32.7	(2.5)	36.5	(3.2)
	31-50	585	22.4	(1.1)	10.9	(1.2)	12.9	(1.2)	16.6	(1.3)	21.4	(1.4)	27.3	(1.7)	33.7	(2.2)	37.9	(2.6)
	51-70	711	18.5	(0.9)	9.5	(1.1)	11.0	(1.1)	14.0	(1.0)	17.6	(1.0)	21.9	(1.2)	26.6	(1.7)	30.0	(2.3)
	>70	342	19.8	(2.3)	9.4	(1.6) ^E	10.9	(1.7)	14.0	(1.9)	18.6	(2.4)	24.5	(3.1)	31.2	(3.9)	35.6	(4.4)
	19+	2022	20.9	(0.6)	10.7	(0.6)	12.4	(0.6)	15.8	(0.7)	20.3	(0.8)	25.7	(1.0)	31.7	(1.3)	35.9	(1.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.12 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	622	18.7	(0.7)	9.1	(1.2)	10.9	(1.0)	14.3	(0.9)	18.7	(0.8)	23.8	(0.9)	28.5	(1.2)	31.9	(1.6)
	4-8	919	23.1	(0.6)	14.9	(1.6)	16.4	(1.4)	19.2	(1.0)	22.7	(0.7)	26.7	(1.1)	30.8	(2.0)	33.5	(2.6)
Male																		
	9-13	579	29.0	(1.1)	19.1	(1.8)	21.0	(1.7)	24.5	(1.4)	29.0	(1.2)	34.3	(1.7)	39.8	(2.7)	43.6	(3.4)
	14-18	634	36.1	(1.5)	21.3	(2.9)	24.3	(2.6)	29.3	(2.1)	35.6	(1.8)	43.3	(2.3)	51.9	(3.7)	57.7	(4.9)
	19-30	578	30.5	(1.2)	16.6	(2.5)	18.9	(2.3)	23.3	(1.9)	29.1	(1.5)	36.1	(1.9)	43.4	(3.2)	48.2	(4.2)
	31-50	693	28.1	(1.1)	13.4	(1.6)	15.7	(1.5)	20.4	(1.3)	26.8	(1.2)	34.3	(1.8)	42.0	(2.8)	47.3	(3.9)
	51-70	596	24.4	(1.1)	10.2	(1.9) ^E	12.8	(1.8)	17.6	(1.6)	23.8	(1.4)	31.1	(1.6)	39.2	(2.2)	44.7	(2.9)
	>70	296	21.3	(1.1)	10.2	(1.0)	12.0	(1.1)	15.5	(1.3)	20.1	(1.4)	25.6	(1.5)	31.1	(1.7)	34.7	(1.8)
	19+	2163	27.2	(0.6)	12.3	(0.8)	14.8	(0.8)	19.7	(0.7)	26.0	(0.7)	33.7	(0.9)	42.1	(1.3)	47.8	(1.6)
Female																		
	9-13	533	23.8	(0.9)	15.3	(1.8)	16.9	(1.7)	19.9	(1.4)	23.5	(1.3)	27.7	(1.6)	31.9	(2.2)	34.8	(2.7)
	14-18	638	23.0	(1.0)	12.1	(0.9)	13.8	(1.0)	17.1	(1.0)	21.9	(1.2)	27.7	(1.6)	34.2	(2.3)	38.8	(3.1)
	19-30	499	24.0	(1.6)	12.3	(1.3)	14.3	(1.4)	18.1	(1.6)	23.0	(1.9)	29.1	(2.4)	35.6	(3.0)	39.9	(3.4)
	31-50	716	21.6	(0.9)	14.8	(2.3)	16.1	(2.1)	18.5	(1.8)	21.6	(1.4)	25.1	(1.4)	28.8	(2.1)	31.2	(2.7)
	51-70	745	20.6	(0.9)	11.0	(1.2)	12.5	(1.1)	15.6	(1.1)	19.9	(1.0)	25.4	(1.4)	31.2	(2.3)	35.3	(3.1)
	>70	510	15.7	(0.7)	7.3	(0.7)	8.7	(0.7)	11.4	(0.7)	14.9	(0.8)	19.2	(1.0)	24.1	(1.4)	27.7	(1.8)
	19+	2470	21.1	(0.6)	11.2	(0.8)	12.9	(0.8)	16.2	(0.8)	20.5	(0.8)	25.8	(0.9)	31.5	(1.2)	35.5	(1.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.13 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	20.5 (0.4)	9.7 (0.6)	11.8 (0.5)	15.5 (0.5)	20.2 (0.5)	25.6 (0.6)	31.4 (0.8)	35.2 (1.0)
	4-8	3235	24.1 (0.4)	14.7 (0.6)	16.5 (0.5)	19.6 (0.4)	23.7 (0.5)	28.4 (0.6)	33.3 (0.9)	36.6 (1.2)
Male	9-13	2080	30.5 (0.6)	17.9 (0.7)	20.2 (0.7)	24.5 (0.7)	30.1 (0.7)	36.6 (0.9)	43.4 (1.2)	47.9 (1.5)
	14-18	2288	35.5 (0.8)	19.7 (1.0)	22.6 (1.0)	28.3 (0.9)	35.6 (0.9)	44.1 (1.2)	53.4 (1.7)	60.1 (2.2)
	19-30	1804	31.1 (0.8)	17.6 (1.2)	19.9 (1.1)	24.2 (1.0)	29.8 (1.0)	36.4 (1.3)	43.2 (1.9)	47.8 (2.3)
	31-50	2596	30.1 (0.7)	14.4 (0.8)	17.0 (0.8)	22.1 (0.8)	28.6 (0.8)	36.8 (1.1)	46.2 (1.7)	52.7 (2.2)
	51-70	2550	26.1 (0.7)	11.5 (0.7)	13.6 (0.7)	18.1 (0.7)	24.4 (0.8)	32.0 (1.0)	40.5 (1.4)	46.6 (1.8)
	>70	1520	21.9 (0.7)	9.9 (0.7)	11.8 (0.7)	15.4 (0.7)	20.4 (0.8)	26.4 (1.1)	33.1 (1.5)	37.8 (2.0)
	19+	8470	28.5 (0.4)	13.1 (0.4)	15.6 (0.4)	20.4 (0.4)	26.9 (0.5)	34.9 (0.6)	43.8 (0.9)	50.1 (1.1)
Female	9-13	1980	25.0 (0.6)	14.5 (0.7)	16.3 (0.6)	19.7 (0.6)	24.2 (0.7)	29.4 (0.8)	34.9 (1.1)	38.6 (1.4)
	14-18	2256	24.3 (0.6)	13.2 (0.7)	15.2 (0.6)	18.9 (0.6)	23.8 (0.7)	29.9 (0.9)	36.5 (1.2)	41.0 (1.5)
	19-30	1854	22.4 (0.7)	11.4 (0.8)	13.2 (0.8)	16.7 (0.7)	21.2 (0.8)	26.6 (1.0)	32.2 (1.4)	35.9 (1.7)
	31-50	2686	22.5 (0.6)	12.4 (0.9)	14.2 (0.9)	17.6 (0.8)	22.2 (0.7)	27.5 (0.8)	33.3 (1.1)	37.2 (1.4)
	51-70	3200	19.8 (0.4)	10.3 (0.5)	11.8 (0.5)	14.9 (0.5)	19.1 (0.5)	24.2 (0.6)	29.6 (0.9)	33.2 (1.1)
	>70	2610	17.6 (0.4)	8.6 (0.4)	10.0 (0.4)	12.9 (0.5)	16.7 (0.5)	21.6 (0.6)	27.0 (0.9)	30.7 (1.1)
	19+	10350	21.1 (0.3)	10.7 (0.3)	12.4 (0.3)	15.8 (0.3)	20.4 (0.4)	25.9 (0.5)	31.9 (0.6)	36.0 (0.8)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

40. Total sugars (g/d): Usual intakes from food

Table 40.1 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Newfoundland and Labrador, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	79	111 (9)	72 (14) ^E	80 (13)	95 (11)	111 (10)	129 (12)	146 (16)	157 (18)
	4-8	127	139 (7)	88 (13)	98 (11)	117 (8)	139 (7)	163 (10)	190 (16)	208 (22)
Male	9-13	111	173 (16)	87 (15) ^E	99 (15)	124 (15)	158 (17)	203 (22)	256 (31)	296 (39)
	14-18	107	178 (16)	99 (22) ^E	112 (21) ^E	137 (18)	172 (17)	216 (23)	266 (36)	300 (47)
	19-30	77	135 (12)	89 (17) ^E	96 (16) ^E	110 (15)	130 (15)	152 (17)	174 (23)	188 (28)
	31-50	145	124 (14)	65 (17) ^E	75 (16) ^E	94 (15)	118 (15)	147 (17)	177 (22)	199 (27)
	51-70	182	96 (8)	46 (10) ^E	54 (10) ^E	70 (9)	91 (9)	117 (13)	143 (19)	160 (25)
	>70	63	94 (10)	51 (12) ^E	57 (12) ^E	69 (12) ^E	89 (13)	115 (18)	143 (26) ^E	163 (33) ^E
	19+	467	115 (7)	58 (7)	67 (7)	86 (7)	112 (8)	142 (10)	174 (14)	194 (17)
Female	9-13	96	126 (10)	56 (16) ^E	69 (16) ^E	94 (14)	125 (13)	162 (16)	198 (22)	222 (27)
	14-18	105	120 (9)	66 (16) ^E	76 (15) ^E	93 (13)	117 (12)	147 (17)	181 (27)	205 (35) ^E
	19-30	91	108 (9)	62 (9)	71 (10)	87 (11)	108 (12)	130 (14)	151 (16)	165 (17)
	31-50	167	73 (6)	22 (6) ^E	31 (6) ^E	49 (6)	70 (7)	97 (9)	123 (14)	141 (17)
	51-70	198	73 (4)	36 (8) ^E	43 (8) ^E	56 (6)	71 (5)	88 (6)	104 (9)	114 (11)
	>70	74	95 (14)	48 (12) ^E	55 (11) ^E	68 (10)	85 (10)	105 (12)	125 (15)	138 (19)
	19+	530	82 (3)	33 (4)	41 (4)	58 (3)	78 (4)	102 (5)	126 (7)	144 (10)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.2 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Prince Edward Island, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	58	99 (10)	60 (12) ^E	68 (12) ^E	81 (11)	97 (12)	114 (13)	131 (16)	141 (18)
	4-8	110	123 (10)	76 (13) ^E	85 (12)	101 (11)	120 (12)	143 (16)	164 (21)	177 (24)
Male	9-13	95	137 (9)	89 (16) ^E	98 (15)	115 (13)	136 (12)	160 (14)	184 (18)	199 (21)
	14-18	87	158 (13)	102 (21) ^E	116 (20) ^E	140 (19)	171 (19)	207 (23)	243 (29)	267 (34)
	19-30	70	158 (21)	84 (27) ^E	97 (26) ^E	122 (25) ^E	155 (26) ^E	195 (35) ^E	238 (45) ^E	267 (53) ^E
	31-50	109	112 (9)	48 (10) ^E	60 (10) ^E	82 (10)	111 (12)	145 (16)	179 (20)	201 (23)
	51-70	128	104 (8)	61 (14) ^E	70 (13) ^E	86 (10)	107 (10)	131 (14)	156 (21)	173 (26)
	>70	65	91 (5)	51 (9) ^E	58 (8)	72 (7)	90 (7)	111 (10)	133 (14)	147 (19)
	19+	372	117 (6)	54 (8)	65 (8)	86 (7)	115 (8)	151 (11)	193 (17)	223 (22)
Female	9-13	75	126 (10)	84 (16) ^E	92 (14)	106 (13)	124 (13)	142 (15)	161 (20)	173 (23)
	14-18	81	129 (9)	87 (17) ^E	96 (15)	113 (12)	132 (10)	153 (12)	174 (17)	187 (22)
	19-30	101	113 (10)	60 (12) ^E	68 (13) ^E	86 (13)	112 (14)	144 (17)	177 (22)	200 (27)
	31-50	116	92 (5)	40 (6)	50 (6)	70 (6)	92 (6)	116 (8)	145 (12)	166 (16)
	51-70	146	87 (5)	47 (4)	54 (4)	67 (5)	85 (6)	106 (8)	131 (11)	149 (14)
	>70	94	72 (6)	33 (9) ^E	39 (9) ^E	50 (8)	65 (8)	82 (10)	100 (13)	112 (16)
	19+	457	92 (3)	44 (4)	52 (4)	68 (4)	89 (4)	114 (5)	142 (8)	162 (9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.3 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Nova Scotia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	112	93 (5)	59 (6)	64 (6)	75 (7)	90 (7)	106 (7)	120 (7)	128 (7)
	4-8	177	129 (7)	76 (11)	87 (10)	107 (8)	131 (8)	158 (10)	184 (13)	200 (15)
Male	9-13	111	157 (10)	107 (16)	116 (14)	133 (12)	154 (11)	177 (16)	200 (23)	214 (29)
	14-18	113	164 (14)	83 (20) ^E	96 (19) ^E	119 (19)	151 (20)	193 (24)	239 (32)	272 (40)
	19-30	91	131 (14)	80 (21) ^E	87 (19) ^E	101 (16)	118 (15)	136 (19)	154 (25)	166 (31) ^E
	31-50	101	129 (12)	56 (15) ^E	69 (14) ^E	95 (13)	128 (13)	169 (17)	218 (27)	255 (36)
	51-70	134	96 (7)	58 (12) ^E	65 (11)	77 (8)	91 (8)	109 (10)	128 (17)	141 (22)
	>70	56	96 (10)	^F	52 (17) ^E	69 (15) ^E	90 (13)	116 (15)	142 (22)	159 (28) ^E
	19+	382	116 (6)	50 (8)	61 (7)	82 (7)	109 (7)	142 (8)	177 (12)	201 (16)
Female	9-13	105	126 (8)	70 (15) ^E	81 (13)	100 (11)	123 (10)	146 (11)	169 (15)	184 (19)
	14-18	120	134 (13)	62 (13) ^E	72 (13) ^E	93 (14)	123 (16)	162 (19)	208 (26)	241 (34)
	19-30	91	128 (16)	76 (21) ^E	88 (21) ^E	110 (21) ^E	135 (21)	156 (22)	177 (27)	193 (31)
	31-50	159	100 (8)	51 (9) ^E	61 (9)	80 (9)	103 (10)	130 (13)	156 (17)	174 (20)
	51-70	174	81 (4)	36 (9) ^E	44 (8) ^E	60 (7)	80 (5)	102 (8)	124 (12)	138 (14)
	>70	80	89 (8)	42 (9) ^E	50 (9) ^E	68 (10)	91 (12)	115 (13)	139 (14)	155 (16)
	19+	504	98 (5)	43 (5)	53 (5)	72 (5)	98 (6)	128 (7)	159 (9)	179 (10)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.4 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, New Brunswick, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	99	109 (5)	74 (11)	82 (9)	94 (8)	110 (7)	130 (9)	149 (12)	162 (15)
	4-8	140	130 (8)	110 (18)	114 (15)	122 (12)	130 (9)	139 (17)	147 (24)	152 (29) ^E
Male	9-13	92	140 (14)	82 (18) ^E	91 (18) ^E	107 (17)	132 (18)	162 (22)	196 (30)	219 (36)
	14-18	107	175 (16)	100 (21) ^E	112 (19) ^E	136 (18)	171 (18)	215 (23)	265 (32)	298 (41)
	19-30	73	151 (23)	73 (17) ^E	86 (16) ^E	110 (16)	143 (20)	184 (30)	232 (45) ^E	265 (57) ^E
	31-50	134	124 (14)	51 (13) ^E	61 (13) ^E	81 (13)	110 (14)	144 (18)	181 (25)	207 (31)
	51-70	131	102 (7)	47 (6)	54 (6)	70 (7)	91 (8)	118 (11)	149 (15)	170 (18)
	>70	55	105 (12)	48 (16) ^E	57 (16) ^E	75 (16) ^E	100 (16)	128 (16)	154 (18)	172 (21)
	19+	393	121 (7)	51 (5)	62 (6)	82 (7)	113 (8)	152 (11)	197 (18)	231 (25)
	Female	9-13	79	123 (8)	76 (8)	85 (9)	102 (9)	124 (11)	147 (12)	169 (14)
14-18	104	113 (9)	62 (14) ^E	72 (12) ^E	90 (11)	112 (10)	137 (13)	162 (18)	178 (22)	
19-30	101	101 (9)	58 (15) ^E	66 (15) ^E	83 (13)	104 (12)	126 (13)	148 (18)	163 (24)	
31-50	143	90 (6)	37 (6) ^E	46 (6)	63 (6)	84 (7)	108 (9)	138 (11)	160 (13)	
51-70	193	79 (4)	48 (9) ^E	54 (8)	63 (6)	75 (5)	89 (6)	102 (10)	111 (12)	
>70	94	69 (4)	35 (7) ^E	39 (6)	47 (6)	59 (5)	73 (7)	90 (12)	102 (16)	
19+	531	86 (3)	39 (3)	47 (3)	62 (4)	81 (4)	103 (5)	127 (6)	145 (7)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.5 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Quebec, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	311	108 (5)	51 (6)	62 (6)	81 (6)	106 (6)	135 (8)	167 (11)	190 (13)
	4-8	485	125 (4)	79 (10)	88 (8)	105 (6)	125 (5)	147 (7)	168 (11)	181 (13)
Male	9-13	277	172 (8)	102 (14)	117 (13)	143 (13)	178 (14)	218 (17)	259 (22)	286 (25)
	14-18	339	194 (12)	106 (15)	124 (14)	156 (13)	196 (14)	243 (18)	295 (24)	330 (30)
	19-30	237	149 (8)	84 (14) ^E	95 (13)	117 (11)	145 (11)	176 (14)	207 (20)	228 (24)
	31-50	423	132 (8)	68 (12) ^E	80 (11)	102 (10)	131 (9)	165 (12)	200 (19)	224 (24)
	51-70	387	109 (4)	52 (4)	62 (4)	81 (5)	106 (5)	135 (6)	165 (7)	184 (8)
	>70	132	98 (7)	42 (9) ^E	52 (9) ^E	72 (9)	100 (10)	134 (13)	165 (16)	187 (20)
	19+	1179	126 (4)	56 (4)	67 (4)	91 (4)	122 (5)	160 (7)	202 (9)	230 (12)
	Female									
9-13	281	132 (5)	78 (6)	89 (6)	108 (6)	131 (7)	156 (7)	180 (9)	196 (10)	
14-18	321	127 (5)	80 (8)	90 (7)	108 (7)	130 (7)	154 (8)	178 (11)	194 (13)	
19-30	249	113 (7)	57 (5)	68 (6)	88 (7)	114 (9)	146 (11)	178 (13)	198 (15)	
31-50	364	99 (7)	47 (7)	55 (7)	72 (7)	94 (8)	123 (10)	158 (14)	185 (19)	
51-70	467	95 (4)	47 (4)	55 (4)	70 (4)	90 (4)	116 (6)	146 (10)	169 (14)	
>70	215	81 (4)	42 (7)	50 (6)	63 (6)	80 (6)	97 (7)	114 (9)	126 (11)	
19+	1295	98 (3)	48 (3)	56 (3)	72 (3)	95 (4)	123 (5)	155 (7)	178 (9)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.6 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Ontario, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	644	99 (3)	51 (6)	60 (5)	77 (4)	97 (3)	121 (4)	146 (7)	164 (10)
	4-8	956	122 (3)	76 (7)	85 (6)	101 (5)	121 (3)	143 (4)	166 (8)	181 (11)
	9-13	589	151 (5)	94 (12)	104 (10)	123 (8)	148 (6)	176 (8)	207 (15)	228 (20)
Male	14-18	639	159 (5)	77 (8)	92 (8)	121 (7)	159 (6)	204 (8)	250 (11)	279 (14)
	19-30	481	129 (5)	82 (17) ^E	91 (15) ^E	107 (11)	127 (7)	150 (8)	172 (15)	185 (20)
	31-50	709	101 (4)	63 (13) ^E	71 (11)	84 (8)	100 (5)	119 (7)	137 (13)	148 (18)
	51-70	758	98 (3)	46 (5)	55 (5)	72 (4)	93 (4)	118 (5)	145 (8)	162 (10)
	>70	734	93 (3)	37 (3)	46 (3)	63 (3)	87 (3)	116 (4)	149 (7)	171 (9)
	19+	2682	105 (2)	51 (4)	61 (4)	79 (3)	102 (3)	129 (3)	157 (5)	176 (7)
	9-13	585	126 (4)	92 (12)	98 (10)	110 (7)	124 (5)	139 (7)	154 (12)	163 (16)
	14-18	645	126 (4)	64 (9)	76 (8)	98 (7)	126 (5)	159 (8)	193 (12)	216 (16)
Female	19-30	514	99 (4)	56 (9)	63 (8)	77 (6)	94 (5)	113 (6)	132 (10)	144 (13)
	31-50	758	89 (3)	35 (6)	44 (5)	63 (4)	87 (4)	113 (4)	139 (7)	158 (9)
	51-70	955	82 (2)	39 (4)	46 (3)	60 (3)	78 (3)	100 (4)	124 (6)	141 (7)
	>70	1345	84 (2)	39 (3)	47 (3)	61 (3)	80 (2)	102 (3)	127 (5)	145 (7)
	19+	3572	89 (2)	39 (2)	48 (2)	64 (2)	84 (2)	109 (2)	135 (3)	153 (4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.7 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Manitoba, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	324	99 (5)	47 (11) ^E	56 (9)	73 (6)	94 (5)	121 (10)	152 (20)	175 (27)
	4-8	425	122 (5)	69 (12) ^E	79 (10)	96 (7)	117 (5)	143 (8)	171 (15)	190 (21)
Male	9-13	274	154 (6)	88 (18) ^E	100 (15)	122 (11)	150 (7)	183 (12)	217 (21)	240 (28)
	14-18	297	185 (10)	94 (14)	111 (13)	143 (12)	183 (13)	231 (18)	282 (25)	317 (31)
	19-30	249	135 (9)	65 (21) ^E	76 (19) ^E	97 (14)	124 (11)	157 (14)	191 (25)	213 (33)
	31-50	309	105 (6)	45 (5)	55 (5)	76 (6)	103 (7)	135 (9)	171 (11)	196 (13)
	51-70	277	97 (6)	41 (10) ^E	50 (9) ^E	69 (8)	94 (7)	124 (9)	156 (14)	178 (19)
	>70	136	92 (7)	38 (9) ^E	46 (9) ^E	62 (8)	86 (8)	117 (11)	153 (19)	179 (26)
	19+	971	108 (4)	40 (5)	51 (5)	72 (5)	101 (4)	138 (6)	179 (9)	208 (12)
	Female	9-13	265	144 (10)	80 (13) ^E	91 (12)	109 (9)	131 (8)	157 (11)	183 (16)
14-18	290	132 (12)	72 (9)	82 (10)	103 (11)	131 (14)	164 (18)	201 (23)	227 (27)	
19-30	197	127 (9)	65 (17) ^E	76 (15) ^E	99 (12)	128 (11)	158 (15)	182 (21)	198 (26)	
31-50	312	87 (6)	43 (6)	50 (6)	63 (6)	82 (6)	106 (9)	134 (13)	155 (17)	
51-70	312	88 (6)	38 (6)	46 (5)	61 (5)	82 (6)	106 (8)	132 (12)	150 (14)	
>70	239	80 (4)	38 (7) ^E	46 (6)	62 (5)	80 (4)	102 (6)	125 (8)	141 (11)	
19+	1060	94 (3)	41 (3)	50 (3)	66 (3)	88 (3)	117 (5)	150 (8)	173 (9)	

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.8 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	99 (5)	52 (8)	61 (7)	77 (7)	96 (6)	119 (7)	143 (10)	159 (12)
	4-8	213	119 (6)	75 (5)	83 (5)	97 (6)	116 (6)	137 (7)	158 (8)	173 (9)
Male	9-13	122	145 (11)	89 (14)	100 (13)	119 (13)	145 (15)	174 (18)	204 (24)	225 (29)
	14-18	150	182 (18)	133 (24) ^E	143 (22)	161 (21)	183 (22)	206 (27)	230 (35)	245 (42) ^E
	19-30	106	137 (14)	50 (14) ^E	63 (15) ^E	89 (15) ^E	125 (17)	169 (22)	217 (31)	250 (39)
	31-50	155	134 (15)	60 (16) ^E	72 (15) ^E	96 (14)	128 (15)	167 (20)	214 (32)	249 (45) ^E
	51-70	122	97 (7)	44 (13) ^E	54 (12) ^E	73 (10)	97 (9)	125 (12)	153 (18)	171 (23)
	>70	88	95 (6)	50 (6)	58 (6)	72 (6)	91 (7)	114 (9)	142 (13)	163 (17)
	19+	471	120 (7)	46 (6)	58 (5)	82 (6)	114 (7)	156 (10)	207 (16)	246 (22)
Female	9-13	103	131 (10)	75 (10)	83 (10)	99 (11)	125 (13)	158 (17)	196 (24)	223 (30)
	14-18	142	145 (12)	98 (17) ^E	109 (17)	129 (17)	153 (18)	181 (22)	209 (28)	226 (33)
	19-30	111	115 (8)	54 (9) ^E	62 (8)	80 (7)	104 (7)	134 (12)	168 (20)	192 (28)
	31-50	146	95 (8)	47 (6)	54 (7)	69 (8)	88 (10)	110 (12)	135 (16)	153 (19)
	51-70	184	91 (9)	58 (13) ^E	65 (12) ^E	76 (11)	90 (10)	105 (11)	122 (15)	133 (18)
	>70	143	88 (4)	46 (5)	54 (5)	68 (4)	86 (5)	106 (6)	128 (8)	143 (11)
	19+	584	97 (4)	46 (5)	55 (5)	70 (5)	91 (5)	117 (6)	147 (9)	170 (12)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.9 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Alberta, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both	1-3	169	95	(6)	48	(10) ^E	58	(9)	75	(7)	97	(6)	122	(8)	150	(15)	172	(22)
	4-8	281	127	(6)	79	(5)	87	(5)	102	(5)	122	(6)	148	(9)	175	(14)	193	(18)
Male	9-13	183	152	(9)	71	(15) ^E	85	(13)	113	(11)	150	(11)	194	(16)	239	(24)	269	(31)
	14-18	187	179	(10)	106	(10)	121	(11)	150	(11)	185	(12)	224	(14)	263	(17)	288	(19)
	19-30	223	134	(7)	74	(16) ^E	84	(14) ^E	103	(11)	127	(8)	154	(12)	182	(20)	201	(26)
	31-50	229	120	(11)	52	(16) ^E	63	(15) ^E	86	(14)	117	(13)	154	(16)	193	(23)	219	(29)
	51-70	197	91	(7)	34	(11) ^E	42	(10) ^E	59	(9)	83	(8)	115	(10)	150	(17)	175	(23)
	>70	72	75	(6)	41	(9) ^E	48	(8) ^E	61	(6)	76	(5)	92	(6)	110	(10)	123	(14)
	19+	721	113	(5)	44	(6)	55	(6)	76	(6)	106	(6)	144	(8)	186	(11)	214	(14)
Female	9-13	165	134	(8)	72	(12) ^E	84	(11)	107	(11)	136	(11)	169	(14)	204	(19)	227	(23)
	14-18	206	120	(7)	66	(7)	76	(7)	94	(8)	117	(9)	143	(11)	170	(12)	188	(14)
	19-30	191	107	(7)	51	(5)	61	(5)	80	(6)	105	(9)	134	(11)	166	(15)	188	(18)
	31-50	258	84	(5)	37	(8) ^E	44	(8) ^E	58	(8)	79	(7)	107	(8)	139	(13)	163	(18)
	51-70	249	85	(5)	52	(10) ^E	58	(9)	70	(7)	84	(6)	100	(7)	116	(10)	127	(13)
	>70	128	67	(5)	27	(7) ^E	33	(7) ^E	47	(7)	64	(7)	82	(9)	103	(13)	120	(17)
	19+	826	88	(3)	42	(4)	50	(4)	64	(4)	85	(4)	111	(5)	139	(7)	159	(9)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.10 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, British Columbia, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	192	95 (4)	53 (3)	60 (4)	71 (5)	88 (6)	107 (7)	129 (8)	144 (9)
	4-8	321	110 (3)	77 (6)	84 (6)	96 (5)	110 (4)	127 (6)	143 (8)	153 (10)
Male	9-13	226	147 (8)	87 (9)	98 (9)	120 (9)	149 (11)	185 (13)	222 (17)	248 (21)
	14-18	262	174 (11)	86 (13)	101 (13)	130 (14)	171 (15)	222 (17)	278 (23)	317 (29)
	19-30	197	145 (10)	69 (13) ^E	83 (13)	108 (12)	142 (13)	183 (16)	224 (23)	252 (28)
	31-50	282	135 (9)	55 (12) ^E	69 (12) ^E	94 (11)	125 (10)	162 (13)	206 (22)	236 (29)
	51-70	234	110 (7)	45 (6)	55 (6)	75 (6)	102 (8)	137 (11)	178 (16)	208 (20)
	>70	119	95 (10)	36 (9) ^E	45 (9) ^E	63 (9)	87 (10)	117 (14)	151 (18)	174 (20)
	19+	832	126 (5)	48 (4)	60 (5)	84 (5)	118 (5)	159 (7)	205 (11)	236 (14)
Female	9-13	226	128 (6)	83 (9)	92 (9)	108 (8)	129 (8)	154 (10)	181 (14)	198 (18)
	14-18	242	121 (8)	55 (9) ^E	67 (9)	89 (9)	117 (9)	151 (11)	188 (16)	214 (20)
	19-30	208	107 (7)	54 (9) ^E	64 (9)	83 (8)	108 (9)	138 (11)	170 (15)	191 (19)
	31-50	263	92 (6)	36 (4)	44 (4)	62 (5)	86 (6)	114 (7)	142 (9)	160 (10)
	51-70	322	78 (4)	31 (5)	39 (5)	53 (4)	73 (5)	97 (6)	127 (9)	149 (11)
	>70	198	84 (6)	42 (6)	48 (6)	60 (6)	78 (8)	102 (11)	126 (17)	143 (22)
	19+	991	90 (3)	38 (3)	46 (3)	62 (3)	84 (3)	112 (4)	142 (6)	163 (7)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.11 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Atlantic Region, 2004^{1,2}

Table 1. Usual Sugars (g/day) Consumed from 1993, by Age Group, Household Population, and Region, 2000										
Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both										
	1-3	348	103 (3)	65 (6)	73 (6)	86 (5)	102 (4)	121 (5)	138 (7)	150 (8)
	4-8	554	131 (4)	85 (6)	95 (6)	112 (5)	132 (5)	154 (6)	177 (8)	192 (10)
Male										
	9-13	409	154 (7)	86 (9)	97 (9)	120 (8)	148 (8)	181 (10)	216 (14)	240 (17)
	14-18	414	170 (8)	87 (9)	101 (9)	126 (9)	162 (10)	210 (13)	265 (18)	304 (22)
	19-30	311	140 (10)	74 (11)	84 (10)	104 (9)	132 (9)	164 (13)	197 (19)	220 (24)
	31-50	489	125 (7)	53 (6)	64 (6)	88 (6)	120 (8)	159 (10)	201 (15)	231 (18)
	51-70	575	98 (4)	51 (6)	59 (6)	73 (5)	93 (5)	119 (7)	147 (10)	164 (13)
	>70	239	98 (6)	40 (5)	49 (5)	68 (7)	93 (8)	124 (9)	156 (11)	176 (12)
	19+	1614	118 (4)	51 (3)	61 (3)	82 (3)	112 (4)	148 (5)	186 (8)	212 (10)
Female										
	9-13	355	125 (5)	71 (6)	81 (6)	101 (6)	126 (6)	152 (7)	177 (8)	194 (10)
	14-18	410	124 (6)	60 (6)	71 (6)	91 (6)	118 (7)	152 (10)	190 (14)	217 (17)
	19-30	384	114 (7)	61 (9)	71 (9)	92 (9)	118 (10)	145 (11)	172 (13)	189 (15)
	31-50	585	90 (4)	36 (4)	46 (4)	65 (4)	90 (5)	118 (7)	149 (9)	171 (10)
	51-70	711	79 (2)	40 (4)	47 (4)	60 (3)	76 (3)	95 (3)	113 (5)	125 (6)
	>70	342	83 (4)	39 (4)	45 (5)	58 (5)	77 (6)	101 (8)	130 (13)	149 (17)
	19+	2022	90 (2)	41 (2)	50 (2)	67 (2)	88 (3)	113 (3)	140 (5)	160 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.12 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Prairie Region, 2004^{1,2}

Sex	Age (years)	Percentiles (and SE) of usual intake								
		n	Mean (SE)	5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	622	97 (4)	45 (5)	55 (4)	73 (4)	96 (4)	121 (5)	150 (8)	172 (12)
	4-8	919	124 (4)	76 (6)	84 (5)	100 (4)	120 (4)	145 (7)	171 (10)	188 (14)
Male	9-13	579	152 (6)	85 (9)	97 (8)	120 (7)	150 (7)	185 (9)	221 (14)	245 (17)
	14-18	634	181 (7)	108 (13)	123 (12)	151 (10)	185 (10)	224 (13)	264 (21)	290 (27)
	19-30	578	135 (6)	64 (9)	75 (9)	97 (7)	125 (6)	159 (9)	195 (14)	219 (17)
	31-50	693	119 (7)	47 (8) ^E	58 (8)	82 (8)	114 (8)	154 (11)	198 (16)	230 (20)
	51-70	596	93 (5)	38 (5)	46 (5)	64 (5)	88 (5)	119 (7)	153 (10)	176 (12)
	>70	296	84 (4)	41 (4)	50 (4)	65 (4)	82 (3)	102 (4)	126 (7)	145 (8)
	19+	2163	113 (3)	44 (3)	55 (3)	76 (3)	106 (4)	145 (5)	189 (7)	220 (9)
Female	9-13	533	135 (5)	77 (7)	88 (6)	108 (7)	134 (7)	164 (9)	196 (11)	217 (12)
	14-18	638	127 (5)	70 (9)	81 (8)	100 (7)	125 (7)	156 (9)	189 (12)	212 (16)
	19-30	499	113 (5)	53 (6)	64 (6)	84 (5)	110 (6)	140 (8)	174 (12)	198 (16)
	31-50	716	87 (4)	43 (6)	50 (6)	64 (5)	83 (5)	107 (6)	133 (8)	151 (11)
	51-70	745	87 (3)	45 (5)	53 (5)	66 (4)	84 (4)	105 (5)	126 (6)	141 (8)
	>70	510	76 (3)	32 (4)	40 (4)	55 (4)	74 (4)	94 (5)	117 (7)	134 (9)
	19+	2470	91 (2)	41 (2)	49 (2)	65 (2)	87 (3)	113 (3)	145 (5)	167 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Table 40.13 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Canada excluding territories, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	2117	100 (2)	51 (3)	61 (2)	77 (2)	98 (2)	122 (3)	149 (4)	168 (5)
	4-8	3235	122 (2)	73 (3)	83 (2)	100 (2)	121 (2)	145 (3)	170 (4)	187 (6)
	9-13	2080	155 (3)	90 (4)	102 (4)	125 (4)	154 (4)	189 (5)	226 (7)	252 (9)
Male	14-18	2288	173 (4)	91 (4)	106 (4)	135 (4)	174 (5)	219 (6)	267 (8)	300 (10)
	19-30	1804	137 (3)	68 (5)	80 (5)	102 (5)	132 (4)	167 (5)	202 (7)	226 (9)
	31-50	2596	117 (3)	46 (3)	58 (3)	80 (3)	112 (4)	151 (4)	192 (6)	220 (9)
	51-70	2550	102 (2)	45 (3)	55 (2)	73 (2)	97 (2)	125 (3)	156 (5)	177 (6)
	>70	1520	93 (3)	41 (3)	49 (3)	66 (3)	89 (3)	117 (4)	147 (6)	168 (7)
	19+	8470	115 (2)	48 (2)	58 (2)	80 (2)	109 (2)	146 (2)	185 (3)	212 (4)
	9-13	1980	129 (2)	76 (3)	86 (3)	104 (3)	127 (3)	153 (4)	181 (5)	199 (6)
	14-18	2256	126 (2)	65 (3)	77 (3)	98 (3)	126 (3)	158 (4)	192 (6)	215 (7)
Female	19-30	1854	107 (3)	52 (3)	61 (3)	79 (3)	104 (3)	133 (4)	163 (5)	183 (6)
	31-50	2686	92 (2)	38 (2)	47 (2)	64 (2)	86 (3)	114 (3)	146 (5)	169 (6)
	51-70	3200	85 (2)	41 (2)	49 (2)	62 (2)	81 (2)	105 (2)	130 (3)	148 (4)
	>70	2610	82 (2)	39 (2)	46 (2)	60 (2)	79 (2)	100 (3)	123 (4)	140 (5)
	19+	10350	92 (1)	41 (1)	50 (1)	66 (1)	87 (1)	113 (2)	143 (3)	164 (3)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Appendix A: Table Footnotes

The following footnotes apply to all of the summary data tables presented in Section II of this report.

1. The survey excludes from its target population those living in the three territories, individuals living on Indian reserves or on Crown lands, residents of institutions, full-time members of the Canadian Forces and residents of certain remote regions.
2. The tables exclude pregnant and breastfeeding females, subject to another set of nutritional recommendations. The sample of pregnant and breastfeeding females is not large enough to allow for reliable estimates.
3. Sample size and mean intake are based on the first 24-hour dietary recall (first day of interview) only.
4. Intakes are based on food consumption only. Intakes from vitamin and mineral supplements are not included. Inferences about the prevalence of nutrient excess or inadequacy based on intakes from food alone may respectively underestimate or overestimate the prevalences based on total nutrient intakes from both food and supplements.
5. The intake distribution (percentiles and percentage above or below a cut-off when applicable) was adjusted to remove within-individual variability using Software for Intake Distribution Estimation (SIDE) (Iowa State University, 1996) and the method presented in Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440-1449.
6. In some cases, within-individual variance was estimated at the regional or national level and applied at the provincial level. For more details, see Section II.4: Measuring Sampling Variability with Bootstrap Replication in Volume 1 of the *Nutrient Intakes from Food* report series.
7. Bootstrapping techniques were used to produce the coefficient of variation (CV) and the standard error (SE).
8. AMDR is the Acceptable Macronutrient Distribution Range, expressed as a percentage of total energy intake. Intakes inside the range (shown in the AMDR columns) are associated with a reduced risk of chronic disease while providing adequate intakes of essential nutrients. For further information on AMDR see the Health Canada publication *Canadian Community Health Survey*,

Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data, Section 2.1.5, p. 27.

The applications of the AMDRs for essential fatty acids to group assessment are not the same as for the other macronutrients. The lower boundaries for the AMDR for linoleic and alpha-linolenic acids are not based on the same type of endpoints as the boundaries for total fat and carbohydrate. The boundaries for fat and carbohydrate are set based on evidence indicating increased risk for coronary heart diseases and the lower bound of the AMDR for both n-6 (linoleic) and n-3 (alpha-linolenic) fatty acids is based on the percent of energy from these fatty acids needed to provide the AI for these nutrients. The AI, in turn, is based on the median intake of both linoleic and alpha-linolenic acid in the United States, where essential fatty acid deficiency is non-existent in the healthy population.

Thus, by definition about half the population has intakes of these fatty acids below the AI and therefore outside the AMDR. In other words, based on the AI, one would conclude that the population is "adequate" with respect to linoleic and alpha-linolenic acids, while based on the AMDR a different conclusion (i.e. that 50% of the population has intakes below the AMDR) would be reached. Therefore, the lower bound of the AMDRs for linoleic and alpha-linolenic acids should not be used in the assessment of population intakes.

9. EAR is the Estimated Average Requirement. The level of intake at the EAR (shown in the EAR columns) is the average daily intake level that is estimated to meet the requirement, as defined by the specified indicator of adequacy, in half of the apparently healthy individuals in a DRI age–sex group. For further information on EAR see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.1, p. 23.
10. AI is the Adequate Intake. The level of intake at the AI (shown in the AI columns) is the recommended average daily intake level based on observed or experimentally determined approximations or estimates of nutrient intake by a group or groups of apparently healthy people that are assumed to be adequate. It is developed when an EAR cannot be determined. The percentage of the population having a usual intake above the AI (shown in the %>AI columns) almost certainly meets their needs. The adequacy of intakes below the AI cannot be assessed, and should not be interpreted as being inadequate. For further information on AI see the Health Canada

publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.3, p. 25.

11. UL is the Tolerable Upper Intake Level. The level of intake at the UL (shown in the UL columns) is the highest average daily intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population. For further information on UL see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.4, p. 26.
12. For a more detailed understanding of DRIs and their interpretation when assessing intakes of particular nutrients, consult the summary of the series of publications on DRIs published by the Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*.
13. In terms of precision, the estimate 0.0 with a standard error of 0.0 refers to a standard error smaller than 0.1%.

Appendix B: Justification for Excluding Nutrients from Volume 2 and Volume 3

Volume 1 of the compendium contained data on 13 nutrients, including 6 nutrients expressed as a percent of total energy. There were originally 31 nutrients scheduled to be released in Volumes 2 and 3 of the compendium, but for a variety of reasons some of these nutrients are not included. Decisions to omit these nutrients were made jointly by representatives from Statistics Canada and Health Canada.

Exclusions and changes to the list of nutrients that were to be included in Volumes 2 and 3 of the compendium are as follows:

Total milligrams of folic acid

Folic acid is found in small amounts in a number of foods. Most respondents consumed a small amount of folic acid, which resulted in a bimodal distribution of folic acid intake. As a result, it was very difficult to normalize the distribution, which meant that SIDE was unable to calculate usual intake.

One of the steps that SIDE uses to estimate usual intake is to transform the data into a normal distribution. Assessing SIDE's ability to perform this transformation rests on measuring the Anderson-Darling (A-D) score for normality. The A-D score is a statistic that measures how close a distribution is to a normal distribution. Any A-D score less than 0.576 is considered to be sufficiently normal for SIDE to continue without warning. Typically, SIDE will be able to transform 95% of the domains without error using the default SIDE options. The remaining 5% of domains will typically score higher than 0.576 but usually less than 1.0. Adjusting the SIDE options will usually reduce the A-D to within the limit. In the case of folic acid, more than half of the provincial domains had an A-D score above the 0.576 threshold and many domains scored higher than 2. The nature of the data simply does not allow SIDE to produce proper estimates for the usual intake of folic acid.

Total grams of alcohol

Alcohol is consumed differently than other nutrients. For most respondents, alcohol is not part of their daily intake of food, but rather is something that is consumed occasionally. In this sense, in terms of analysis, alcohol behaves more like a food than a nutrient. In order for SIDE to estimate the usual intake of foods, many recalls are needed to capture enough occurrences of the particular food. Thus, two recalls are not enough to calculate the usual intake of alcohol.

Percent of energy from alcohol

The difficulty in estimating a usual intake for alcohol causes similar problems for expressing that intake as a percent of total energy.

Caffeine

Caffeine also is consumed differently than other nutrients. The usual intake of caffeine could not be calculated due to the same issues as folic acid and alcohol. Many respondents reported zero or small levels of caffeine intake. Therefore, it is difficult for SIDE to properly model the data with only two dietary recalls.

Based on the changes above, the list of nutrients included in Volume 1 and the revised list of nutrients included in Volumes 2 and 3 are as follows:

List of Nutrients Included in the Three-Volume Set		
Volume 1	Volume 2	Volume 3
Total Energy	Folate (DFE)	Folacin
Percentage of total energy intake from fats	Iron	Linolenic acid (g, % energy)
Percentage of total energy intake from protein	Linoleic acid (g, % energy)	Moisture
Percentage of total energy intake from carbohydrates	Magnesium	Naturally occurring folate
Percentage of total energy intake from saturated fats	Niacin	Protein
Percentage of total energy intake from monounsaturated fats	Phosphorus	Total carbohydrates
Percentage of total energy intake from polyunsaturated fats	Potassium	Total fats
Total dietary fibre	Riboflavin	Total monounsaturated fats
Cholesterol	Thiamin	Total polyunsaturated fats
Vitamin A	Vitamin B ₆	Total saturated fats
Vitamin C	Vitamin B ₁₂	Total sugars
Calcium	Vitamin C by smoking status	
Sodium	Vitamin D	
	Zinc	

Appendix C: References

Department of Statistics and Center for Agricultural and Rural Development, Iowa State University: *A User's Guide to SIDE, Software for Intake Distribution Estimation Version 1.0*. Technical Report 96-TR 30. Ames, IA: Iowa State University Statistical Laboratory, 1996. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr30.pdf

Dodd KW: *A Technical Guide to C-SIDE, Software for Intake Distribution Estimation*. Technical Report 96-TR 32, Dietary Assessment Research Series Report 9. Ames, IA: Department of Statistics and Center for Agricultural and Rural Development, Iowa State University, 2006. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr32.pdf

Health Canada: *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*. Ottawa: Publications, Health Canada, 2006. Available at: www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_focus-volet_escc-eng.php

Institute of Medicine: *Dietary Reference Intakes: A Risk Assessment Model for Establishing Upper Intake Levels for Nutrients*. Washington, DC: National Academy Press, 1998. Available at: <http://books.nap.edu/catalog/6432.html>

Institute of Medicine: *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin and Choline*. Washington, DC: National Academy Press, 1998. Available at: www.nap.edu/catalog.php?record_id=6015

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academy Press, 2000. Available at: <http://books.nap.edu/catalog/9956.html>

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Planning*. Washington, DC: National Academy Press, 2003. Available at: <http://books.nap.edu/catalog/10609.html>

Institute of Medicine: *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington, DC: National Academy Press, 2004. Available at: <http://books.nap.edu/catalog/10925.html>

Institute of Medicine: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. Washington, DC:

National Academy Press, 2005. Available at:

<http://books.nap.edu/catalog/10490.html>

Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: National Academy Press, 2006.

Available at: www.nap.edu/topics.php?topic=380

Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440–1449

Nusser SM, Fuller WA, Guenther PM: Estimating usual dietary intake distributions: adjusting for measurement error and nonnormality in 24-hour food intake data. In L Lyberg, P Biemer, M Collins et al., eds. *Survey Measurement and Process Quality*, New York: John Wiley & Sons, 1997.

Statistics Canada: *Canadian Community Health Survey (CCHS) Cycle 2.2—Nutrition: Software for Intake Distribution Estimation (SIDE) Documentation*. Ottawa, 2007.

Available at: www.statcan.ca/english/sdds/document/5049_D22_T9_V1_E.pdf