



Health
Canada

Santé
Canada

Statistics
Canada

Statistique
Canada

Canadian Community Health Survey Cycle 2.2, Nutrition (2004)

Nutrient Intakes from Food

Provincial, Regional and National Summary Data Tables
Volume 3

Revised February 2009

Note:

This PDF contains the 12 data tables for Saskatchewan as well as the Appendices.



Canada 

Table of Contents (for the full report)

	Acknowledgements	i
	List of Tables.....	iii
	List of Appendices	vii
	List of Abbreviations.....	viii
I	Introduction.....	1
II	Summary Data Tables	3
	<i>(table numbering continued from Volume 2)</i>	
	29. Folic acid ($\mu\text{g/d}$): Usual intakes from food	3
	30. Linolenic acid (g/d): Usual intakes from food.....	17
	31. Percentage of total energy intake from linolenic acid.....	31
	32. Moisture (g/d): Usual intakes from food	45
	33. Naturally occurring folate ($\mu\text{g/d}$): Usual intakes from food.....	59
	34. Protein (g/d): Usual intakes from food.....	73
	35. Total carbohydrates (g/d): Usual intakes from food	87
	36. Total fats (g/d): Usual intakes from food	101
	37. Total monounsaturated fats (g/d): Usual intakes from food.....	115
	38. Total polyunsaturated fats (g/d): Usual intakes from food	129
	39. Total saturated fats (g/d): Usual intakes from food	143
	40. Total sugars (g/d): Usual intakes from food.....	157

List of Appendices

Appendix A: Table Footnotes.....	171
Appendix B: Justification for Excluding Nutrients from Volume 2 and Volume 3.....	175
List of Nutrients Included in the Three-Volume Set.....	176
Appendix C: References.....	177

Table 29.8 Folic acid (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2,3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	189 (9)	118 (13)	131 (12)	153 (11)	179 (11)	211 (14)	247 (21)	274 (28)
	4-8	213	275 (12)	174 (21)	193 (18)	226 (14)	269 (13)	317 (19)	365 (28)	397 (36)
Male	9-13	122	345 (35)	202 (32)	227 (31)	274 (31)	340 (36)	428 (52)	527 (79)	594 (99) ^E
	14-18	150	443 (44)	264 (51) ^E	295 (48)	353 (44)	430 (44)	521 (62)	618 (98)	683 (127) ^E
	19-30	106	435 (41)	202 (42) ^E	238 (41) ^E	308 (38)	405 (40)	525 (53)	655 (78)	744 (100)
	31-50	155	388 (24)	220 (45) ^E	248 (41)	301 (34)	379 (30)	473 (44)	568 (70)	630 (90)
	51-70	122	368 (46)	208 (33)	234 (31)	278 (30)	337 (41)	438 (85) ^E	616 (199) ^E	^F
	>70	88	316 (18)	192 (28)	213 (26)	252 (23)	304 (21)	368 (26)	438 (40)	487 (52)
	19+	471	385 (19)	187 (16)	219 (15)	281 (14)	361 (16)	467 (24)	610 (50)	723 (81)
Female	9-13	103	324 (25)	237 (35)	254 (33)	285 (32)	322 (32)	363 (38)	404 (48)	430 (55)
	14-18	142	296 (18)	175 (17)	196 (17)	235 (18)	285 (21)	348 (29)	423 (39)	480 (49)
	19-30	111	288 (19)	160 (28) ^E	183 (24)	224 (20)	278 (20)	345 (31)	419 (51)	472 (69)
	31-50	146	315 (21)	203 (39) ^E	222 (35)	257 (30)	302 (27)	359 (30)	424 (46)	471 (62)
	51-70	184	314 (13)	251 (37)	265 (32)	289 (23)	317 (16)	348 (24)	377 (41)	396 (54)
	>70	143	248 (13)	178 (23)	193 (21)	218 (18)	251 (16)	289 (19)	328 (26)	354 (32)
	19+	584	299 (10)	184 (15)	204 (15)	243 (13)	293 (13)	352 (15)	416 (20)	461 (25)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for folic acid.

³ There are two chemical forms in foods that contribute to folate bioactivity: naturally occurring folate called “food folate” and the added synthetic form of folate called “folic acid.” The term “folic acid” is a measuring unit referring to the simple arithmetic sum of the content of both food folate and folic acid in foods (in micrograms).

For additional footnotes common to all tables, see Appendix A.

Table 30.8 Linolenic acid (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI	(SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)			
Both	1-3	129	1.06 (0.11)	0.63 (0.15) ^E	0.71 (0.14) ^E	0.88 (0.14)	1.11 (0.16)	1.39 (0.23) ^E	1.70 (0.35) ^E	1.92 (0.45) ^E	0.7	90.8	(9.7)
	4-8	213	1.40 (0.10)	0.81 (0.12)	0.90 (0.11)	1.07 (0.10)	1.31 (0.09)	1.60 (0.13)	1.92 (0.20)	2.14 (0.27)	0.9	89.8	(6.2)
Male	9-13	122	2.23 (0.57) ^E	0.93 (0.29) ^E	1.10 (0.30) ^E	1.45 (0.33) ^E	2.02 (0.43) ^E	2.84 (0.65) ^E	3.91 (1.03) ^E	4.74 (1.38) ^E	1.2	86.1	(10.4)
	14-18	150	2.23 (0.18)	1.39 (0.14)	1.55 (0.16)	1.84 (0.20)	2.23 (0.25)	2.72 (0.32)	3.26 (0.41)	3.63 (0.48)	1.6	87.7	(8.8)
	19-30	106	2.87 (0.43)	^F	1.12 (0.35) ^E	1.63 (0.35) ^E	2.44 (0.41)	3.59 (0.62) ^E	5.00 (1.08) ^E	6.07 (1.57) ^E	1.6	76.1	(10.7)
	31-50	155	2.10 (0.16)	0.99 (0.29) ^E	1.15 (0.27) ^E	1.47 (0.23)	1.95 (0.20)	2.55 (0.26)	3.19 (0.42)	3.63 (0.56)	1.6	68.1	(13.1) ^E
	51-70	122	2.10 (0.17)	1.71 (0.24)	1.84 (0.26)	2.08 (0.28)	2.37 (0.32)	2.69 (0.35)	3.00 (0.39)	3.20 (0.42)	1.6	97.5	(11.7)
	>70	88	2.37 (0.30)	1.02 (0.31) ^E	1.20 (0.31) ^E	1.58 (0.31) ^E	2.09 (0.35) ^E	2.77 (0.50) ^E	3.58 (0.76) ^E	4.14 (1.00) ^E	1.6	74.0	(14.5) ^E
	19+	471	2.31 (0.13)	1.06 (0.15)	1.24 (0.15)	1.61 (0.15)	2.17 (0.16)	2.91 (0.22)	3.76 (0.35)	4.38 (0.48)	1.6	75.5	(6.5)
Female	9-13	103	1.55 (0.16)	1.06 (0.23) ^E	1.15 (0.22) ^E	1.34 (0.22)	1.59 (0.22)	1.88 (0.28)	2.20 (0.42) ^E	2.42 (0.56) ^E	1.0	97.0	(8.6)
	14-18	142	1.72 (0.13)	0.86 (0.17) ^E	1.01 (0.16)	1.31 (0.16)	1.73 (0.19)	2.24 (0.27)	2.78 (0.37)	3.16 (0.46)	1.1	86.0	(8.0)
	19-30	111	1.72 (0.14)	1.10 (0.28) ^E	1.22 (0.26) ^E	1.45 (0.23)	1.76 (0.22)	2.13 (0.29)	2.52 (0.45) ^E	2.79 (0.60) ^E	1.1	94.9	(9.7)
	31-50	146	1.55 (0.14)	0.90 (0.12)	1.00 (0.13)	1.17 (0.14)	1.39 (0.15)	1.64 (0.18)	1.89 (0.21)	2.05 (0.24)	1.1	81.9	(9.9)
	51-70	184	2.10 (0.22)	^F	0.98 (0.28) ^E	1.32 (0.26) ^E	1.84 (0.25)	2.57 (0.32)	3.46 (0.54)	4.14 (0.77) ^E	1.1	85.3	(8.6)
	>70	143	1.54 (0.15)	0.82 (0.26) ^E	0.95 (0.24) ^E	1.20 (0.22) ^E	1.57 (0.22)	2.07 (0.30)	2.66 (0.48) ^E	3.09 (0.65) ^E	1.1	81.4	(12.4)
	19+	584	1.73 (0.09)	0.88 (0.16) ^E	1.01 (0.15)	1.27 (0.13)	1.63 (0.12)	2.09 (0.15)	2.61 (0.25)	2.99 (0.35)	1.1	85.2	(6.6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 31.8 Percentage of total energy intake from linolenic acid, by DRI age–sex group, household population, Saskatchewan, 2004^{1–3}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	0.64 (0.07)	0.44 (0.08)^E	0.48 (0.08)	0.55 (0.08)	0.65 (0.09)	0.76 (0.12)	0.90 (0.17)^E	0.99 (0.21)^E
	4-8	213	0.61 (0.03)	0.47 (0.02)	0.49 (0.02)	0.54 (0.02)	0.59 (0.03)	0.65 (0.03)	0.72 (0.04)	0.76 (0.04)
Male	9-13	122	0.75 (0.10)	0.41 (0.08)^E	0.46 (0.08)^E	0.56 (0.08)	0.70 (0.09)	0.88 (0.14)	1.09 (0.21)^E	1.24 (0.27)^E
	14-18	150	0.67 (0.04)	0.48 (0.04)	0.51 (0.04)	0.58 (0.05)	0.66 (0.06)	0.76 (0.07)	0.86 (0.09)	0.93 (0.11)
	19-30	106	0.88 (0.11)	0.54 (0.07)	0.59 (0.08)	0.70 (0.10)	0.83 (0.12)	1.00 (0.15)	1.19 (0.19)	1.32 (0.22)^E
	31-50	155	0.77 (0.07)	0.56 (0.11)^E	0.59 (0.10)	0.65 (0.08)	0.73 (0.07)	0.82 (0.09)	0.92 (0.14)	0.98 (0.19)^E
	51-70	122	0.89 (0.08)	0.62 (0.07)	0.68 (0.08)	0.80 (0.10)	0.95 (0.12)	1.11 (0.14)	1.27 (0.17)	1.37 (0.19)
	>70	88	1.05 (0.13)	0.49 (0.12)^E	0.56 (0.12)^E	0.69 (0.12)^E	0.90 (0.14)	1.20 (0.20)	1.57 (0.31)^E	1.85 (0.41)^E
	19+	471	0.86 (0.04)	0.54 (0.10)^E	0.60 (0.09)	0.71 (0.07)	0.84 (0.06)	1.01 (0.07)	1.19 (0.11)	1.30 (0.15)
Female	9-13	103	0.70 (0.09)	0.41 (0.09)^E	0.46 (0.09)^E	0.55 (0.08)	0.68 (0.09)	0.86 (0.13)	1.09 (0.22)^E	1.27 (0.32)^E
	14-18	142	0.72 (0.06)	0.49 (0.05)	0.53 (0.05)	0.61 (0.06)	0.72 (0.07)	0.85 (0.09)	0.98 (0.12)	1.07 (0.14)
	19-30	111	0.80 (0.06)	0.65 (0.12)^E	0.69 (0.12)^E	0.76 (0.10)	0.85 (0.10)	0.95 (0.11)	1.05 (0.14)	1.12 (0.18)
	31-50	146	0.73 (0.06)	0.46 (0.06)	0.50 (0.06)	0.58 (0.06)	0.68 (0.07)	0.80 (0.08)	0.93 (0.09)	1.01 (0.11)
	51-70	184	0.99 (0.09)	0.51 (0.13)^E	0.58 (0.12)^E	0.71 (0.11)	0.89 (0.11)	1.15 (0.14)	1.46 (0.23)	1.70 (0.31)^E
	>70	143	0.84 (0.06)	0.58 (0.10)^E	0.63 (0.10)	0.72 (0.09)	0.85 (0.08)	1.01 (0.10)	1.18 (0.17)	1.29 (0.23)^E
	19+	584	0.83 (0.04)	0.51 (0.08)	0.56 (0.08)	0.66 (0.07)	0.80 (0.05)	0.97 (0.06)	1.17 (0.10)	1.31 (0.15)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Linolenic acid is an alternative name for α -linolenic acid (n-3).

³ AMDR is the Acceptable Macronutrient Distribution Range. For additional detail, see footnote 8 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 32.8 Moisture (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake							AI ³	% >AI (SE)
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)		
Both	1-3	129	1461 (64)	910 (76)	1010 (73)	1186 (77)	1458 (88)	1784 (124)	2073 (161)	2258 (188)	1300	64.1 (8.2)
	4-8	213	1655 (57)	1137 (122)	1240 (108)	1427 (85)	1660 (69)	1920 (87)	2179 (132)	2347 (168)	1700	45.6 (8.9) ^E
Male	9-13	122	1968 (112)	1195 (155)	1319 (157)	1586 (161)	1958 (158)	2368 (159)	2754 (196)	3002 (239)	2400	^F
	14-18	150	3075 (161)	1760 (247)	1995 (229)	2447 (197)	3052 (191)	3780 (270)	4558 (430)	5087 (564)	3300	40.3 (9.2) ^E
	19-30	106	3161 (204)	1677 (255)	1901 (245)	2334 (225)	2918 (209)	3640 (251)	4446 (370)	5015 (478)	3700	23.4 (6.9) ^E
	31-50	155	2951 (178)	1608 (227)	1802 (220)	2191 (214)	2745 (220)	3467 (269)	4304 (403)	4911 (528)	3700	^F
	51-70	122	2672 (131)	1802 (220)	1955 (195)	2240 (158)	2607 (155)	3035 (235)	3481 (377)	3779 (494)	3700	^F
	>70	88	2499 (110)	1509 (194)	1713 (171)	2082 (143)	2516 (136)	2972 (152)	3448 (217)	3789 (298)	3700	^F
	19+	471	2872 (97)	1587 (77)	1770 (84)	2161 (96)	2729 (114)	3365 (146)	4146 (224)	4774 (310)	3700	16.7 (3.5) ^E
Female	9-13	103	2017 (117)	1331 (143)	1463 (137)	1702 (131)	2005 (140)	2379 (173)	2811 (233)	3129 (290)	2100	42.6 (11.1) ^E
	14-18	142	2428 (131)	1518 (155)	1706 (155)	2047 (158)	2478 (175)	2988 (212)	3544 (276)	3932 (336)	2300	60.5 (9.6)
	19-30	111	2867 (149)	1670 (248)	1851 (235)	2207 (213)	2710 (205)	3360 (262)	4063 (386)	4517 (481)	2700	50.5 (11.6) ^E
	31-50	146	3328 (187)	2216 (357)	2444 (322)	2835 (270)	3284 (230)	3745 (238)	4168 (295)	4424 (345)	2700	81.1 (11.1)
	51-70	184	2810 (107)	1941 (214)	2113 (189)	2432 (147)	2837 (123)	3305 (172)	3785 (273)	4103 (350)	2700	58.6 (9.5)
	>70	143	2333 (89)	1494 (141)	1667 (130)	1973 (118)	2338 (119)	2729 (143)	3102 (177)	3335 (203)	2700	26.6 (7.7) ^E
	19+	584	2938 (83)	1818 (103)	2020 (102)	2401 (102)	2905 (109)	3506 (133)	4110 (169)	4481 (193)	2700	60.2 (5.4)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² The term “moisture” includes water from all food and beverage sources.

³ AI is the Adequate Intake. For additional detail, see footnote 10 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 33.8 Naturally occurring folate (µg/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	121 (6)	68 (10)	78 (9)	94 (8)	114 (8)	139 (10)	172 (15)	200 (22)
	4-8	213	149 (8)	89 (12)	100 (11)	119 (9)	143 (9)	172 (12)	202 (17)	221 (22)
Male	9-13	122	189 (13)	122 (19)	133 (18)	156 (16)	187 (15)	225 (22)	267 (35)	295 (47)
	14-18	150	260 (46) ^E	157 (39) ^E	178 (40) ^E	218 (44) ^E	260 (48) ^E	301 (57) ^E	346 (83) ^E	378 (108) ^E
	19-30	106	255 (21)	99 (29) ^E	122 (28) ^E	171 (25)	241 (25)	332 (35)	435 (60)	508 (83)
	31-50	155	258 (21)	137 (38) ^E	158 (35) ^E	199 (31)	255 (27)	325 (34)	401 (53)	454 (70)
	51-70	122	279 (46)	142 (32) ^E	163 (31) ^E	198 (32)	246 (43) ^E	336 (78) ^E	^F	^F
	>70	88	245 (16)	130 (21)	147 (19)	180 (17)	226 (17)	287 (25)	361 (42)	417 (59)
	19+	471	262 (15)	120 (13)	143 (12)	187 (11)	243 (12)	321 (19)	427 (41)	517 (72)
Female	9-13	103	193 (23)	124 (23) ^E	135 (23) ^E	155 (23)	182 (26)	216 (32)	251 (41)	275 (48) ^E
	14-18	142	166 (13)	86 (15) ^E	99 (14)	124 (14)	158 (16)	203 (22)	258 (35)	300 (45)
	19-30	111	184 (14)	91 (15) ^E	107 (14)	136 (13)	173 (15)	222 (22)	285 (38)	335 (54)
	31-50	146	196 (12)	114 (24) ^E	126 (22) ^E	148 (19)	177 (17)	216 (19)	260 (29)	293 (40)
	51-70	184	223 (12)	152 (26) ^E	167 (23)	191 (18)	222 (16)	256 (21)	293 (33)	318 (44)
	>70	143	176 (8)	111 (13)	123 (12)	147 (10)	177 (10)	211 (13)	246 (18)	269 (23)
	19+	584	198 (7)	106 (10)	122 (9)	150 (9)	187 (9)	234 (11)	289 (15)	330 (21)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for naturally occurring folate.

For additional footnotes common to all tables, see Appendix A.

Table 34.8 Protein (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	54 (3)	34 (4)	38 (4)	45 (4)	54 (4)	64 (5)	74 (6)	81 (7)
	4-8	213	65 (2)	48 (5)	51 (4)	57 (3)	65 (3)	73 (4)	81 (6)	86 (7)
	9-13	122	88 (6)	68 (6)	73 (6)	81 (7)	91 (8)	102 (9)	114 (11)	122 (13)
Male	14-18	150	114 (6)	67 (8)	75 (8)	91 (7)	111 (8)	135 (9)	159 (12)	175 (15)
	19-30	106	108 (7)	47 (10) ^E	58 (10) ^E	78 (8)	104 (8)	133 (11)	164 (16)	183 (20)
	31-50	155	97 (5)	50 (8) ^E	59 (8)	75 (7)	95 (6)	117 (7)	139 (9)	153 (11)
	51-70	122	92 (5)	60 (11) ^E	67 (10)	79 (8)	94 (7)	111 (9)	127 (13)	137 (16)
	>70	88	85 (5)	50 (7)	56 (7)	67 (6)	83 (6)	102 (8)	123 (13)	137 (17)
	19+	471	97 (3)	48 (4)	57 (4)	73 (3)	95 (4)	119 (5)	144 (6)	161 (8)
	9-13	103	72 (4)	52 (7)	56 (6)	63 (6)	71 (5)	81 (6)	91 (8)	98 (10)
	14-18	142	72 (4)	44 (4)	49 (4)	59 (4)	71 (5)	84 (6)	98 (7)	107 (9)
Female	19-30	111	67 (3)	38 (6)	43 (5)	53 (5)	64 (5)	78 (6)	91 (7)	99 (9)
	31-50	146	79 (4)	36 (7) ^E	42 (7) ^E	59 (6)	75 (5)	89 (5)	102 (8)	115 (9)
	51-70	184	73 (3)	51 (3)	55 (3)	63 (4)	72 (4)	82 (5)	93 (7)	100 (7)
	>70	143	68 (4)	38 (5)	43 (5)	53 (5)	68 (5)	86 (7)	108 (10)	123 (13)
	19+	584	73 (2)	40 (3)	46 (3)	57 (3)	70 (3)	87 (3)	102 (5)	112 (6)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² Although DRIs for protein have been established on a “per kg body weight” basis, no DRIs have been established for the absolute amount of protein.

For additional footnotes common to all tables, see Appendix A.

Table 35.8 Total carbohydrates (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004¹

Sex	Age (years)	n Mean (SE)		Percentiles (and SE) of usual intake						EAR ²	% <EAR (SE)	
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)			
Both	1-3	129	198 (9)	122 (12)	137 (11)	164 (10)	197 (11)	233 (13)	269 (16)	293 (18)	100	F
	4-8	213	265 (9)	197 (18)	212 (15)	236 (10)	264 (8)	294 (14)	325 (23)	347 (29)	100	<3
Male	9-13	122	332 (19)	241 (26)	261 (25)	298 (22)	343 (22)	394 (27)	444 (36)	476 (45)	100	0.0 (0.0)
	14-18	150	394 (29)	281 (40)	305 (36)	346 (30)	395 (29)	450 (37)	505 (52)	541 (64)	100	<3
	19-30	106	333 (26)	167 (32) ^E	195 (31)	249 (28)	317 (28)	396 (34)	475 (46)	527 (56)	100	<3
	31-50	155	303 (22)	150 (31) ^E	179 (28)	229 (24)	289 (23)	362 (29)	449 (47)	516 (66)	100	<3
	51-70	122	232 (12)	137 (23) ^E	157 (20)	193 (16)	236 (16)	281 (22)	320 (30)	343 (36)	100	F
	>70	88	240 (12)	152 (15)	167 (14)	196 (14)	235 (15)	280 (19)	329 (27)	361 (34)	100	<3
	19+	471	283 (11)	139 (11)	166 (10)	214 (10)	273 (11)	346 (15)	430 (24)	492 (34)	100	<3
Female	9-13	103	289 (16)	202 (21)	219 (20)	251 (18)	289 (19)	333 (24)	378 (32)	407 (38)	100	<3
	14-18	142	298 (15)	202 (26)	226 (25)	267 (24)	314 (24)	366 (26)	421 (30)	459 (36)	100	<3
	19-30	111	242 (12)	149 (15)	165 (14)	194 (12)	233 (13)	278 (18)	325 (28)	356 (36)	100	<3
	31-50	146	221 (11)	158 (23)	169 (20)	189 (17)	214 (14)	243 (15)	272 (21)	291 (27)	100	<3
	51-70	184	217 (14)	151 (24)	165 (22)	189 (18)	217 (15)	248 (16)	278 (21)	298 (26)	100	<3
	>70	143	204 (8)	136 (10)	151 (9)	175 (9)	204 (11)	239 (14)	276 (18)	300 (21)	100	<3
	19+	584	221 (6)	141 (9)	156 (8)	182 (8)	216 (8)	257 (8)	299 (11)	328 (13)	100	<3

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² EAR is the Estimated Average Requirement. For additional detail, see footnote 9 in Appendix A.

For additional footnotes common to all tables, see Appendix A.

Table 36.8 Total fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	53 (3)	33 (5)	37 (5)	45 (4)	54 (4)	65 (5)	76 (6)	83 (8)
	4-8	213	71 (4)	52 (6)	56 (6)	62 (5)	71 (4)	80 (5)	89 (7)	95 (9)
Male	9-13	122	90 (9)	65 (9)	70 (9)	79 (9)	91 (9)	104 (12)	117 (15)	126 (18)
	14-18	150	105 (6)	79 (12)	85 (10)	94 (8)	105 (8)	117 (10)	129 (14)	137 (17)
	19-30	106	106 (8)	53 (14) ^E	63 (13) ^E	82 (11)	105 (10)	132 (12)	160 (18)	178 (24)
	31-50	155	98 (7)	53 (11) ^E	62 (10)	78 (8)	97 (8)	120 (11)	144 (18)	161 (23)
	51-70	122	87 (6)	48 (11) ^E	56 (9)	71 (8)	89 (7)	113 (12)	137 (19)	152 (24)
	>70	88	77 (5)	43 (8) ^E	49 (7)	61 (6)	75 (6)	91 (7)	107 (10)	118 (12)
	19+	471	95 (4)	46 (5)	56 (5)	72 (4)	94 (5)	119 (6)	147 (9)	166 (11)
Female	9-13	103	74 (5)	53 (9) ^E	57 (9)	65 (8)	76 (7)	87 (8)	98 (9)	105 (11)
	14-18	142	71 (4)	43 (4)	49 (4)	59 (4)	70 (5)	84 (7)	97 (9)	107 (11)
	19-30	111	68 (4)	43 (9) ^E	48 (8) ^E	56 (7)	67 (6)	79 (8)	90 (11)	98 (14)
	31-50	146	65 (3)	37 (7) ^E	42 (6)	50 (5)	60 (4)	72 (5)	84 (7)	91 (9)
	51-70	184	67 (3)	45 (7)	49 (6)	57 (5)	67 (4)	79 (6)	90 (9)	97 (12)
	>70	143	55 (4)	31 (7) ^E	35 (7) ^E	43 (6)	55 (5)	70 (7)	84 (10)	94 (12)
	19+	584	65 (2)	37 (3)	42 (3)	51 (3)	63 (3)	77 (3)	90 (4)	99 (5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for the absolute amount of total fats.

For additional footnotes common to all tables, see Appendix A.

Table 37.8 Total monounsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	19.9 (1.2)	11.0 (1.8)	12.7 (1.7)	15.9 (1.6)	20.1 (1.7)	24.9 (2.1)	29.7 (2.7)	32.9 (3.2)
	4-8	213	27.8 (1.9)	20.4 (2.7)	21.8 (2.4)	24.3 (2.0)	27.4 (1.8)	30.9 (2.3)	34.3 (3.5)	36.5 (4.4)
Male	9-13	122	34.1 (2.5)	23.1 (3.1)	25.1 (3.0)	29.1 (2.8)	34.4 (3.0)	40.9 (4.0)	47.6 (5.5)	52.0 (6.7)
	14-18	150	40.5 (2.6)	26.8 (2.3)	29.8 (2.4)	34.8 (2.8)	40.4 (3.4)	46.6 (4.4)	53.7 (5.2)	58.5 (5.7)
	19-30	106	45.4 (3.7)	19.1 (6.2) ^E	23.6 (5.8) ^E	32.4 (5.0)	44.1 (4.7)	57.8 (6.1)	71.8 (9.5)	81.0 (12.7)
	31-50	155	40.1 (3.1)	21.6 (5.0) ^E	25.1 (4.5) ^E	31.4 (3.8)	39.4 (3.7)	48.7 (5.0)	58.5 (7.4)	65.2 (9.3)
	51-70	122	36.0 (2.4)	21.8 (4.3) ^E	24.9 (3.8)	30.6 (3.1)	37.8 (3.2)	46.0 (4.8)	54.2 (7.3)	59.6 (9.3)
	>70	88	30.2 (1.8)	16.2 (3.2) ^E	18.6 (2.9)	23.0 (2.4)	28.9 (2.2)	35.6 (3.0)	42.6 (4.6)	47.2 (5.8)
	19+	471	39.0 (1.7)	19.1 (2.0)	22.7 (1.9)	29.4 (1.9)	38.3 (2.1)	49.1 (2.7)	60.7 (3.7)	68.6 (4.6)
Female	9-13	103	27.9 (1.9)	20.2 (3.7) ^E	21.9 (3.5)	24.8 (3.1)	28.4 (2.8)	32.3 (3.0)	36.1 (3.7)	38.5 (4.3)
	14-18	142	28.8 (1.5)	18.7 (2.2)	20.9 (2.0)	24.6 (1.8)	29.1 (2.1)	34.0 (2.8)	39.1 (3.9)	42.4 (4.7)
	19-30	111	27.9 (1.8)	17.8 (3.8) ^E	19.7 (3.5) ^E	23.2 (2.9)	27.7 (2.6)	32.7 (3.2)	37.9 (4.7)	41.2 (5.9)
	31-50	146	25.1 (1.3)	13.2 (2.7) ^E	15.1 (2.4)	18.5 (2.1)	22.8 (1.8)	28.1 (1.9)	33.4 (2.7)	36.9 (3.4)
	51-70	184	26.8 (1.3)	19.9 (3.2)	21.4 (2.8)	24.1 (2.1)	27.3 (1.7)	30.7 (2.3)	34.1 (3.6)	36.2 (4.7)
	>70	143	21.9 (1.7)	12.7 (3.1) ^E	14.5 (2.9) ^E	17.9 (2.5)	22.6 (2.3)	28.4 (3.0)	34.7 (4.6)	39.0 (6.0)
	19+	584	25.6 (0.8)	14.6 (1.3)	16.6 (1.2)	20.3 (1.1)	25.0 (1.1)	30.3 (1.2)	35.7 (1.6)	39.2 (2.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total monounsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 38.8 Total polyunsaturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	129	7.0	(0.5)	4.1	(0.9) ^E	4.6	(0.8) ^E	5.8	(0.7)	7.2	(0.6)	8.8	(0.8)	10.4	(1.1)	11.5	(1.4)
	4-8	213	11.5	(0.9)	6.5	(0.8)	7.3	(0.8)	8.8	(0.8)	10.8	(0.8)	13.4	(1.2)	16.2	(1.7)	18.1	(2.2)
Male																		
	9-13	122	18.2	(5.2) ^E	8.6	(2.4) ^E	9.7	(2.5) ^E	13.2	(2.8) ^E	16.5	(3.5) ^E	21.7	(5.0) ^E	26.7	(6.7) ^E	31.1	(8.4) ^E
	14-18	150	19.1	(1.4)	12.6	(1.2)	13.9	(1.3)	16.3	(1.5)	19.4	(1.9)	22.9	(2.4)	26.5	(3.1)	28.9	(3.6)
	19-30	106	18.9	(1.8)	8.4	(2.7) ^E	10.1	(2.6) ^E	13.4	(2.3) ^E	18.0	(2.1)	23.5	(2.9)	29.5	(4.7)	33.5	(6.5) ^E
	31-50	155	17.5	(1.6)	9.8	(2.1) ^E	11.2	(2.0) ^E	13.9	(1.9)	17.2	(2.0)	21.3	(2.5)	25.7	(3.5)	28.7	(4.3)
	51-70	122	15.5	(1.3)	8.9	(2.0) ^E	10.2	(1.8) ^E	12.7	(1.6)	16.0	(1.6)	20.0	(2.2)	24.4	(3.6)	27.6	(4.9) ^E
	>70	88	14.3	(1.1)	7.1	(1.2) ^E	8.2	(1.2)	10.4	(1.1)	13.5	(1.1)	17.3	(1.8)	21.6	(2.9)	24.6	(3.8)
	19+	471	16.9	(0.8)	8.0	(1.0)	9.6	(0.9)	12.5	(0.9)	16.3	(1.0)	21.1	(1.3)	27.0	(1.9)	31.4	(2.6)
Female																		
	9-13	103	11.4	(0.8)	8.7	(1.5) ^E	9.3	(1.4)	10.3	(1.2)	11.6	(1.1)	13.1	(1.2)	14.5	(1.5)	15.5	(1.8)
	14-18	142	12.1	(0.6)	7.5	(1.1)	8.4	(1.0)	10.0	(0.9)	12.1	(0.9)	14.5	(1.2)	17.1	(1.8)	18.8	(2.3)
	19-30	111	12.2	(0.7)	9.3	(1.8) ^E	9.9	(1.6) ^E	11.0	(1.3)	12.4	(1.1)	14.0	(1.6)	15.5	(2.6) ^E	16.6	(3.5) ^E
	31-50	146	10.8	(0.8)	5.5	(1.8) ^E	6.2	(1.6) ^E	7.7	(1.3) ^E	9.6	(1.1)	12.0	(1.1)	14.5	(1.7)	16.3	(2.3)
	51-70	184	12.8	(0.8)	6.9	(1.8) ^E	7.9	(1.6) ^E	9.8	(1.4)	12.4	(1.2)	15.6	(1.4)	19.2	(2.3)	21.7	(3.2)
	>70	143	10.5	(0.8)	6.9	(1.4) ^E	7.7	(1.3) ^E	9.1	(1.2)	10.9	(1.2)	13.1	(1.5)	15.5	(2.1)	17.0	(2.6)
	19+	584	11.6	(0.4)	6.4	(0.9)	7.2	(0.9)	8.9	(0.7)	11.1	(0.6)	13.8	(0.7)	16.7	(1.1)	18.7	(1.5)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total polyunsaturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 39.8 Total saturated fats (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean	(SE)	Percentiles (and SE) of usual intake													
					5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)							
Both																		
	1-3	129	21.4	(1.3)	10.4	(1.8) ^E	12.6	(1.7)	16.4	(1.6)	21.2	(1.6)	26.9	(2.0)	33.3	(2.9)	37.9	(3.7)
	4-8	213	24.3	(1.1)	16.2	(2.4)	17.7	(2.1)	20.6	(1.6)	24.3	(1.2)	28.4	(1.6)	32.5	(2.6)	35.3	(3.4)
Male																		
	9-13	122	29.5	(2.1)	21.2	(3.0)	22.8	(2.8)	25.7	(2.5)	29.5	(2.5)	34.0	(3.3)	38.4	(4.4)	41.2	(5.3)
	14-18	150	35.0	(2.1)	25.5	(4.0)	27.5	(3.6)	31.0	(3.0)	35.1	(2.6)	39.7	(3.3)	44.2	(4.8)	47.1	(6.1)
	19-30	106	33.3	(2.3)	18.8	(2.4)	21.5	(2.4)	26.5	(2.7)	33.0	(3.0)	40.4	(3.6)	47.8	(4.4)	52.6	(5.1)
	31-50	155	31.5	(2.6)	18.0	(3.6) ^E	20.3	(3.3)	24.8	(2.9)	31.0	(2.9)	38.5	(4.0)	46.5	(6.4)	52.2	(8.7) ^E
	51-70	122	27.3	(2.0)	15.2	(3.0) ^E	17.7	(2.7)	22.2	(2.4)	28.0	(2.6)	35.0	(3.7)	42.0	(5.5)	46.4	(6.8)
	>70	88	24.3	(1.9)	12.5	(2.4) ^E	14.5	(2.3)	18.4	(2.1)	23.6	(2.2)	29.6	(2.8)	36.0	(4.0)	40.2	(4.9)
	19+	471	30.0	(1.3)	14.3	(1.5)	17.0	(1.4)	22.1	(1.4)	29.3	(1.6)	38.1	(2.0)	47.3	(3.1)	53.8	(4.3)
Female																		
	9-13	103	26.6	(2.1)	19.3	(3.3) ^E	20.8	(3.0)	23.7	(2.8)	27.2	(2.8)	31.2	(3.3)	35.2	(4.2)	37.9	(5.0)
	14-18	142	22.6	(1.5)	12.0	(1.4)	14.1	(1.4)	17.7	(1.5)	22.1	(1.9)	27.4	(2.7)	33.3	(3.7)	37.6	(4.6)
	19-30	111	21.5	(1.6)	12.4	(1.2)	14.0	(1.3)	17.0	(1.6)	20.5	(2.0)	24.5	(2.4)	28.5	(2.9)	31.1	(3.1)
	31-50	146	21.7	(1.5)	12.6	(2.4) ^E	14.0	(2.2)	16.5	(1.9)	19.7	(1.8)	23.7	(2.0)	28.2	(2.9)	31.3	(3.8)
	51-70	184	21.0	(1.3)	13.3	(2.6) ^E	14.7	(2.3)	17.4	(1.9)	20.8	(1.6)	24.7	(2.3)	28.7	(3.6)	31.4	(4.7)
	>70	143	16.4	(1.4)	7.2	(2.2) ^E	8.6	(2.1) ^E	11.5	(2.0) ^E	15.6	(1.8)	20.8	(2.1)	26.8	(3.0)	31.0	(3.8)
	19+	584	20.6	(0.8)	10.8	(1.0)	12.5	(0.9)	15.7	(0.8)	19.6	(0.9)	24.3	(1.2)	29.5	(1.7)	33.0	(2.0)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total saturated fats.

For additional footnotes common to all tables, see Appendix A.

Table 40.8 Total sugars (g/d): Usual intakes from food, by DRI age–sex group, household population, Saskatchewan, 2004^{1,2}

Sex	Age (years)	n	Mean (SE)	Percentiles (and SE) of usual intake						
				5th (SE)	10th (SE)	25th (SE)	50th (SE)	75th (SE)	90th (SE)	95th (SE)
Both	1-3	129	99 (5)	52 (8)	61 (7)	77 (7)	96 (6)	119 (7)	143 (10)	159 (12)
	4-8	213	119 (6)	75 (5)	83 (5)	97 (6)	116 (6)	137 (7)	158 (8)	173 (9)
Male	9-13	122	145 (11)	89 (14)	100 (13)	119 (13)	145 (15)	174 (18)	204 (24)	225 (29)
	14-18	150	182 (18)	133 (24) ^E	143 (22)	161 (21)	183 (22)	206 (27)	230 (35)	245 (42) ^E
	19-30	106	137 (14)	50 (14) ^E	63 (15) ^E	89 (15) ^E	125 (17)	169 (22)	217 (31)	250 (39)
	31-50	155	134 (15)	60 (16) ^E	72 (15) ^E	96 (14)	128 (15)	167 (20)	214 (32)	249 (45) ^E
	51-70	122	97 (7)	44 (13) ^E	54 (12) ^E	73 (10)	97 (9)	125 (12)	153 (18)	171 (23)
	>70	88	95 (6)	50 (6)	58 (6)	72 (6)	91 (7)	114 (9)	142 (13)	163 (17)
	19+	471	120 (7)	46 (6)	58 (5)	82 (6)	114 (7)	156 (10)	207 (16)	246 (22)
Female	9-13	103	131 (10)	75 (10)	83 (10)	99 (11)	125 (13)	158 (17)	196 (24)	223 (30)
	14-18	142	145 (12)	98 (17) ^E	109 (17)	129 (17)	153 (18)	181 (22)	209 (28)	226 (33)
	19-30	111	115 (8)	54 (9) ^E	62 (8)	80 (7)	104 (7)	134 (12)	168 (20)	192 (28)
	31-50	146	95 (8)	47 (6)	54 (7)	69 (8)	88 (10)	110 (12)	135 (16)	153 (19)
	51-70	184	91 (9)	58 (13) ^E	65 (12) ^E	76 (11)	90 (10)	105 (11)	122 (15)	133 (18)
	>70	143	88 (4)	46 (5)	54 (5)	68 (4)	86 (5)	106 (6)	128 (8)	143 (11)
	19+	584	97 (4)	46 (5)	55 (5)	70 (5)	91 (5)	117 (6)	147 (9)	170 (12)

Data source: Statistics Canada, Canadian Community Health Survey, Cycle 2.2, Nutrition (2004) - Share File

Symbol Legend

^E Data with a coefficient of variation (CV) from 16.6% to 33.3%; interpret with caution.

<3 Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval entirely between 0 and 3%; interpret with caution.

^F Data with a coefficient of variation (CV) greater than 33.3% with a 95% confidence interval not entirely between 0 and 3%; suppressed due to extreme sampling variability.

Footnotes

¹ Intakes are based on food consumption only. For additional detail, see footnote 4 in Appendix A.

² No DRIs have been established for total sugars.

For additional footnotes common to all tables, see Appendix A.

Appendix A: Table Footnotes

The following footnotes apply to all of the summary data tables presented in Section II of this report.

1. The survey excludes from its target population those living in the three territories, individuals living on Indian reserves or on Crown lands, residents of institutions, full-time members of the Canadian Forces and residents of certain remote regions.
2. The tables exclude pregnant and breastfeeding females, subject to another set of nutritional recommendations. The sample of pregnant and breastfeeding females is not large enough to allow for reliable estimates.
3. Sample size and mean intake are based on the first 24-hour dietary recall (first day of interview) only.
4. Intakes are based on food consumption only. Intakes from vitamin and mineral supplements are not included. Inferences about the prevalence of nutrient excess or inadequacy based on intakes from food alone may respectively underestimate or overestimate the prevalences based on total nutrient intakes from both food and supplements.
5. The intake distribution (percentiles and percentage above or below a cut-off when applicable) was adjusted to remove within-individual variability using Software for Intake Distribution Estimation (SIDE) (Iowa State University, 1996) and the method presented in Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440-1449.
6. In some cases, within-individual variance was estimated at the regional or national level and applied at the provincial level. For more details, see Section II.4: Measuring Sampling Variability with Bootstrap Replication in Volume 1 of the *Nutrient Intakes from Food* report series.
7. Bootstrapping techniques were used to produce the coefficient of variation (CV) and the standard error (SE).
8. AMDR is the Acceptable Macronutrient Distribution Range, expressed as a percentage of total energy intake. Intakes inside the range (shown in the AMDR columns) are associated with a reduced risk of chronic disease while providing adequate intakes of essential nutrients. For further information on AMDR see the Health Canada publication *Canadian Community Health Survey*,

Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data, Section 2.1.5, p. 27.

The applications of the AMDRs for essential fatty acids to group assessment are not the same as for the other macronutrients. The lower boundaries for the AMDR for linoleic and alpha-linolenic acids are not based on the same type of endpoints as the boundaries for total fat and carbohydrate. The boundaries for fat and carbohydrate are set based on evidence indicating increased risk for coronary heart diseases and the lower bound of the AMDR for both n-6 (linoleic) and n-3 (alpha-linolenic) fatty acids is based on the percent of energy from these fatty acids needed to provide the AI for these nutrients. The AI, in turn, is based on the median intake of both linoleic and alpha-linolenic acid in the United States, where essential fatty acid deficiency is non-existent in the healthy population.

Thus, by definition about half the population has intakes of these fatty acids below the AI and therefore outside the AMDR. In other words, based on the AI, one would conclude that the population is "adequate" with respect to linoleic and alpha-linolenic acids, while based on the AMDR a different conclusion (i.e. that 50% of the population has intakes below the AMDR) would be reached. Therefore, the lower bound of the AMDRs for linoleic and alpha-linolenic acids should not be used in the assessment of population intakes.

9. EAR is the Estimated Average Requirement. The level of intake at the EAR (shown in the EAR columns) is the average daily intake level that is estimated to meet the requirement, as defined by the specified indicator of adequacy, in half of the apparently healthy individuals in a DRI age–sex group. For further information on EAR see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.1, p. 23.
10. AI is the Adequate Intake. The level of intake at the AI (shown in the AI columns) is the recommended average daily intake level based on observed or experimentally determined approximations or estimates of nutrient intake by a group or groups of apparently healthy people that are assumed to be adequate. It is developed when an EAR cannot be determined. The percentage of the population having a usual intake above the AI (shown in the %>AI columns) almost certainly meets their needs. The adequacy of intakes below the AI cannot be assessed, and should not be interpreted as being inadequate. For further information on AI see the Health Canada

publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.3, p. 25.

11. UL is the Tolerable Upper Intake Level. The level of intake at the UL (shown in the UL columns) is the highest average daily intake level that is likely to pose no risk of adverse health effects to almost all individuals in the general population. For further information on UL see the Health Canada publication *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*, Section 2.1.4, p. 26.
12. For a more detailed understanding of DRIs and their interpretation when assessing intakes of particular nutrients, consult the summary of the series of publications on DRIs published by the Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*.
13. In terms of precision, the estimate 0.0 with a standard error of 0.0 refers to a standard error smaller than 0.1%.

Appendix B: Justification for Excluding Nutrients from Volume 2 and Volume 3

Volume 1 of the compendium contained data on 13 nutrients, including 6 nutrients expressed as a percent of total energy. There were originally 31 nutrients scheduled to be released in Volumes 2 and 3 of the compendium, but for a variety of reasons some of these nutrients are not included. Decisions to omit these nutrients were made jointly by representatives from Statistics Canada and Health Canada.

Exclusions and changes to the list of nutrients that were to be included in Volumes 2 and 3 of the compendium are as follows:

Total milligrams of folic acid

Folic acid is found in small amounts in a number of foods. Most respondents consumed a small amount of folic acid, which resulted in a bimodal distribution of folic acid intake. As a result, it was very difficult to normalize the distribution, which meant that SIDE was unable to calculate usual intake.

One of the steps that SIDE uses to estimate usual intake is to transform the data into a normal distribution. Assessing SIDE's ability to perform this transformation rests on measuring the Anderson-Darling (A-D) score for normality. The A-D score is a statistic that measures how close a distribution is to a normal distribution. Any A-D score less than 0.576 is considered to be sufficiently normal for SIDE to continue without warning. Typically, SIDE will be able to transform 95% of the domains without error using the default SIDE options. The remaining 5% of domains will typically score higher than 0.576 but usually less than 1.0. Adjusting the SIDE options will usually reduce the A-D to within the limit. In the case of folic acid, more than half of the provincial domains had an A-D score above the 0.576 threshold and many domains scored higher than 2. The nature of the data simply does not allow SIDE to produce proper estimates for the usual intake of folic acid.

Total grams of alcohol

Alcohol is consumed differently than other nutrients. For most respondents, alcohol is not part of their daily intake of food, but rather is something that is consumed occasionally. In this sense, in terms of analysis, alcohol behaves more like a food than a nutrient. In order for SIDE to estimate the usual intake of foods, many recalls are needed to capture enough occurrences of the particular food. Thus, two recalls are not enough to calculate the usual intake of alcohol.

Percent of energy from alcohol

The difficulty in estimating a usual intake for alcohol causes similar problems for expressing that intake as a percent of total energy.

Caffeine

Caffeine also is consumed differently than other nutrients. The usual intake of caffeine could not be calculated due to the same issues as folic acid and alcohol. Many respondents reported zero or small levels of caffeine intake. Therefore, it is difficult for SIDE to properly model the data with only two dietary recalls.

Based on the changes above, the list of nutrients included in Volume 1 and the revised list of nutrients included in Volumes 2 and 3 are as follows:

List of Nutrients Included in the Three-Volume Set		
Volume 1	Volume 2	Volume 3
Total Energy	Folate (DFE)	Folacin
Percentage of total energy intake from fats	Iron	Linolenic acid (g, % energy)
Percentage of total energy intake from protein	Linoleic acid (g, % energy)	Moisture
Percentage of total energy intake from carbohydrates	Magnesium	Naturally occurring folate
Percentage of total energy intake from saturated fats	Niacin	Protein
Percentage of total energy intake from monounsaturated fats	Phosphorus	Total carbohydrates
Percentage of total energy intake from polyunsaturated fats	Potassium	Total fats
Total dietary fibre	Riboflavin	Total monounsaturated fats
Cholesterol	Thiamin	Total polyunsaturated fats
Vitamin A	Vitamin B ₆	Total saturated fats
Vitamin C	Vitamin B ₁₂	Total sugars
Calcium	Vitamin C by smoking status	
Sodium	Vitamin D	
	Zinc	

Appendix C: References

Department of Statistics and Center for Agricultural and Rural Development, Iowa State University: *A User's Guide to SIDE, Software for Intake Distribution Estimation Version 1.0*. Technical Report 96-TR 30. Ames, IA: Iowa State University Statistical Laboratory, 1996. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr30.pdf

Dodd KW: *A Technical Guide to C-SIDE, Software for Intake Distribution Estimation*. Technical Report 96-TR 32, Dietary Assessment Research Series Report 9. Ames, IA: Department of Statistics and Center for Agricultural and Rural Development, Iowa State University, 2006. Available at:

www.card.iastate.edu/publications/DBS/PDFFiles/96tr32.pdf

Health Canada: *Canadian Community Health Survey, Cycle 2.2, Nutrition (2004)—A Guide to Accessing and Interpreting the Data*. Ottawa: Publications, Health Canada, 2006. Available at: www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_focus-volet_escc-eng.php

Institute of Medicine: *Dietary Reference Intakes: A Risk Assessment Model for Establishing Upper Intake Levels for Nutrients*. Washington, DC: National Academy Press, 1998. Available at: <http://books.nap.edu/catalog/6432.html>

Institute of Medicine: *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin and Choline*. Washington, DC: National Academy Press, 1998. Available at: www.nap.edu/catalog.php?record_id=6015

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academy Press, 2000. Available at: <http://books.nap.edu/catalog/9956.html>

Institute of Medicine: *Dietary Reference Intakes: Applications in Dietary Planning*. Washington, DC: National Academy Press, 2003. Available at: <http://books.nap.edu/catalog/10609.html>

Institute of Medicine: *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington, DC: National Academy Press, 2004. Available at: <http://books.nap.edu/catalog/10925.html>

Institute of Medicine: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. Washington, DC:

National Academy Press, 2005. Available at:

<http://books.nap.edu/catalog/10490.html>

Institute of Medicine: *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: National Academy Press, 2006.

Available at: www.nap.edu/topics.php?topic=380

Nusser SM, Carriquiry AL, Dodd KW, Fuller WA: A semiparametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* 1996; 91: 1440–1449

Nusser SM, Fuller WA, Guenther PM: Estimating usual dietary intake distributions: adjusting for measurement error and nonnormality in 24-hour food intake data. In L Lyberg, P Biemer, M Collins et al., eds. *Survey Measurement and Process Quality*, New York: John Wiley & Sons, 1997.

Statistics Canada: *Canadian Community Health Survey (CCHS) Cycle 2.2—Nutrition: Software for Intake Distribution Estimation (SIDE) Documentation*. Ottawa, 2007.

Available at: www.statcan.ca/english/sdds/document/5049_D22_T9_V1_E.pdf