## Residential Telephone Service Survey

# Microdata User Guide for the re-weighted May 1999 data

Special Surveys Division issued January 2001

# Table of Contents

1.0	Note to	o users 1
2.0	Introd	luction 3
3.0	Backg	ground 5
4.0	Objec	tives 7
5.0	Conce	epts and Definitions 9
6.0	6.1 6.2 6.3 6.4 6.5 6.6	y Methodology         11           Population Coverage         11           Sample Design         11           6.2.1 Primary Stratification         12           6.2.2 Types of Areas         12           6.2.3 Secondary Stratification         13           6.2.4 Cluster Delineation and Selection         13           6.2.5 Dwelling Selection         14           6.2.6 Person Selection         14           Sample Size         17           Sample Rotation         17           Modifications to the L.F.S design for the Supplement         18           Sample size by Province for the Supplement         18
7.0	Data (7.1 7.2 7.3 7.4 7.5	Interviewing for the LFS
8.0		Service Survey       21         Processing       23         Data Capture       23         Editing       23         Coding of Open-ended Questions       23         Creation of Derived Variables       23         Weighting       24         Suppression of Confidential Information       24
9.0	<b>Data (</b> 9.1 9.2	Quality         27           Response Rates         27           Survey Errors         28

14.0	Recor	Layout and Univariates 75	5
13.0	Quest	onnaires and Code Sheets	1
	12.2	Weighting Procedures for the Residential Telephone Service Survey	
12.0	Weigh	ing6 Weighting Procedures for the LFS6	<b>7</b> 7
	11.6	CV Tables 53	3
	11.5	Release cut-off's for the RTSS 5	1
	11.4	Coefficients of Variation for Quantitative Estimates 50	
		11.3.1 Example of using the CV tables to do a t-test 50	
	11.3	How to use the CV tables to do a t-test	9
		confidence limits	9
	11.2	11.2.1 Example of using the CV tables to obtain	J
	11.2	Categorical Estimates	
		11.1.1 Examples of using the C.V. tables for	1
	11.1	How to use the C.V. tables for Categorical Estimates 42	2
11.0		imate Sampling Variability Tables 4	
			_
	10.4	CV Release Guidelines	
	10.3	Guidelines for Statistical Analysis	
		10.2.3 Tabulation of Quantitative Estimates 30	
		10.2.2 Tabulation of Categorical Estimates	
		10.2.1 Definitions of types of estimates: Categorical vs.  Quantitative	5
	10.2	Sample Weighting Guidelines for Tabulation 34	4
	10.1	Rounding Guidelines	
10.0		nes for Tabulation, Analysis and Release	
		·	
		9.2.4 Non-response	
		9.2.3 Data Processing	
		9.2.1 The Frame	
		9.2.1 The Frame	Ω

## 1.0 Note to users

The Residential Telephone Service Survey (RTSS) is administered to a sub-sample of the dwellings in the Labour Force Survey (LFS) sample, and therefore its weighting and population estimates are closely tied to those of the LFS.

In the past the LFS calculated its population totals based on population projections using the 1991 Census. In January 2000, the Labour Force Survey began calculating its population totals based on population projections using the 1996 Census. This has resulted in a decrease in the estimate of the total population. For RTSS, this resulted in population totals reported in the May 2000 RTSS to be lower than those reported in the August 1999 RTSS.

Because of this, users were not able to make comparisons between the <u>counts</u> seen in the May 2000 RTSS and any previous RTSS. Users could, however, continue to compare <u>rates</u> as rates are robust to changes in totals.

To address this issue, the weights on the May 1999 RTSS microdata file were re-calculated in October 2000, using the 1996 Census population projections. This means that users can now compare the May 2000 data to the re-weighted May 1999 data.

This Microdata User Guide should be used with the re-weighted 1999 May RTSS data file.

# 2.0 Introduction

The latest quarter of the Residential Telephone Service Survey was conducted by Statistics Canada in May 1999 with the cooperation and support of Bell Canada. This manual has been produced to facilitate the manipulation of the microdata file of the survey results.

Any questions about the data set or its use should be directed to:

Statistics Canada
Tracey Bushnik
Special Surveys Division, Statistics Canada
Section B38
1st floor, Jean Talon Building
Tunney's Pasture
Ottawa, Ontario K1A 0T6
(613) 951-3017

Bell Canada Lynn Solvason Regulatory Matters Floor 5 - 105 Hotel de Ville Hull, Quebec (819) 773-5582

## 3.0 Background

Statistics Canada was approached by Stentor Resource Centre Inc. to conduct a quarterly survey in order to monitor the phone penetration rates across Canada. The management of the survey was transfered from Stentor to Bell Canada in the Fall 1998. The latest quarter was conducted in May 1999 as a supplement to the Labour Force Survey. Bell Canada and other companies are from time to time negotiating local service pricing options for phone rates with the Canadian Radio-Television and Telecommunication Commission. Penetration rates are the most reliable indicator of affordability as there is no price range that can be identified as affordable or not affordable. As a result, the importance of monitoring any changes in phone penetration rates and analysing the reasons for non-subscribers is necessary to properly guide regulators in decisions about rate increases, decreases or subsidies.

Concern has been expressed that the current mechanism for monitoring penetration rates is not adequate in providing timely results to indicate whether Canadian penetration rates fall as a result of increases in local rates. At present, data on penetration rates are available from the Household Facilities and Equipment Survey (HFE) but only on an annual basis. Given the changes that are and will be occurring in the basic residential telephone rates, an annual survey was not adequate to accurately reflect the impact that these changes are having on Canadian telephone subscribership.

## 4.0 Objectives

There are two main objectives which Bell Canada has outlined. They are:

- to collect information on penetration rates across
   Canada (and make them available by province);
- (ii) to collect information on non-subscriber characteristics.

To accommodate these goals, and to ensure that the survey is focused on fulfilling these objectives, Bell Canada submitted an analysis plan which outlined their data needs. This plan was used to design the questionnaire and to justify the variables requested.

# 5.0 Concepts and Definitions

This chapter outlines concepts and definitions of interest to the users. Users are referred to Chapter 12 of this document for a copy of the actual survey questions used.

#### Number of telephone numbers for the residence:

Includes cellular telephone numbers and telephone numbers used for business even if the business is not within the residence or if the employer is paying for the persons's telephone service within that person's household. It includes cellular telephones from work that are brought home. Pagers are excluded.

#### Income:

Household income has been measured against the 1992-base Statistics Canada Low Income Cut-Offs - LICO (see catalogue 13-551-XPB for the full definition of LICO). For the purpose of the RTSS, the low income values used to assessed the level of income were rounded to the nearest \$500. The total income was collected for the entire household, regardless of family structure. LICOs normally apply to economic families and unattached individuals. Respondents were asked to self-report if their total household income was above or below the modified LICO, with no additional prompting for precision

Although Statistics Canada's LICOs are often referred to as poverty lines, they do not have an officially recognized status, nor does Statistics Canada promote their use as poverty lines. Since the LICOs are recognized Statistics Canada income measures and that many modifications were made to them for the purpose of the RTSS, we recommend that the term 'LICO' not be used to refer to the RTSS income measure as this could be misleading to unadvised readers.

## 6.0 Survey Methodology

The latest quarter of the Residential Telephone Service Survey was administered in May 1999 to a sub-sample of the dwellings in the Labour Force Survey (LFS) sample, and therefore its sample design is closely tied to that of the LFS. The LFS design is briefly described in Sections 5.1 to 5.4<sup>1</sup>. Sections 5.5 and 5.6 describe how the Residential Telephone Service Survey departed from the basic LFS design in May 1999.

## 6.1

## **Population Coverage**

The LFS is a monthly household survey whose sample of individuals is representative of the civilian, non-institutionalized population 15 years of age or older in Canada's ten provinces. Specifically excluded from the survey's coverage are residents of the Yukon<sup>2</sup> and Northwest Territories, persons living on Indian Reserves, full-time members of the Canadian Armed Forces and inmates of institutions. These groups together represent an exclusion of approximately 2% of the population aged 15 or over.

#### 6.2

## Sample Design

The LFS has undergone an extensive redesign, culminating in the introduction of a new design at the end of 1994. The LFS sample is based upon a stratified, multi-stage design employing probability sampling at all stages of the design. The design principles are the same for each province. A diagram summarizing the design stages appears at the end of this section.

A detailed description of the previous LFS design is available in the Statistics Canada publication entitled **Methodology of the Canadian Labour Force Survey**, 1984-1990 (catalogue #71-526).

Since 1992, the LFS has been administered in the Yukon, using an alternative methodology that accommodates some of the operational difficulties inherent to remote locales. To improve reliability due to small sample size, estimates are available on a three month average basis only. These estimates are not included in national totals.

#### 6.2.1

## **Primary Stratification**

Provinces are divided into economic regions and employment insurance regions. Economic regions (ERs) are geographic areas of more or less homogeneous economic structure formed on the basis of federal provincial agreements. They are relatively stable over time. Employment insurance economic regions (EIERs) are also geographic areas, and are roughly the same size and number as ERs, but they do not share the same definitions. Labour force estimates are produced for the EIER regions for the use of Human Resources Development Canada.

The intersections of the two types of regions form the first level of stratification for the LFS. These ER/EIER intersections are treated as primary strata and further stratification is carried out within them (see section 5.2.3). Note that a third set of regions, Census Metropolitan Areas (CMAs), is also respected by stratification in the current LFS design, since each CMA is also an EIER.

### 6.2.2

## **Types of Areas**

The primary strata (ER/EIER intersections) are further disaggregated into 3 types of areas: rural, urban, and remote areas. Urban and rural areas are loosely based on the Census definitions of urban and rural, with some exceptions to allow for the formation of strata in some areas. Urban areas include the largest CMAs down to the smallest villages categorized by the 1991 Census as urban (1000 people or more), while rural areas are made up of areas not designated as urban or remote.

All urban areas are further subdivided into two types: those using an apartment list frame and an area frame, as well as those using only an area frame.

Approximately 1% of the LFS population is found in remote areas of provinces which are less accessible to LFS interviewers than other areas. For administrative purposes, this portion of the population is sampled separately through the remote area frame. Some populations, not congregated in places of 25 or more people, are excluded from the sampling frame.

#### 6.2.3

## Secondary Stratification

In urban areas with sufficiently large numbers of apartment buildings, the strata are subdivided into apartment frames and area frames. The apartment list frame is a register which is based upon information supplied by CMHC and is maintained in the 18 largest cities across Canada. The purpose of this is to ensure better representation of apartment dwellers in the sample as well as to minimize the effect of growth in clusters, due to construction of new apartment buildings. In the major cities, the apartment strata are further stratified into low income strata and regular strata.

Where it is possible and/or necessary, the urban area frame is further stratified into regular strata, high income strata, and low population density strata. Most urban areas fall into the regular urban strata, which, in fact, cover the majority of Canada's population. High income strata are found in major urban areas, while low density urban strata consist of small towns that are geographically scattered.

In rural areas, the population density can vary greatly from relatively high population density areas to low population density areas, resulting in the formation of strata that reflect these variations. The different stratification strategies for rural areas were based not only on concentration of population, but also on cost-efficiency and interviewer constraints.

In each province, remote settlements are sampled proportional to the number of dwellings in the settlement, with no further stratification taking place. Dwellings are selected using systematic sampling in each of the places sampled.

### 6.2.4

## Cluster Delineation and Selection

Households in final strata are not selected directly. Instead, each stratum is divided into clusters, and then a sample of clusters is selected within the stratum. Dwellings are then sampled from selected clusters. Different methods are used to define the clusters, depending on the type of stratum.

Within each urban stratum in the urban area frame, a number of geographically contiguous groups of dwellings, or clusters, are formed based upon 1991 Census counts. These clusters are generally a set of one or more city blocks or block faces. The selection of a sample of clusters (always 6 or a multiple of 6 clusters) from each of these secondary strata represents the first stage of sampling in most urban areas. In some other urban areas, Census Enumeration Areas (EAs) are used as clusters. In the

low density urban strata, a three stage design is followed. Under this design, two towns within a stratum are sampled, and then six or 24 clusters within each town are sampled.

For urban apartment strata, instead of defining clusters, the apartment building is the primary sampling unit. Apartment buildings are sampled from the list frame with probability proportional to the number of units in each building.

Within each of the secondary strata in rural areas, where necessary, further stratification is carried out in order to reflect the differences among a number of socio-economic characteristics within each stratum. Within each rural stratum, six EAs or two or three groups of EAs are sampled as clusters.

#### 6.2.5

## **Dwelling Selection**


In all three types of areas (urban, rural and remote areas) selected clusters are first visited by enumerators in the field and a listing of all private dwellings in the cluster is prepared. From the listing, a sample of dwellings is then selected. The sample yield depends on the type of stratum. For example, in the urban area frame, sample yields are either 6 or 8 dwellings, depending on the size of the city. In the urban apartment frame, each cluster yields 5 dwellings, while in the rural areas and EA parts of cities, each cluster yields 10 dwellings. In all clusters, dwellings are sampled systematically. This represents the final stage of sampling.

### 6.2.6

## **Person Selection**

Demographic information is obtained for all persons for whom the selected dwelling is the usual place of residence. LFS information is obtained for all civilian household members 15 years of age or older. Response burden is minimized for the elderly (70 years of age or older) by carrying forward their responses for the initial interview to the subsequent five months in the survey.

#### Labour Force Survey Sample Design - 1995+



= level of stratification

EIR - Employment Insurance Region

ER - Economic Region

{%} - percentage of total sample

EA - Census Enumeration Area cluster - set of block faces

= stage of sampling

## 6.3 Sample Size

The sample size of eligible persons in the LFS is determined so as to meet the statistical precision requirements for various labour force characteristics at the provincial and sub-provincial level, and to meet the requirements of federal, provincial and municipal governments as well as a host of other data users.

The monthly LFS sample consists of approximately 59,000 dwellings. After excluding dwellings found to be vacant, dwellings demolished or converted to non-residential uses, dwellings containing only ineligible persons, dwellings under construction, and seasonal dwellings, about 52,350 dwellings remain which are occupied by one or more eligible persons. From these dwellings, LFS information is obtained for approximately 102,000 civilians aged 15 or over.

## 6.4 Sample Rotation

The LFS employs a panel design whereby the entire monthly sample of dwellings consists of 6 panels, or rotation groups, of approximately equal size. Each of these panels is, by itself, representative of the entire LFS population. All dwellings in a rotation group remain in the LFS sample for 6 consecutive months after which time they are replaced (rotated out of the sample) by a new panel of dwellings selected from the same or similar clusters.

This rotation pattern was adopted to minimize any problems of nonresponse or respondent burden that would occur if households were to remain in the sample for longer than 6 months. It also has the statistical advantage of providing a common sample base for short-term month-tomonth comparisons of LFS characteristics, since five of the six rotation groups in the LFS sample are common from month to month.

Because of the rotation group feature, it is possible to readily conduct supplementary surveys using the LFS design but employing less than the full size sample.

#### 6.5

## Modifications to the L.F.S design for the Supplement

The Residential Telephone Service Survey used five of the six rotation groups in the May 1999 LFS sample. For the RTSS, the coverage of the LFS was set at the household level. Unlike the LFS where information is collected for all eligible household members, the Residential Telephone Service Survey only collected information from one household member who reported the information at the household level.

#### 6.6

## Sample size by Province for the Supplement

The following table shows the number of household in the LFS sampled rotations who were eligible for the Residential Telephone Service Survey supplement.

PROVINCE	SAMPLE SIZE

Newfoundland	1586
Prince Edward Island	1146
Nova Scotia	2882
New Brunswick	2446
Quebec	8374
Ontario	12615
Manitoba	3166
Saskatchewan	3283
Alberta	3291
British Columbia	3972
CANADA	42761

## 7.0 Data Collection

Data collection for the LFS is carried out each month using the computerassisted method during the week following the LFS reference week, usually the third week of the month.

#### 7.1

## Interviewing for the LFS

Statistics Canada interviewers, who are part-time employees hired and trained specifically to carry out the LFS, contact each of the sampled dwellings to obtain the required labour force information. Each interviewer contacts approximately 70 dwellings per month.

Dwellings new to the sample are contacted through a personal visit. The interviewer first obtains socio-demographic information for each household member and then obtains labour force information for all eligible members. All interviews are conducted using a notebook computer. Provided there is a telephone in the dwelling and permission has been granted, subsequent interviews are conducted by telephone. As a result, approximately 85% of all dwellings are interviewed by telephone. In these subsequent monthly interviews, as they are called, the interviewer confirms the sociodemographic information collected in the first month and collects the labour force information for the current month.

In all dwellings, information about all household members is obtained from a knowledgeable household member - usually the person at home when the interviewer calls. Such 'proxy' reporting, which accounts for approximately 55% of the information collected, is used to avoid the high cost and extended time requirements that would be involved in repeat visits or calls necessary to obtain information directly from each respondent.

At the conclusion of the LFS monthly interviews, interviewers introduce the supplementary survey, if any, to be administered to some or all household members that month.

If, during the course of the six months that a dwelling normally remains in the sample, an entire household moves out and is replaced by a new household, information is obtained about the new household for the remainder of the six-month period.

#### 7.2

## **Supervision and Control**

All LFS interviewers are under the supervision of a staff of senior interviewers who are responsible for ensuring that interviewers are familiar with the concepts and procedures of the LFS and its many supplementary surveys, and also for periodically monitoring their interviewers and reviewing their completed documents. The senior interviewers are, in turn, under the supervision of the LFS program managers, located in each of the six Statistics Canada regional offices.

#### 7.3

## Non-Response to the LFS

Interviewers are instructed to make all reasonable attempts to obtain LFS interviews with members of eligible households. For individuals who at first refuse to participate in the LFS, a letter is sent from the Regional Office to the dwelling address stressing the importance of the survey and the household's cooperation. This is followed by a second call (or visit) from the interviewer. For cases in which the timing of the interviewer's call (or visit) is inconvenient, an appointment is arranged to call back at a more convenient time. For cases in which there is no one home, numerous call backs are made. Under no circumstances are sampled dwellings replaced by other dwellings for reasons of non-response.

Each month, after all attempts to obtain interviews have been made, a small number of non-responding households remain. For households non-responding to the LFS and for which LFS information was obtained in the previous month, this information is brought forward and used as the current month's LFS information. No supplementary survey information is collected for these households.

### 7.4

## Data Collection Modifications for Residential Telephone Service Survey

Information for the Residential Telephone Service Survey was obtained from a knowledgable household member. Upon completion of the Labour Force Survey interview, the interviewer introduced the RTSS and proceeded with the interview with the respondent's permission.

The Residential Telephone Service Survey was programmed to appear on the list of surveys to be completed on the notebook computer after the demographic component for the LFS had been completed. Any RTSS component not completed at the time the LFS was transmitted to one of the Statistics Canada regional offices was left incomplete and transmitted with the LFS.

#### 7.5

## Non-Response to the Residential Telephone Service Survey

For households responding to the LFS, the next stage of data collection was to administer the Residential Telephone Service Survey. In total, 42,761 households were eligible for the supplementary survey; the Residential Telephone Service Survey interview was completed for 40,340 of these households for a response rate of 94.3%. Most non-response cases for the supplementary survey in May 1999 are directly linked to the computer-assisted collection method. Some questionnaires for RTSS were not created because the application was not properly installed on some laptop computers or they were not completed before the LFS questionnaires from which they are dependent were transmitted. More detailed information on response rates is presented in Chapter 8 (Data Quality).

# 8.0 Data Processing

The main output of the Residential Telephone Service Survey is a "clean" microdata file. This section presents a brief summary of the processing steps involved in producing this file.

#### 8.1

## **Data Capture**

Capture of survey data was done directly on notebook computers by interviewers at the time of collection. A partly edited version of the computer record was electronically transmitted to Ottawa for further processing. In total, 49,325 interviews were captured and transmitted for the survey.

#### 8.2

## **Editing**

The type of error treated involved a lack of information in questions which should have been answered. For this type of error, a non-response or "not-stated" code was assigned to the item.

### 8.3

## **Coding of Open-ended Questions**

No data items on the questionnaire were recorded by interviewers in an open-ended format. A total of two partially open-ended questions were included in the survey. These were items relating to reasons households do not have telephone service for their residence and why they cancelled their telephone service.

### 8.4

## **Creation of Derived Variables**

A number of data items on the microdata file have been derived by combining items on the questionnaire in order to facilitate data analysis. CMA, for example, is actually a combination of Census Metropolitan Area (CMA) and Census Agglomeration(CA). The CAs have been recoded to 0, while the CMAs remain the same. A 'size of urban area' variable was also

created. This variable provides a population size code based on 1991 Census definitions for every urban/non-urban area in the LFS sample frame.

## 8.5 Weighting

The principle behind estimation in a probability sample such as the LFS is that each person in the sample "represents", besides himself or herself, several other persons not in the sample. For example, in a simple random 2% sample of the population, each person in the sample represents 50 persons in the population. The same principle also applies to households.

The weighting phase is a step which calculates, for each record, what this number is. This weight appears on the microdata file, and must be used to derive meaningful estimates from the survey. For example, if the number of households with one or more telephone numbers for their residence is to be estimated, it is done by selecting the records referring to those households in the sample with that characteristic and summing the weights entered on those records.

Details of the method used to calculate these weights are presented in Chapter 11.

# 8.6 Suppression of Confidential Information

It should be noted that the 'Public Use' microdata files described above differ in a number of important respects from the survey 'master' files held by Statistics Canada. These differences are the result of actions taken to protect the anonymity of individual survey respondents. Users requiring access to information excluded from the microdata files may purchase custom tabulations. Estimates generated will be released to the user, subject to meeting the guidelines for analysis and release outlined in Section 9 of this document.

#### **Province** - Suppression of Geographic Identifiers

The survey master data file includes explicit geographic identifiers for province, stratum and Census Metropolitan Area. It is also possible to obtain, where sample sizes permit, estimates by urban size class. The survey public-use microdata files usually do not contain any geographic identifiers below the provincial level. However, since the RTSS is a household based survey, the variables CMA and urban size class will be on

the microdata file. Note that, in some provinces, urban size class values may be suppressed for certain unique areas where disclosure risk may be high.

None of the questions in the RTSS survey were completely open-ended. The two questions that were partially open-ended (Q02 and Q03) were regrouped in the CAI application and therefore appear as such on the master and microdata files.

## 9.0 Data Quality

### 9.1

## **Response Rates**

The following table summarizes the response rates to the Labour Force Survey and to the Residential Telephone Service Survey in May 1999.

	Household response rate for full LFS (05 99) (*1)	Household response rate for LFS rotations (1, 2, 3, 4, 6) (*1)	Household response rate to Residential Telephone Service Survey (*2)
Newfoundland	95.0%	95.1%	96.0%
Prince Edward Island	96.5%	97.0%	96.3%
Nova Scotia	94.3%	94.7%	95.2%
New Brunswick	94.3%	95.0%	91.4%
Quebec	94.7%	95.3%	94.4%
Ontario	95.8%	96.5%	94.5%
Manitoba	96.3%	96.6%	94.3%
Saskatchewan	96.4%	96.8%	94.2%
Alberta	96.8%	96.9%	94.3%
British Columbia	95.0%	95.6%	93.9%
CANADA	95.5%	96.0%	94.3%

#### Note:

- (\*1) Response rate is number of responding households as a percentage of number of eligible households.
- (\*2) Response rate is number of households responding to the Residential Telephone Service Survey as a percentage of number of households responding to LFS in rotations sampled.

#### 9.2

## **Survey Errors**

The estimates derived from this survey are based on a sample of households. Somewhat different figures might have been obtained if a complete census had been taken using the same questionnaire, interviewers, supervisors, processing methods, etc. as those actually used. The difference between the estimates obtained from the sample and the results from a complete count taken under similar conditions is called the sampling error of the estimate.

Errors which are not related to sampling may occur at almost every phase of a survey operation. Interviewers may misunderstand instructions, respondents may make errors in answering questions, the answers may be incorrectly entered on the questionnaire and errors may be introduced in the processing and tabulation of the data. These are all examples of non-sampling errors.

### 9.2.1

### The Frame

Because the RTSS was a supplement to the LFS, the frame used was the LFS frame. Any non-response to the LFS had an impact on the RTSS frame. Because non-response to the LFS is quite low (usually less than 5%) this impact was minimal. The quality of the sampling variables in the frame was very high. The RTSS sample consisted of five rotation groups from the LFS. No records were dropped due to missing rotation group number or any other type of sampling variable.

Note that the LFS frame excludes about 2% of all households in the 10 provinces of Canada. Therefore, the RTSS frame also excludes the same proportion of households in the same geographical area. It is likely that this exclusion introduces little, if any, significant bias into the survey data.

All variables in the LFS frame are updated monthly.

Some variables on the sampling frame play a critical role with respect to software application used in the survey. For example, in the RTSS CAI application, each record must have accurate stratum, cluster and rotation group codes. These variables are always of very high quality each month in the LFS.

At times, duplication of records occurs. 54 such records were dropped from the RTSS data file for the May 1999 collection. Some duplicates are created at the time of collection but most come from double transmissions of data from the collection environment to the processing server.

#### 9.2.2

#### **Data Collection**

Interviewer training consisted of reading the RTSS Procedures Manual, practising with the RTSS training cases on the laptop computer, and discussing any questions with senior interviewers before the start of the survey. A description of the background and objectives of the survey was provided, as well as a glossary of terms and a set of questions and answers. Interviewers collected RTSS information at the same time that LFS information was collected. The collection period ran from May 17 - 26, 1999.

#### 9.2.3

## **Data Processing**

During processing of the data, 6,869 RTSS records did not match to corresponding records in the LFS. Thus they were coded as out-of-scope and were dropped from further processing. When supplementary survey records do not match to host survey records they must be dropped since a weight cannot be derived for them.

Conversely, 311 records in the LFS were found that should have matched to an RTSS record but did not. These records were coded as in-scope, since they were eligible records from the frame which, for one reason or another, did not have corresponding RTSS records. These records were considered to be non-responding records, and were used in the weighting process to adjust for non-response.

Data processing of the RTSS was straightforward since there were only five questions on the CAI application. Any record that contained a refusal or don't know in the first question (Q01) was coded as a non-response. Note that 291 records were treated this way. Since the data was collected using a CAI instrument, data quality before processing was very high. Very few changes were made to the data during editing.

No imputation was done for this survey.

## 9.2.4

## Non-response

Over a large number of observations, randomly occurring errors will have little effect on estimates derived from the survey. However, errors occurring systematically will contribute to biases in the survey estimates. Considerable time and effort was made to reduce non-sampling errors in the survey. Quality assurance measures were implemented at each step of the data collection and processing cycle to monitor the quality of the data. These measures included the use of highly skilled interviewers, extensive training of interviewers with respect to the survey procedures and questionnaire, observation of interviewers to detect problems of questionnaire design or misunderstanding of instructions, procedures to ensure that data capture errors were minimized and coding and edit quality checks to verify the processing logic.

A major source of non-sampling errors in surveys is the effect of <u>non-response</u> on the survey results. The extent of non-response varies from partial non-response (failure to answer just one or some questions) to total non-response. Total non-response occurred because the interviewer was either unable to contact the respondent, no member of the household was able to provide the information, or the respondent refused to participate in the survey. Total non-response was handled by adjusting the weight of households who responded to the survey to compensate for those who did not respond.

In most cases, partial non-response to the survey occurred when the respondent did not understand or misinterpreted a question, refused to answer a question, or could not recall the requested information.

Item non-response was very low for the RTSS. Q02, Q03, Q04\_1, Q04\_2, Q04\_3, Q04\_4, Q04\_5 and Q04\_6 all had non-response rates which were less than .01%. Q08, which was the income class question, had a non-response rate of 2.1%, which is considered to be quite low, especially for an income related question.

Since it is an unavoidable fact that estimates from a sample survey are subject to sampling error, sound statistical practice calls for researchers to provide users with some indication of the magnitude of this sampling error. This section of the documentation outlines the <a href="mailto:measures">measures</a> of sampling error which Statistics Canada commonly uses and which it urges users producing estimates from this microdata file to use also.

The basis for measuring the potential size of sampling errors is the standard error of the estimates derived from survey results.

However, because of the large variety of estimates that can be produced from a survey, the standard error of an estimate is usually expressed relative to the estimate to which it pertains. This resulting measure, known as the coefficient of variation (CV) of an estimate, is obtained by dividing the standard error of the estimate by the estimate itself and is expressed as a percentage of the estimate.

For example, suppose that, based upon the survey results, one estimates that 1.7% of Canadian households did not have telephone service to their residence during the month of May 1999, and this estimate is found to have a standard error of .00078. Then the coefficient of variation of the estimate is calculated as:

$$\left(\frac{.00078}{.017}\right) \times 100\% = 4.6\%$$

## 10.0 Guidelines for Tabulation, Analysis and Release

This section of the documentation outlines the guidelines to be adhered to by users tabulating, analysing, publishing or otherwise releasing any data derived from the survey microdata file. With the aid of these guidelines, users of microdata should be able to produce the same figures as those produced by Statistics Canada and, at the same time, will be able to develop currently unpublished figures in a manner consistent with these established guidelines.

## 10.1

## **Rounding Guidelines**

In order that estimates for publication or other release derived from this microdata file correspond to those produced by Statistics Canada, users are urged to adhere to the following guidelines regarding the rounding of such estimates:

- Estimates in the main body of a statistical table a) are to be rounded to the nearest hundred units using the normal rounding technique. In normal rounding, if the first or only digit to be dropped is 0 to 4, the last digit to be retained is not changed. If the first or only digit to be dropped is 5 to 9, the last digit to be retained is raised by one. For example, in normal rounding to the nearest 100, if the last two digits are between 00 and 49, they are changed to 00 and the preceding digit (the hundreds digit) is left unchanged. If the last digits are between 50 and 99 they are changed to 00 and the preceding digit is incremented by 1.
- b) Marginal sub-totals and totals in statistical tables are to be derived from their corresponding unrounded components and then are to be rounded themselves to the nearest 100 units using normal rounding.

- c) Averages, proportions, rates and percentages are to be computed from unrounded components (i.e. numerators and/or denominators) and then are to be rounded themselves to one decimal using normal rounding. In normal rounding to a single digit, if the final or only digit to be dropped is 0 to 4, the last digit to be retained is not changed. If the first or only digit to be dropped is 5 to 9, the last digit to be retained is increased by 1.
  - d) Sums and differences of aggregates (or ratios) are to be derived from their corresponding unrounded components and then are to be rounded themselves to the nearest 100 units (or the nearest one decimal) using normal rounding.
  - e) In instances where, due to technical or other limitations, a rounding technique other than normal rounding is used resulting in estimates to be published or otherwise released which differ from corresponding estimates published by Statistics Canada, users are urged to note the reason for such differences in the publication or release document(s).
  - f) Under no circumstances are unrounded estimates to be published or otherwise released by users.
     Unrounded estimates imply greater precision than actually exists.

## 10.2

# Sample Weighting Guidelines for Tabulation

The sample design used for the Residential Telephone Service Survey was not self-weighting. When producing simple estimates, including the production of ordinary statistical tables, users must apply the proper sampling weight.

If proper weights are not used, the estimates derived from the microdata file cannot be considered to be representative of the survey population, and will not correspond to those produced by Statistics Canada.

Users should also note that some software packages may not allow the generation of estimates that exactly match those available from Statistics Canada, because of their treatment of the weight field.

## 10.2.1

# **Definitions of types of estimates:** Categorical vs. Quantitative

Before discussing how the Residential Telephone Service Survey data can be tabulated and analysed, it is useful to describe the two main types of point estimates of population characteristics which can be generated from the microdata file for the RTSS.

## **Categorical Estimates**

Categorical estimates are estimates of the number, or percentage of the surveyed population possessing certain characteristics or falling into some defined category. The number of households which did not have telephone service for their residence during the reference month or the proportion of households which had two or more telephone lines for their residence are examples of such estimates. An estimate of the number of households possessing a certain characteristic may also be referred to as an estimate of an aggregate.

## **Examples of Categorical Questions:**

Q: How many different telephone numbers are there for your residence?

R: 0, 1, 2 or more

Q: In 1998, was your total annual family income before taxes and deductions less or more than LICO?

R: Less, more

#### **Quantitative Estimates**

Quantitative estimates are estimates of totals or of means, medians and other measures of central tendency of quantities based upon some or all of the members of the surveyed population. They also specifically involve estimates of the form  $X/\hat{Y}$  where X is an estimate of surveyed population quantity total and  $\hat{Y}$  is an estimate of the number of persons in the surveyed population contributing to that total quantity. Note that there were no true quantitative questions in the RTSS application.

An example of a quantitative estimate is the average number of weeks for which unemployment insurance was collected for absences due to illness (taken from an unemployment survey). The numerator is an estimate of the total number of weeks for which unemployment insurance was collected for all persons experiencing an absence due to illness, and its denominator is the number of persons reporting an absence due to illness.

## **Examples of Quantitative Questions:**

Q: How many consecutive weeks was this last absence?
R: |\_| | Weeks
Q: How many separate periods of 2 or more weeks were you unable to work due to your own illness, accident or pregnancy?
R: |\_| | Periods

## 10.2.2

## **Tabulation of Categorical Estimates**

Estimates of the number of people with a certain characteristic can be obtained from the microdata file by summing the final weights of all records possessing the characteristic(s) of interest. Proportions and ratios of the form X/Y are obtained by:

- (a) summing the final weights of records having the characteristic of interest for the numerator (X),
- (b) summing the final weights of records having the characteristic of interest for the denominator (Y), then
- © dividing the numerator estimate by the denominator estimate.

## 10.2.3

## **Tabulation of Quantitative Estimates**

Estimates of quantities can be obtained from the microdata file by multiplying the value of the variable of interest by the final weight for each record, then summing this quantity over all records of interest. For example, using an unemployment survey, to obtain an estimate of the total number of weeks of employment insurance received by people whose last absence was due to pregnancy, multiply the value reported for weeks received EI by the final weight for the record, then sum this value over all records which report last absence due to pregnancy.

To obtain a weighted average of the form X/Y, the numerator (X) is calculated as for a quantitative estimate and the denominator (Y) is calculated as for a categorical estimate. For example, to estimate the <u>average</u> number of weeks EI was received by people whose last absence was due to pregnancy,

- estimate the total number of weeks as described above.
- (b) estimate the number of people in this category by summing the final weights of all records which report last absence due to pregnancy, then © divide estimate (a) by estimate (b).

## 10.3

## **Guidelines for Statistical Analysis**

The Residential Telephone Services Survey is based upon a complex sample design, with stratification, multiple stages of selection, and unequal probabilities of selection of respondents. Using data from such complex surveys presents problems to analysts because the survey design and the selection probabilities affect the estimation and variance calculation procedures that should be used. In order for survey estimates and analyses to be free from bias, the survey weights must be used.

While many analysis procedures found in statistical packages allow weights to be used, the meaning or definition of the weight in these procedures differ from that which is appropriate in a sample survey framework, with the result that while in many cases the estimates produced by the packages are correct, the variances that are calculated are poor. Variances for simple estimates such as totals, proportions and ratios (for qualitative variables) are provided in the accompanying Sampling Variability Tables.

For other analysis techniques (for example linear regression, logistic regression and analysis of variance), a method exists which can make the variances calculated by the standard packages more meaningful, by incorporating the unequal probabilities of selection. The method rescales the weights so that there is an average weight of 1.

For example, suppose that analysis of all male respondents is required. The steps to rescale the weights are as follows:

- select all respondents from the file who reported SEX=male
- Calculate the AVERAGE weight for these records by summing the original person weights from the microdata file for these records and then dividing by the number of respondents who reported SEX=male
- for each of these respondents, calculate a RESCALED weight equal to the original person weight divided by the AVERAGE weight
- perform the analysis for these respondents using the RESCALED weight.

However, because the stratification and clustering of the sample's design are still not taken into account, the variance estimates calculated in this way are likely to be under-estimates.

The calculation of truly meaningful variance estimates requires detailed knowledge of the design of the survey. Such detail cannot be given in this microdata file because of confidentiality. Variances that take the complete sample design into account can be calculated for many statistics by Statistics Canada on a cost recovery basis.

## 10.4

## **CV Release Guidelines**

Before releasing and/or publishing any estimate from the Residential Telephone Service Survey, users should first determine the quality level of the estimate. The quality levels are *acceptable*, *marginal* and *unacceptable*. Data quality is affected by both sampling and non-sampling errors as discussed in section 8. However for this purpose, the quality level of an estimate will be determined only on the basis of sampling error as reflected by the coefficient of variation as shown in the table below. Nonetheless, users should be sure to read section 8 to be more fully aware of the quality characteristics of these data.

First, the number of respondents who contribute to the calculation of the estimate should be determined. If this number is less than 30, the weighted estimate should be considered to be of unacceptable quality.

For weighted estimates based on sample sizes of 30 or more, users should determine the coefficient of variation of the estimate and follow the guidelines below. These quality level guidelines should be applied to weighted rounded estimates.

All estimates can be considered releasable. However, those of marginal or unacceptable quality level must be accompanied by a warning to caution subsequent users.

## **Quality Level Guidelines**

Quality Level of Estimate	Guidelines
1. Acceptable	Estimates have: a sample size of 30 or more, and low coefficients of variation in the range 0.0% - 16.5%  No warning is required.
2. Marginal	Estimates have: a sample size of 30 or more, and high coefficients of variation in the range 16.6% - 33.3%.  Estimates should be flagged with the letter M (or some similar identifier). They should be accompanied by a warning to caution subsequent users about the high levels of error, associated with the estimates.
3. Unacceptable	Estimates have: a sample size of less than 30, or very high coefficients of variation in excess of 33.3%.  Statistics Canada recommends not to release estimates of unacceptable quality. However, if the user chooses to do so then estimates should be flagged with the letter U (or some similar identifier) and the following warning should accompany the estimates:  "The user is advised that (specify the data) do not meet Statistics Canada's quality standards for this statistical program. Conclusions based on these data will be unreliable, and most likely invalid. These data and any consequent findings should not be published. If the user chooses to publish these data or findings, then this disclaimer must be published with the data."

## 11.0 Approximate Sampling Variability Tables

In order to supply coefficients of variation which would be applicable to a wide variety of categorical estimates produced from this microdata file and which could be readily accessed by the user, a set of Approximate Sampling Variability Tables has been produced. These "look-up" tables allow the user to obtain an approximate coefficient of variation based on the size of the estimate calculated from the survey data.

The coefficients of variation (C.V.) are derived using the variance formula for simple random sampling and incorporating a factor which reflects the multi-stage, clustered nature of the sample design. This factor, known as the design effect, was determined by first calculating design effects for a wide range of characteristics and then choosing from among these a conservative value to be used in the look-up tables which would then apply to the entire set of characteristics.

The table below shows the design effects, sample sizes and population counts by province which were used to produce the Approximate Sampling Variability Tables.

PROVINCE	DESIGN EFFECT	SAMPLE SIZE	POPULATION
Newfoundland	1.1	1523	190297
Prince Edward Island	1.54	1104	51071
Nova Scotia	1.85	2743	361986
New Brunswick	1.09	2235	281149
Quebec	1.96	7906	2980703
Ontario	1.41	11917	4277134
Manitoba	1.32	2986	424627
Saskatchewan	1.61	3092	383846
Alberta	1.04	3103	1079912
British Columbia	1.48	3731	1551915
Atlantic Provinces	1.43	7605	884503
Prairies	1.19	9181	1888385
Canada	1.94	40340	11582640

All coefficients of variation in the Approximate Sampling Variability Tables are <u>approximate</u> and, therefore, unofficial. Estimates of actual variance for specific variables may be obtained from Statistics Canada on a cost-recovery basis. The use of actual variance estimates would allow users to release otherwise unreleaseable estimates, i.e., estimates with coefficients of variation in the 'confidential' range.

<u>Remember</u>: if the number of observations on which an estimate is based is less than 30, the weighted estimate should not be released regardless of the value of the coefficient of variation for this estimate. This is because the formulas used for estimating the variance do not hold true for small sample sizes.

## 11.1

# How to use the C.V. tables for Categorical Estimates

The following rules should enable the user to determine the approximate coefficients of variation from the Sampling Variability Tables for estimates of the number, proportion or percentage of the surveyed population possessing a certain characteristic and for ratios and differences between such estimates.

## Rule 1: Estimates of Numbers Possessing a Characteristic (Aggregates)

The coefficient of variation depends only on the size of the estimate itself. On the Sampling Variability Table for the appropriate geographic area, locate the estimated number in the left-most column of the table (headed "Numerator of Percentage") and follow the asterisks (if any) across to the first figure encountered. This figure is the approximate coefficient of variation.

## Rule 2: Estimates of Proportions or Percentages Possessing a Characteristic

The coefficient of variation of an estimated proportion or percentage depends on both the size of the proportion or percentage and the size of the total upon which the proportion or percentage is based. Estimated proportions or percentages are relatively more reliable than the corresponding estimates of the numerator of the proportion or percentage, when the proportion or percentage is based upon a sub-group of the population. For example, the <u>proportion</u> of "households which did not have telephone service for their residence during the reference period" is more reliable than the estimated <u>number</u> of "households which did not have telephone service for their residence during the reference period". (Note that in the tables the CV's decline in value reading from left to right).

When the proportion or percentage is based upon the total population of the geographic area covered by the table, the CV of the proportion or percentage is the same as the CV of the numerator of the proportion or percentage. In this case, Rule 1 can be used.

When the proportion or percentage is based upon a subset of the total population (e.g. those in a particular sex or age group), reference should be made to the proportion or percentage (across the top of the table) and to the numerator of the proportion or percentage (down the left side of the table). The intersection of the appropriate row and column gives the coefficient of variation.

## Rule 3: Estimates of Differences Between Aggregates or Percentages

The standard error of a difference between two estimates is approximately equal to the square root of the sum of squares of each standard error considered separately. That is, the standard error of a difference  $(\mathring{d} = \mathring{X}_1 - \mathring{X}_2)$  is:

$$\sigma_{\hat{d}} = \sqrt{(\hat{X}_1 \alpha_1)^2 + (\hat{X}_2 \alpha_2)^2}$$

where  $X_1$  is estimate 1,  $X_2$  is estimate 2, and  $\alpha_1$  and  $\alpha_2$  are the coefficients of variation of  $X_1$  and  $X_2$  respectively. The coefficient of variation of  $\hat{\alpha}$  is given by  $\sigma_{\hat{\alpha}}/\hat{\alpha}$ . This formula is accurate for the difference between separate and uncorrelated characteristics, but is only approximate otherwise.

#### Rule 4: Estimates of Ratios

In the case where the numerator is a subset of the denominator, the ratio should be converted to a percentage and Rule 2 applied. This would apply, for example, to the case where the denominator is the number of "households which did not have telephone service for their residence during the reference period" and the numerator is the number of "households which did not have telephone service to their residence during the reference period because they could not afford it".

In the case where the numerator is not a subset of the denominator, as for example, the ratio of the number of "households in Quebec whose total annual income for 1998 was bellow the low income cut off" as compared to the number of "households in Ontario whose total annual income for 1998 was bellow the low income cut off", the standard deviation of the ratio of the estimates is approximately equal to the square root of the sum of squares of each coefficient of variation considered separately multiplied by R. That is, the standard error of a ratio ( $R = X_1 / X_2$ ) is:

$$\sigma_{\hat{R}} = \hat{R} \sqrt{\alpha_1^2 + \alpha_2^2}$$

where  $\alpha_1$  and  $\alpha_2$  are the coefficients of variation of  $X_1$  and  $X_2$  respectively. The coefficient of variation of R is given by  $\sigma_R/R$ . The formula will tend to overstate the error, if  $X_1$  and  $X_2$  are positively correlated and understate the error if  $X_1$  and  $X_2$  are negatively correlated.

#### Rule 5: Estimates of Differences of Ratios

In this case, Rules 3 and 4 are combined. The CV's for the two ratios are first determined using Rule 4, and then the CV of their difference is found using Rule 3.

## 11.1.1

# **Examples of using the C.V. tables for Categorical Estimates**

The following 'real life' examples are included to assist users in applying the foregoing rules.

## Example 1 : Estimates of Numbers Possessing a Characteristic (Aggregates)

Suppose that a user estimates that 194,951 households did not have telephone service for their residence during the reference period. How does the user determine the coefficient of variation of this estimate?

- (1) Refer to the CV table for CANADA.
- (2) The estimated aggregate (194,951) does not appear in the left-hand column (the 'Numerator of Percentage' column), so it is necessary to use the figure closest to it, namely 200,000.
- (3) The coefficient of variation for an estimated aggregate is found by referring to the first non asterisk entry on that row, namely, 5.2%.
- (4) So the approximate coefficient of variation of the estimate is 5.2%.

The finding that there were 194,951 households did not have telephone service for their residence during the reference period is publishable with no qualifications.

## Example 2: Estimates of Proportions or Percentages Possessing a Characteristic

Suppose that the user estimates that 109,276/194,951=56.1% of households which did not have telephone service for their residence during the reference period reported that they could not afford telephone service. How does the user determine the coefficient of variation of this estimate?

- (1) Refer to the table for CANADA.
- (2) Because the estimate is a percentage which is based on a subset of the total population (i.e.,households which did not have telephone service for their residence during the reference period), it is necessary to use both the percentage (56.1%) and the numerator portion of the percentage (109,276) in determining the coefficient of variation.
- (3) The numerator, 109,276, does not appear in the left-hand column (the 'Numerator of Percentage' column) so it is necessary to use the figure closet to it, namely 100,000. Similarly, the percentage estimate does not appear as any of the column headings, so it is necessary to use the figure closest to it, 50.0%.
- (4) The figure at the intersection of the row and column used, namely 5.3% is the coefficient of variation to be used.
- (5) So the approximate coefficient of variation of the estimate is 5.3%. The finding that 56.1% of households which did not have telephone service for their residence during the reference period could not afford telephone service can be published with no qualifications.

## Example 3: Estimates of Differences Between Aggregates or Percentages

Suppose that a user estimates that 796,933/2,980,703=26.7% of households in Quebec reported that their total annual income for 1998 was less than LICO, while 799,482/4,277,134=18.7% of households in Ontario reported that their total annual income for 1998 was less than LICO. How does the user determine the coefficient of variation of the difference between these two estimates?

- (1) Using the QUEBEC and ONTARIO CV table in the same manner as described in example 1 gives the CV of the estimate for households in Quebec as 2.6, and the CV of the estimate for households in Ontario as 2.3%.
- (2) Using rule 3, the standard error of a difference  $(\hat{a} = X_1 X_2)$  is:

$$\sigma_{\hat{d}} = \sqrt{(\hat{X}_1 \alpha_1)^2 + (\hat{X}_2 \alpha_2)^2}$$

where  $X_1$  is estimate 1,  $X_2$  is estimate 2, and  $\alpha_1$  and  $\alpha_2$  are the coefficients of variation of  $X_1$  and  $X_2$  respectively.

That is, the standard error of the difference  $\hat{d} = (.267 - .187) = .080$  is:

$$\sigma_{\hat{d}} = \sqrt{[(.267)(.026)]^2 + [(.187)(.023)]^2}$$

$$= \sqrt{(.0000481) + (.0000184)}$$

$$= .0082$$

- (3) The coefficient of variation of  $\hat{a}$  is given by  $\sigma_{\hat{a}}/\hat{a} = .0082/.080 = .103$ .
- (4) So the approximate coefficient of variation of the difference between the estimates is 10.3%. This estimate is publishable with no qualifications.

## **Example 4: Estimates of Ratios**

Suppose that the user estimates that 796,933 households in Quebec reported that their total annual income for 1998 was less than LICO, while 799,482 households in Ontario reported that their total annual income for 1998 was less than LICO. The user is interested in comparing the estimate of Quebec households versus that of Ontario households in the form of a ratio. How does the user determine the coefficient of variation of this estimate?

- (1) First of all, this estimate is a ratio estimate, where the numerator of the estimate (=  $X_1$ ) is the number of households in Quebec which reported that their total annual income for 1998 was less than LICO. The denominator of the estimate (=  $X_2$ ) is the number of households in Ontario which reported that their total annual income for 1998 was less than LICO.
- (2) Refer to the tables for QUEBEC and ONTARIO.
- (3) The numerator of this ratio estimate is 796,933. The figure closest to it is 750,000. The coefficient of variation for this estimate is found by referring to the first non-asterisk entry on that row in the QUEBEC table, namely, 2.6%.
- (4) The denominator of this ratio estimate is 799,482. The figure closest to it is 750,000. The coefficient of variation for this estimate is found by referring to the first non-asterisk entry on that row in the ONTARIO table, namely, 2.3%.
- (5) So the approximate coefficient of variation of the ratio estimate is given by rule 4, which is,

$$\alpha_{\hat{R}} = \sqrt{\alpha_1^2 + \alpha_2^2}$$

where  $\alpha_{\scriptscriptstyle 1}$  and  $\alpha_{\scriptscriptstyle 2}$  are the coefficients of variation of  $X_{\scriptscriptstyle 1}$  and  $X_{\scriptscriptstyle 2}$  respectively.

That is,

$$\alpha_{\hat{R}} = \sqrt{(.026)^2 + (.023)^2}$$
$$= 0.035$$

The obtained ratio of Quebec versus Ontario households whose total annual income for 1998 is less than the LICO is 796,933/799,482 - which is 0.997:1. The coefficient of variation of this estimate is 3.5%, which is releasable with no qualifications.

## 11.2

# How to use the CV tables to obtain Confidence Limits

Although coefficients of variation are widely used, a more intuitively meaningful measure of sampling error is the confidence interval of an estimate. A confidence interval constitutes a statement on the level of confidence that the true value for the population lies within a specified range of values. For example a 95% confidence interval can be described as follows:

If sampling of the population is repeated indefinitely, each sample leading to a new confidence interval for an estimate, then in 95% of the samples the interval will cover the true population value.

Using the standard error of an estimate, confidence intervals for estimates may be obtained under the assumption that under repeated sampling of the population, the various estimates obtained for a population characteristic are normally distributed about the true population value. Under this assumption, the chances are about 68 out of 100 that the difference between a sample estimate and the true population value would be less than one standard error, about 95 out of 100 that the difference would be less than two standard errors, and about 99 out of 100 that the differences would be less than three standard errors. These different degrees of confidence are referred to as the confidence levels.

Confidence intervals for an estimate,  $\hat{X}$ , are generally expressed as two numbers, one below the estimate and one above the estimate, as  $(\hat{X}-k, \hat{X}+k)$  where k is determined depending upon the level of confidence desired and the sampling error of the estimate.

Confidence intervals for an estimate can be calculated directly from the Approximate Sampling Variability Tables by first determining from the appropriate table the coefficient of variation of the estimate  $\hat{X}$ , and then using the following formula to convert to a confidence interval CI:

$$CI_X = [\hat{X} - t\hat{X}\alpha_{\hat{X}}, \hat{X} + t\hat{X}\alpha_{\hat{X}}]$$

where  $\alpha_{\hat{X}}$  is the determined coefficient of variation of  $\hat{X},$  and

t = 1 if a 68% confidence interval is desired

t = 1.6 if a 90% confidence interval is desired

t = 2 if a 95% confidence interval is desired

t = 3 if a 99% confidence interval is desired.

Note:

Release guidelines which apply to the estimate also apply to the confidence interval. For example, if the estimate is not releasable, then the confidence interval is not releasable either.

## 11.2.1

# Example of using the CV tables to obtain confidence limits

A 95% confidence interval for the estimated proportion of households which did not have telephone service for their residence during the reference period because they could not afford telephone service (from Example 2, section 10.2) would be calculated as follows.

 $\hat{X} = 56.1\%$  (or expressed as a proportion = .561)

t = 2

 $\alpha_{X}$  = 5.3% (.053 expressed as a proportion) is the coefficient of variation of this estimate as determined from the tables.

 $CI_x = \{.561 - (2) (.561) (.053), .561 + (2) (.561) (.053)\}$ 

 $CI_x = \{.561 - .059, .561 + .059\}$ 

 $CI_x = \{.502.620\}$ 

With 95% confidence it can be said that between 50.2% and 62.0% of households which did not have telephone service for their residence during the reference period reported that they could not afford telephone service.

## 11.3

# How to use the CV tables to do a t-test

Standard errors may also be used to perform hypothesis testing, a procedure for distinguishing between population parameters using sample estimates. The sample estimates can be numbers, averages, percentages, ratios, etc. Tests may be performed at various levels of significance, where a level of significance is the probability of concluding that the characteristics are different when, in fact, they are identical.

Let  $\mathbf{X_1}$  and  $\mathbf{X_2}$  be sample estimates for 2 characteristics of interest. Let the standard error on the difference  $\mathbf{X_1}$  -  $\mathbf{X_2}$  be  $\sigma_{\mathbf{\tilde{G}}}$ .

If 
$$=\frac{\hat{X_1}-\hat{X_2}}{\sigma_{\hat{d}}}$$
 is between -2 and 2, then no conclusion

about the difference between the characteristics is justified at the 5% level of significance. If however, this ratio is smaller than -2 or larger than +2, the observed difference is significant at the 0.05 level. That is to say that the characteristics are significant.

## 11.3.1

# Example of using the CV tables to do a t-test

Let us suppose we wish to test, at a 5% level of significance, the hypothesis that there is no difference between the proportion of households in Quebec which reported that their total annual income for 1998 was less than LICO, and the proportion of households in Ontario which reported that their total annual income for 1998 was less than LICO. From example 3, section 10.2, the standard error of the difference between these two estimates was found to be = .0082. Hence,

$$t = \frac{\hat{X}_1 - \hat{X}_2}{\sigma_{\hat{x}}} = \frac{.267 - .187}{.0082} = \frac{.080}{.0082} = 9.76.$$

Since t = 9.76 is greater than 2, it must be concluded that there is a significant difference between the two estimates at the 0.05 level of significance.

## 11.4

## **Coefficients of Variation for Quantitative Estimates**

For quantitative estimates, special tables would have to be produced to determine their sampling error. Since all of the variables for the RTSS are primarily categorical in nature, this has not been done.

As a general rule, however, the coefficient of variation of a quantitative total will be larger than the coefficient of variation of the corresponding category estimate (i.e., the estimate of the number of persons contributing to the quantitative estimate). If the corresponding category estimate is not releasable, the quantitative estimate will not be either. For example, in an absence from work survey, the coefficient of variation of the total number of weeks absent from work would be greater than the coefficient of variation of the corresponding proportion of paid workers with an absence. Hence if the coefficient of variation of the proportion is not releasable, then the coefficient of variation of the corresponding quantitative estimate will also not be releasable.

Coefficients of variation of such estimates can be derived as required for a specific estimate using a technique known as pseudo replication. This involves dividing the records on the microdata files into subgroups (or replicates) and determining the variation in the estimate from replicate to replicate. Users wishing to derive coefficients of variation for quantitative estimates may contact Statistics Canada for advice on the allocation of records to appropriate replicates and the formulae to be used in these calculations.

## 11.5

## Release cut-off's for the RTSS

The minimum size of the estimate at the provincial, regional and Canada levels are specified in the table below. Estimates smaller than the minimum size given in the "Not Releasable" column may not be released under any circumstances.

## Table of Release Cut-offs

Province	Unqualified	Qualified	Confidential	Not Releasable
Newfoundland	5,000 & +	2,000 - 4,900	1,000 - 1,900	under 1,000
Prince Edward Island	2,500 & +	1,000 - 2,400	500 - 900	under 500
Nova Scotia	9,000 & +	4,000 - 8,900	2,000 - 3,900	under 2,000
New Brunswick	5,000 & +	2,000 - 4,900	1,000 - 1,900	under 1,000
Quebec	27,000 & +	12,000 - 26,900	6,500 -11,900	under 6,500
Ontario	18,500 & +	8,000 - 18,400	4,500 - 7,900	under 4,500
Manitoba	7,000 & +	3,000 - 6,900	1,500 - 2,900	under 1,500
Saskatchewan	7,000 & +	3,000 - 6,900	2,000 - 2,900	under 2,000
Alberta	13,000 & +	6,000 - 12,900	3,500 - 5,900	under 3,500
British Columbia	22,500 & +	10,000 - 22,400	5,500 - 9,900	under 5,500
Atlantic Provinces	6,000 & +	2,500 - 5,900	1,500 - 2,400	under 1,500
Prairie Provinces	9,000 & +	4,000 - 8,900	2,000 - 3,900	under 2,000
CANADA	20,500 & +	9,000 - 20,400	5,000 - 8,900	under 5,000

# 11.6 CV Tables

RESIDENTIAL TELEPHONE SERVICES SURVEY - 0599 (re-weighted)

Approximate Sampling Variability Tables for NEWFOUNDLAND

NUMERATOR O PERCENTAGE						ESTIMATE	D PERCEN	ΓAGE						
('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	******	36.7	36.6	36.0	35.0	34.0	33.0	32.0	30.9	29.8	28.6	26.1	20.2	11.7
2	*****		25.8	25.4	24.8	24.1	23.4	22.6	21.8	21.1	20.2	18.5	14.3	8.3
3	*****		21.1	20.8	20.2	19.7	19.1	18.5	17.8	17.2	16.5	15.1	11.7	6.7
4	*****			18.0	17.5	17.0	16.5	16.0	15.4	14.9	14.3	13.1	10.1	5.8
5	*****	******	*****	16.1	15.7	15.2	14.8	14.3	13.8	13.3	12.8	11.7	9.0	5.2
6	*****			14.7	14.3	13.9	13.5	13.1	12.6	12.2	11.7	10.7	8.3	4.8
7	*****	******	*****	13.6	13.2	12.9	12.5	12.1	11.7	11.3	10.8	9.9	7.6	4.4
8	*****	******	*****	12.7	12.4	12.0	11.7	11.3	10.9	10.5	10.1	9.2	7.2	4.1
9	*****	******	*****	12.0	11.7	11.3	11.0	10.7	10.3	9.9	9.5	8.7	6.7	3.9
10	*****	******	******	*****	11.1	10.8	10.4	10.1	9.8	9.4	9.0	8.3	6.4	3.7
11	*****	******	******	*****	10.6	10.3	10.0	9.6	9.3	9.0	8.6	7.9	6.1	3.5
12	*****	******	******	*****	10.1	9.8	9.5	9.2	8.9	8.6	8.3	7.5	5.8	3.4
13	*****	******	******	*****	9.7	9.4	9.2	8.9	8.6	8.3	7.9	7.2	5.6	3.2
14	*****	******	******	*****	9.4	9.1	8.8	8.5	8.3	8.0	7.6	7.0	5.4	3.1
15	*****	******	******	*****	9.0	8.8	8.5	8.3	8.0	7.7	7.4	6.7	5.2	3.0
16	*****	******	******	*****	8.8	8.5	8.3	8.0	7.7	7.4	7.2	6.5	5.1	2.9
17	*****	******	******	*****	8.5	8.3	8.0	7.8	7.5	7.2	6.9	6.3	4.9	2.8
18	*****	******	******	*****	8.3	8.0	7.8	7.5	7.3	7.0	6.7	6.2	4.8	2.8
19	*****	******	******	*****	8.0	7.8	7.6	7.3	7.1	6.8	6.6	6.0	4.6	2.7
20	******	******	******	*****	*****	7.6	7.4	7.2	6.9	6.7	6.4	5.8	4.5	2.6
21	*****	******	******	*****	*****	7.4	7.2	7.0	6.7	6.5	6.2	5.7	4.4	2.5
22	*****	******	******	****	*****	7.3	7.0	6.8	6.6	6.3	6.1	5.6	4.3	2.5
23	*****	******	******	****	*****	7.1	6.9	6.7	6.4	6.2	6.0	5.4	4.2	2.4
24	*****	******	******	*****	*****	6.9	6.7	6.5	6.3	6.1	5.8	5.3	4.1	2.4
25	*****	******	******	*****	*****	6.8	6.6	6.4	6.2	6.0	5.7	5.2	4.0	2.3
30	*****	******	******	*****	*****	*****	6.0	5.8	5.6	5.4	5.2	4.8	3.7	2.1
35	*****	******	******	*****	*****	*****	5.6	5.4	5.2	5.0	4.8	4.4	3.4	2.0
40	*****	******	******	*****	*****	*****	*****	5.1	4.9	4.7	4.5	4.1	3.2	1.8
45	*****	******	******	****	*****	*****	*****	4.8	4.6	4.4	4.3	3.9	3.0	1.7
50	*****	******	******	*****	*****	*****	*****	*****	4.4	4.2	4.0	3.7	2.9	1.7
55	*****	******	******	****	*****	*****	*****	*****	4.2	4.0	3.9	3.5	2.7	1.6
60	*****	******	******	****	*****	*****	*****	*****	*****	3.8	3.7	3.4	2.6	1.5
65	*****	******	******	*****	*****	*****	*****	*****	*****	3.7	3.5	3.2	2.5	1.4
70	*****	******	******	*****	*****	*****	*****	*****	******	****	3.4	3.1	2.4	1.4
75	*****	******	******	*****	*****	*****	*****	*****	******	****	3.3	3.0	2.3	1.3
80	*****	******	******	*****	*****	*****	*****	*****	******	******		2.9	2.3	1.3
85	*****	******	******	*****	*****	****	*****	*****	******	******	*****	2.8	2.2	1.3
90	*****	******	******	*****	*****	****	*****	*****	******	******	*****	2.8	2.1	1.2
95	*****	******	******	*****	*****	****	*****	*****	******	******	*****	2.7	2.1	1.2
100	*****	******	******	*****	*****	****	*****	*****	******	******	******		2.0	1.2
125	*****	******	******	*****	******	*****	*****	*****	******	******	******	*****	1.8	1.0
150	*****													1.0

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for PRINCE EDWARD ISLAND

NUMERATOR O														
PERCENTAGE														
('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	******	*****	26.1	25.7	25.0	24.3	23.6	22.9	22.1	21.3	20.4	18.7	14.5	8.3
2	******	*****	*****	18.2	17.7	17.2	16.7	16.2	15.6	15.1	14.5	13.2	10.2	5.9
3	******	*****	******	*****	14.5	14.1	13.6	13.2	12.8	12.3	11.8	10.8	8.3	4.8
4	******	*****	*****	*****	12.5	12.2	11.8	11.4	11.0	10.6	10.2	9.3	7.2	4.2
5	******	*****	*****	*****	11.2	10.9	10.6	10.2	9.9	9.5	9.1	8.3	6.5	3.7
6	******	*****	*****	*****	*****	9.9	9.6	9.3	9.0	8.7	8.3	7.6	5.9	3.4
7	******	*****	*****	*****	*****	9.2	8.9	8.6	8.3	8.0	7.7	7.1	5.5	3.2
8	******	*****	*****	*****	*****	*****	8.3	8.1	7.8	7.5	7.2	6.6	5.1	3.0
9	******	*****	*****	*****	******	*****	7.9	7.6	7.4	7.1	6.8	6.2	4.8	2.8
10	******	*****	******	*****	*****	*****	7.5	7.2	7.0	6.7	6.5	5.9	4.6	2.6
11	******	*****	*****	*****	******	*****	*****	6.9	6.7	6.4	6.2	5.6	4.4	2.5
12	******	*****	*****	*****	*****	*****	*****	6.6	6.4	6.1	5.9	5.4	4.2	2.4
13	******	*****	*****	*****	******	*****	*****	*****	6.1	5.9	5.7	5.2	4.0	2.3
14	******	*****	*****	*****	******	*****	*****	*****	5.9	5.7	5.5	5.0	3.9	2.2
15	******	*****	*****	*****	******	*****	*****	*****	5.7	5.5	5.3	4.8	3.7	2.2
16	******	*****	*****	*****	******	*****	*****	*****	*****	5.3	5.1	4.7	3.6	2.1
17	******	*****	******	*****	******	*****	*****	*****	*****	5.2	5.0	4.5	3.5	2.0
18	******	*****	*****	*****	******	*****	*****	*****	******	****	4.8	4.4	3.4	2.0
19	******	*****	******	*****	******	*****	*****	*****	******	*****	4.7	4.3	3.3	1.9
20	******	*****	*****	*****	******	*****	*****	*****	*****	*****	4.6	4.2	3.2	1.9
21	******	*****	******	*****	******	*****	*****	*****	******	******	*****	4.1	3.2	1.8
22	******	*****	*****	*****	******	*****	*****	*****	*****	******	*****	4.0	3.1	1.8
23	******	*****	*****	*****	******	*****	*****	*****	*****	******	*****	3.9	3.0	1.7
24	******	*****	*****	*****	******	*****	*****	*****	*****	******	*****	3.8	3.0	1.7
25	******	*****	*****	*****	******	*****	*****	*****	*****	******	*****	3.7	2.9	1.7
30	******	*****	*****	*****	******	*****	*****	*****	*****	******	******	*****	2.6	1.5
35	******	*****	******	*****	******	*****	*****	*****	******	******	******	*****	2.4	1.4
40	******	*****	******	*****	******	*****	*****	*****	******	******	******	******	*****	1.3
45	******	*****	*****	*****	*****	****	*****	*****	*****	******	*****	******	*****	1.2

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for NOVA SCOTIA

NUMERATOR O	F				ĺ	ESTIMATE	) PERCENT	TAGE						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	*****	49.0	48.7	48.0	46.7	45.4	44.0	42.6	41.2	39.7	38.1	34.8	27.0	15.6
2	******	34.6	34.5	33.9	33.0	32.1	31.1	30.1	29.1	28.1	27.0	24.6	19.1	11.0
3	******	28.3	28.1	27.7	27.0	26.2	25.4	24.6	23.8	22.9	22.0	20.1	15.6	9.0
4	*****		24.4	24.0	23.3	22.7	22.0	21.3	20.6	19.8	19.1	17.4	13.5	7.8
5	*****		21.8	21.5	20.9	20.3	19.7	19.1	18.4	17.7	17.1	15.6	12.1	7.0
6	*****	*****	19.9	19.6	19.1	18.5	18.0	17.4	16.8	16.2	15.6	14.2	11.0	6.4
7	*****		18.4	18.1	17.6	17.2	16.6	16.1	15.6	15.0	14.4	13.2	10.2	5.9
8	*****			17.0	16.5	16.0	15.6	15.1	14.6	14.0	13.5	12.3	9.5	5.5
9	*****			16.0	15.6	15.1	14.7	14.2	13.7	13.2	12.7	11.6	9.0	5.2
10	*****			15.2	14.8	14.4	13.9	13.5	13.0	12.5	12.1	11.0	8.5	4.9
11	*****			14.5	14.1	13.7	13.3	12.9	12.4	12.0	11.5	10.5	8.1	4.7
12	*****			13.8	13.5	13.1	12.7	12.3	11.9	11.5	11.0	10.0	7.8	4.5
13	*****			13.3	13.0	12.6	12.2	11.8	11.4	11.0	10.6	9.7	7.5	4.3
14	*****			12.8	12.5	12.1	11.8	11.4	11.0	10.6	10.2	9.3	7.2	4.2
15	*****			12.4	12.1	11.7	11.4	11.0	10.6	10.2	9.8	9.0	7.0	4.0
16	*****			12.0	11.7	11.3	11.0	10.7	10.3	9.9	9.5	8.7	6.7	3.9
17	*****			11.6	11.3	11.0	10.7	10.3	10.0	9.6	9.2	8.4	6.5	3.8
18	*****			11.3	11.0	10.7	10.4	10.0	9.7	9.4	9.0	8.2	6.4	3.7
19	*****				10.7	10.4	10.1	9.8	9.4	9.1	8.7	8.0	6.2	3.6
20	*****				10.4	10.1	9.8	9.5	9.2	8.9	8.5	7.8	6.0	3.5
21	*****				10.2	9.9	9.6	9.3	9.0	8.7	8.3	7.6	5.9	3.4
22	*****				10.0	9.7	9.4	9.1	8.8	8.5	8.1	7.4	5.7	3.3
23	*****				9.7	9.5	9.2	8.9	8.6	8.3	8.0	7.3	5.6	3.2
24	*****				9.5	9.3	9.0	8.7	8.4	8.1	7.8	7.1	5.5	3.2
25	*****				9.3	9.1	8.8	8.5	8.2	7.9	7.6	7.0	5.4	3.1
30	*****				8.5	8.3	8.0	7.8	7.5	7.2	7.0	6.4	4.9	2.8
35	*****				7.9	7.7	7.4	7.2	7.0	6.7	6.4	5.9	4.6	2.6
40	*****					7.2	7.0	6.7	6.5	6.3	6.0	5.5	4.3	2.5
45	*****					6.8	6.6	6.4	6.1	5.9	5.7	5.2	4.0	2.3
50	*****					6.4	6.2	6.0	5.8	5.6	5.4	4.9	3.8	2.2
55	*****						5.9	5.7	5.6	5.4	5.1	4.7	3.6	2.1
60 65	*****						5.7 5.5	5.5	5.3	5.1	4.9	4.5	3.5	2.0
70	*****						5.5	5.3 5.1	5.1 4.9	4.9 4.7	4.7 4.6	4.3 4.2	3.3	1.9 1.9
70 75	*****							4.9	4.9		4.6	4.2	3.2 3.1	1.9
75 80	*****							4.9	4.6	4.6 4.4	4.4	3.9	3.1	1.0
85	*****							4.6	4.5	4.4	4.3	3.8	2.9	1.7
90	*****							4.5	4.3	4.3	4.1	3.7	2.8	1.7
95	*****								4.2	4.2	3.9	3.7	2.8	1.6
100	*****								4.2	4.1	3.8	3.5	2.7	1.6
125	*****								–	3.5	3.4	3.5	2.7	1.0
150	****											2.8	2.4	1.4
200	*****												1.9	1.1
250	****												1.7	1.1
300	*****													0.9

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for NEW BRUNSWICK

NUMERATOR O	F				1	ESTIMATE	) PERCEN	ΓAGE						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	*****	36.7	36.5	35.9	35.0	34.0	33.0	31.9	30.9	29.7	28.6	26.1	20.2	11.7
2	******	25.9	25.8	25.4	24.7	24.0	23.3	22.6	21.8	21.0	20.2	18.4	14.3	8.2
3	*****	*****	21.1	20.8	20.2	19.6	19.0	18.4	17.8	17.2	16.5	15.1	11.7	6.7
4	*****	*****	18.3	18.0	17.5	17.0	16.5	16.0	15.4	14.9	14.3	13.0	10.1	5.8
5	*****	*****	16.3	16.1	15.6	15.2	14.8	14.3	13.8	13.3	12.8	11.7	9.0	5.2
6	*****	******	****	14.7	14.3	13.9	13.5	13.0	12.6	12.1	11.7	10.6	8.2	4.8
7	*****	*****	*****	13.6	13.2	12.9	12.5	12.1	11.7	11.2	10.8	9.9	7.6	4.4
8	*****	******	*****	12.7	12.4	12.0	11.7	11.3	10.9	10.5	10.1	9.2	7.1	4.1
9	*****	******	****	12.0	11.7	11.3	11.0	10.6	10.3	9.9	9.5	8.7	6.7	3.9
10	*****	******	*****	11.4	11.1	10.8	10.4	10.1	9.8	9.4	9.0	8.2	6.4	3.7
11	*****	******	****	10.8	10.5	10.3	9.9	9.6	9.3	9.0	8.6	7.9	6.1	3.5
12	*****	******	*****	10.4	10.1	9.8	9.5	9.2	8.9	8.6	8.2	7.5	5.8	3.4
13	*****	******	****	10.0	9.7	9.4	9.1	8.9	8.6	8.2	7.9	7.2	5.6	3.2
14	*****	******	****	9.6	9.4	9.1	8.8	8.5	8.2	7.9	7.6	7.0	5.4	3.1
15	*****	******	*****	*****	9.0	8.8	8.5	8.2	8.0	7.7	7.4	6.7	5.2	3.0
16	*****	******	******	*****	8.7	8.5	8.2	8.0	7.7	7.4	7.1	6.5	5.1	2.9
17	*****	******	*****	*****	8.5	8.2	8.0	7.7	7.5	7.2	6.9	6.3	4.9	2.8
18	*****	******	******	*****	8.2	8.0	7.8	7.5	7.3	7.0	6.7	6.1	4.8	2.7
19	*****	******	*****	*****	8.0	7.8	7.6	7.3	7.1	6.8	6.6	6.0	4.6	2.7
20	*****	*****	******	*****	7.8	7.6	7.4	7.1	6.9	6.6	6.4	5.8	4.5	2.6
21	*****	******	*****	*****	7.6	7.4	7.2	7.0	6.7	6.5	6.2	5.7	4.4	2.5
22	*****	******	*****	*****	7.5	7.2	7.0	6.8	6.6	6.3	6.1	5.6	4.3	2.5
23	*****	******	*****	*****	7.3	7.1	6.9	6.7	6.4	6.2	6.0	5.4	4.2	2.4
24	*****	******	*****	*****	7.1	6.9	6.7	6.5	6.3	6.1	5.8	5.3	4.1	2.4
25	*****	******	*****	*****	7.0	6.8	6.6	6.4	6.2	5.9	5.7	5.2	4.0	2.3
30	*****	******	*****	*****	*****	6.2	6.0	5.8	5.6	5.4	5.2	4.8	3.7	2.1
35	*****	******	*****	*****	*****	5.7	5.6	5.4	5.2	5.0	4.8	4.4	3.4	2.0
40	*****	******	*****	******	*****	5.4	5.2	5.1	4.9	4.7	4.5	4.1	3.2	1.8
45	*****	******	*****	*****	*****	*****	4.9	4.8	4.6	4.4	4.3	3.9	3.0	1.7
50	*****	******	*****	******	*****	*****	4.7	4.5	4.4	4.2	4.0	3.7	2.9	1.6
55	******	******	******	*****	*****	*****	4.4	4.3	4.2	4.0	3.9	3.5	2.7	1.6
60	******							4.1	4.0	3.8	3.7	3.4	2.6	1.5
65	*****	******	*****	*****	******	*****	*****	4.0	3.8	3.7	3.5	3.2	2.5	1.4
70	*****	*****	******	******	*****	*****	*****	3.8	3.7	3.6	3.4	3.1	2.4	1.4
75	*****	******	*****	*****	******	*****	******	*****	3.6	3.4	3.3	3.0	2.3	1.3
80	*****	*****	*****	*****	******	*****	******	*****	3.4	3.3	3.2	2.9	2.3	1.3
85	*****	******	*****	*****	******	*****	******	*****	*****	3.2	3.1	2.8	2.2	1.3
90	*****									3.1	3.0	2.7	2.1	1.2
95	*****									3.1	2.9	2.7	2.1	1.2
100	*****										2.9	2.6	2.0	1.2
125	*****											2.3	1.8	1.0
150	*****	*****	******	*****	******	*****	******	*****	******	******	*****	*****	1.6	1.0
200	*****	*****	******	*****	******	*****	******	*****	******	******	*****	******	*****	0.8
250	*****	*****	******	*****	******	******	******	******	******	*******	******	******	*****	0.7

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for QUEBEC

NUMERATOR O					İ	ESTIMATE	D PERCEN	ΓAGE						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1 2	85.8 60.7	85.4 60.4	85.0 60.1	83.7 59.2	81.4 57.6	79.1 56.0	76.8 54.3	74.3 52.6	71.8 50.8	69.2 48.9	66.5 47.0	60.7 42.9	47.0 33.2	27.1 19.2
3	*****	49.3	49.1	48.3	47.0	45.7	44.3	42.9	41.5	40.0	38.4	35.0	27.1	15.7
4	*****	42.7	42.5	41.8	40.7	39.6	38.4	37.2	35.9	34.6	33.2	30.4	23.5	13.6
5	*****	38.2	38.0	37.4	36.4	35.4	34.3	33.2	32.1	31.0	29.7	27.1	21.0	12.1
6	*****	34.9	34.7	34.2	33.2	32.3	31.3	30.4	29.3	28.3	27.1	24.8	19.2	11.1
7	*****	32.3	32.1	31.6	30.8	29.9	29.0	28.1	27.1	26.2	25.1	22.9	17.8	10.3
8	******	30.2	30.0	29.6	28.8	28.0	27.1	26.3	25.4	24.5	23.5	21.5	16.6	9.6
9	******	28.5	28.3	27.9	27.1	26.4	25.6	24.8	23.9	23.1	22.2	20.2	15.7	9.0
10 11	*****	27.0 25.8	26.9 25.6	26.5 25.2	25.8 24.6	25.0 23.9	24.3 23.2	23.5 22.4	22.7 21.7	21.9 20.9	21.0 20.0	19.2 18.3	14.9 14.2	8.6 8.2
12	*****	24.7	24.5	24.2	23.5	22.8	22.2	21.5	20.7	20.9	19.2	17.5	13.6	7.8
13	*****	23.7	23.6	23.2	22.6	22.0	21.3	20.6	19.9	19.2	18.4	16.8	13.0	7.5
14	*****	22.8	22.7	22.4	21.8	21.2	20.5	19.9	19.2	18.5	17.8	16.2	12.6	7.3
15	*****	22.1	21.9	21.6	21.0	20.4	19.8	19.2	18.5	17.9	17.2	15.7	12.1	7.0
16	******	21.4	21.2	20.9	20.4	19.8	19.2	18.6	18.0	17.3	16.6	15.2	11.8	6.8
17	******	20.7	20.6	20.3	19.8	19.2	18.6	18.0	17.4	16.8	16.1	14.7	11.4	6.6
18	*****	20.1	20.0	19.7	19.2	18.7	18.1	17.5	16.9	16.3	15.7	14.3	11.1	6.4
19	******	19.6	19.5	19.2	18.7	18.2	17.6	17.1	16.5	15.9	15.3	13.9	10.8	6.2
20	*****	19.1	19.0	18.7	18.2	17.7	17.2	16.6	16.1	15.5	14.9	13.6	10.5	6.1
21	*****	18.6	18.5	18.3	17.8	17.3	16.8	16.2	15.7	15.1	14.5	13.2	10.3	5.9
22	*****	18.2	18.1	17.8	17.4	16.9	16.4	15.9	15.3	14.8	14.2	12.9	10.0	5.8
23	******	17.8	17.7	17.4	17.0	16.5	16.0	15.5	15.0	14.4	13.9	12.7	9.8	5.7
24	*****	17.4	17.3	17.1	16.6	16.2	15.7	15.2	14.7	14.1	13.6	12.4	9.6	5.5
25	******	17.1	17.0	16.7	16.3	15.8	15.4	14.9	14.4	13.8	13.3	12.1	9.4	5.4
30 35	******		15.5 14.4	15.3	14.9	14.5	14.0	13.6	13.1	12.6	12.1	11.1	8.6 7.9	5.0
35 40	*****		14.4	14.1 13.2	13.8 12.9	13.4 12.5	13.0 12.1	12.6 11.8	12.1 11.4	11.7 10.9	11.2 10.5	10.3 9.6	7.9 7.4	4.6 4.3
45	*****		12.7	12.5	12.9	11.8	11.4	11.0	10.7	10.3	9.9	9.0	7.4	4.3
50	*****		12.0	11.8	11.5	11.2	10.9	10.5	10.7	9.8	9.4	8.6	6.6	3.8
55	*****	*****	11.5	11.3	11.0	10.7	10.4	10.0	9.7	9.3	9.0	8.2	6.3	3.7
60	*****	*****		10.8	10.5	10.2	9.9	9.6	9.3	8.9	8.6	7.8	6.1	3.5
65	*****	*****	****	10.4	10.1	9.8	9.5	9.2	8.9	8.6	8.2	7.5	5.8	3.4
70	*****	*****	*****	10.0	9.7	9.5	9.2	8.9	8.6	8.3	7.9	7.3	5.6	3.2
75	*****	*****	****	9.7	9.4	9.1	8.9	8.6	8.3	8.0	7.7	7.0	5.4	3.1
80	*****	*****	*****	9.4	9.1	8.8	8.6	8.3	8.0	7.7	7.4	6.8	5.3	3.0
85	*****			9.1	8.8	8.6	8.3	8.1	7.8	7.5	7.2	6.6	5.1	2.9
90	*****			8.8	8.6	8.3	8.1	7.8	7.6	7.3	7.0	6.4	5.0	2.9
95	*****			8.6	8.4	8.1	7.9	7.6	7.4	7.1	6.8	6.2	4.8	2.8
100	*****			8.4	8.1	7.9	7.7	7.4	7.2	6.9	6.6	6.1	4.7	2.7
125 150	*****			7.5	7.3	7.1	6.9	6.6	6.4	6.2	5.9	5.4	4.2	2.4
200	*****				6.6 5.8	6.5 5.6	6.3 5.4	6.1 5.3	5.9 5.1	5.7 4.9	5.4 4.7	5.0 4.3	3.8 3.3	2.2 1.9
250	*****				5.0	5.0	4.9	4.7	4.5	4.9	4.7	3.8	3.0	1.7
300	*****	*****	******	****		4.6	4.4	4.3	4.1	4.0	3.8	3.5	2.7	1.6
350	*****	*****	*****	****	*****	4.2	4.1	4.0	3.8	3.7	3.6	3.2	2.5	1.5
400	*****	*****	*****	*****	*****	4.0	3.8	3.7	3.6	3.5	3.3	3.0	2.4	1.4
450	*****	*****	******	*****	*****	*****	3.6	3.5	3.4	3.3	3.1	2.9	2.2	1.3
500	*****						3.4	3.3	3.2	3.1	3.0	2.7	2.1	1.2
750	******								2.6	2.5	2.4	2.2	1.7	1.0
1000	******									2.2	2.1	1.9	1.5	0.9
1500	*****												1.2	0.7
2000	*****	******	******	*****	******	*****	******	*****	******	*******	*****	*****	1.1	0.6

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for ONTARIO

NUMERATOR O						ESTIMATE!	D PERCEN	ГАСЕ						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1 2	71.0 50.2	70.7 50.0	70.3 49.7	69.2 49.0	67.4 47.7	65.5 46.3	63.5 44.9	61.5 43.5	59.4 42.0	57.3 40.5	55.0 38.9	50.2 35.5	38.9 27.5	22.5 15.9
3	41.0	40.8	40.6	40.0	38.9	37.8	36.7	35.5	34.3	33.1	31.8	29.0	22.5	13.0
4	35.5	35.3	35.2	34.6	33.7	32.7	31.8	30.8	29.7	28.6	27.5	25.1	19.5	11.2
5	******	31.6	31.5	31.0	30.1	29.3	28.4	27.5	26.6	25.6	24.6	22.5	17.4	10.0
6 7	******	28.9 26.7	28.7	28.3 26.2	27.5 25.5	26.7 24.8	25.9 24.0	25.1	24.3	23.4	22.5	20.5 19.0	15.9	9.2
/ 8	*****	25.0	26.6 24.9	26.2	25.5 23.8	24.8	24.0	23.3 21.8	22.5 21.0	21.6 20.2	20.8 19.5	19.0	14.7 13.8	8.5 7.9
9	******	23.6	23.4	23.1	22.5	21.8	21.2	20.5	19.8	19.1	18.3	16.7	13.0	7.5
10	*****	22.4	22.2	21.9	21.3	20.7	20.1	19.5	18.8	18.1	17.4	15.9	12.3	7.1
11	*****	21.3	21.2	20.9	20.3	19.7	19.2	18.5	17.9	17.3	16.6	15.1	11.7	6.8
12	******	20.4	20.3	20.0	19.5	18.9	18.3	17.8	17.2	16.5	15.9	14.5	11.2	6.5
13	******	19.6	19.5	19.2	18.7	18.2	17.6	17.1	16.5	15.9	15.3	13.9	10.8	6.2
14	*****	18.9	18.8	18.5	18.0	17.5	17.0	16.4	15.9	15.3	14.7	13.4	10.4	6.0
15	******	18.3	18.2	17.9	17.4	16.9	16.4	15.9	15.3	14.8	14.2	13.0	10.0	5.8
16	******	17.7	17.6	17.3	16.8	16.4	15.9	15.4	14.9	14.3	13.8	12.6	9.7	5.6
17 18	*****	17.1 16.7	17.1 16.6	16.8 16.3	16.3 15.9	15.9 15.4	15.4 15.0	14.9 14.5	14.4 14.0	13.9 13.5	13.3 13.0	12.2 11.8	9.4 9.2	5.4 5.3
19	*****	16.7	16.1	15.9	15.5	15.4	14.6	14.5	13.6	13.3	12.6	11.5	8.9	5.2
20	******	15.8	15.7	15.5	15.1	14.6	14.2	13.8	13.3	12.8	12.3	11.2	8.7	5.0
21	*****	15.4	15.3	15.1	14.7	14.3	13.9	13.4	13.0	12.5	12.0	11.0	8.5	4.9
22	*****	15.1	15.0	14.8	14.4	14.0	13.5	13.1	12.7	12.2	11.7	10.7	8.3	4.8
23	******	14.7	14.7	14.4	14.1	13.7	13.2	12.8	12.4	11.9	11.5	10.5	8.1	4.7
24	******	14.4	14.4	14.1	13.8	13.4	13.0	12.6	12.1	11.7	11.2	10.3	7.9	4.6
25	*****	14.1	14.1	13.8	13.5	13.1	12.7	12.3	11.9	11.5	11.0	10.0	7.8	4.5
30	******	12.9	12.8	12.6	12.3	12.0	11.6	11.2	10.9	10.5	10.0	9.2	7.1	4.1
35 40	******	11.9 11.2	11.9	11.7 10.9	11.4	11.1 10.4	10.7	10.4 9.7	10.0 9.4	9.7	9.3	8.5 7.9	6.6 6.2	3.8
45	*****		11.1 10.5	10.9	10.7 10.0	9.8	10.0 9.5	9.7	9.4 8.9	9.1 8.5	8.7 8.2	7.5	5.8	3.6 3.3
50	*****		9.9	9.8	9.5	9.3	9.0	8.7	8.4	8.1	7.8	7.3	5.5	3.2
55	*****	****	9.5	9.3	9.1	8.8	8.6	8.3	8.0	7.7	7.4	6.8	5.2	3.0
60	*****	****	9.1	8.9	8.7	8.5	8.2	7.9	7.7	7.4	7.1	6.5	5.0	2.9
65	*****		8.7	8.6	8.4	8.1	7.9	7.6	7.4	7.1	6.8	6.2	4.8	2.8
70	*****		8.4	8.3	8.1	7.8	7.6	7.4	7.1	6.8	6.6	6.0	4.7	2.7
75	*****		8.1	8.0	7.8	7.6	7.3	7.1	6.9	6.6	6.4	5.8	4.5	2.6
80	*****		7.9	7.7	7.5	7.3	7.1	6.9	6.6	6.4	6.2	5.6	4.4	2.5
85 90	******		7.6	7.5	7.3	7.1	6.9	6.7	6.4	6.2	6.0	5.4	4.2	2.4
90 95	*****			7.3 7.1	7.1 6.9	6.9 6.7	6.7 6.5	6.5 6.3	6.3 6.1	6.0 5.9	5.8 5.6	5.3 5.2	4.1 4.0	2.4 2.3
100	*****			6.9	6.7	6.5	6.4	6.2	5.9	5.7	5.5	5.0	3.9	2.2
125	*****	*****	****	6.2	6.0	5.9	5.7	5.5	5.3	5.1	4.9	4.5	3.5	2.0
150	*****	*****	*****	5.7	5.5	5.3	5.2	5.0	4.9	4.7	4.5	4.1	3.2	1.8
200	*****	*****	****	4.9	4.8	4.6	4.5	4.4	4.2	4.0	3.9	3.6	2.8	1.6
250	*****	*****	******	*****	4.3	4.1	4.0	3.9	3.8	3.6	3.5	3.2	2.5	1.4
300	*****				3.9	3.8	3.7	3.6	3.4	3.3	3.2	2.9	2.2	1.3
350	*****				3.6	3.5	3.4	3.3	3.2	3.1	2.9	2.7	2.1	1.2
400	*****				3.4	3.3	3.2	3.1	3.0	2.9	2.8	2.5	1.9	1.1
450	******					3.1	3.0	2.9	2.8	2.7	2.6	2.4	1.8	1.1
500 750	******					2.9	2.8 2.3	2.8 2.2	2.7 2.2	2.6 2.1	2.5 2.0	2.2 1.8	1.7	1.0
1000	*****							1.9	1.9	1.8	1.7	1.8	1.4 1.2	0.8 0.7
1500	*****										1.7	1.8	1.2	0.7
2000	*****											1.1	0.9	0.5
3000	******	*****	******	*****	*****	*****	*****	*****	*****	******	*****			0.4

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for MANITOBA

NUMERATOR O					İ	ESTIMATE	) PERCENT	ГАСЕ						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	*****	43.0	42.7	42.1	41.0	39.8	38.6	37.4	36.1	34.8	33.4	30.5	23.6	13.7
2	*****	30.4	30.2	29.8	29.0	28.1	27.3	26.4	25.5	24.6	23.6	21.6	16.7	9.7
3	*****	24.8	24.7	24.3	23.6	23.0	22.3	21.6	20.9	20.1	19.3	17.6	13.7	7.9
4	*****	21.5	21.4	21.0	20.5	19.9	19.3	18.7	18.1	17.4	16.7	15.3	11.8	6.8
5	*****	****	19.1	18.8	18.3	17.8	17.3	16.7	16.2	15.6	15.0	13.7	10.6	6.1
6	*****	*****	17.4	17.2	16.7	16.2	15.8	15.3	14.7	14.2	13.7	12.5	9.7	5.6
7	*****	*****	16.2	15.9	15.5	15.0	14.6	14.1	13.7	13.2	12.6	11.5	8.9	5.2
8	******	*****	15.1	14.9	14.5	14.1	13.7	13.2	12.8	12.3	11.8	10.8	8.4	4.8
9	******	*****	*****	14.0	13.7	13.3	12.9	12.5	12.0	11.6	11.1	10.2	7.9	4.6
10	******	*****	*****	13.3	13.0	12.6	12.2	11.8	11.4	11.0	10.6	9.7	7.5	4.3
11	******	*****	*****	12.7	12.3	12.0	11.6	11.3	10.9	10.5	10.1	9.2	7.1	4.1
12	******	*****	*****	12.1	11.8	11.5	11.1	10.8	10.4	10.0	9.7	8.8	6.8	3.9
13	******	*****	*****	11.7	11.4	11.0	10.7	10.4	10.0	9.7	9.3	8.5	6.6	3.8
14	******	*****	*****	11.2	10.9	10.6	10.3	10.0	9.7	9.3	8.9	8.2	6.3	3.6
15	******	*****	*****	10.9	10.6	10.3	10.0	9.7	9.3	9.0	8.6	7.9	6.1	3.5
16	******	*****	*****	10.5	10.2	10.0	9.7	9.3	9.0	8.7	8.4	7.6	5.9	3.4
17	******	*****	*****	10.2	9.9	9.7	9.4	9.1	8.8	8.4	8.1	7.4	5.7	3.3
18	******	*****	*****	9.9	9.7	9.4	9.1	8.8	8.5	8.2	7.9	7.2	5.6	3.2
19	******	*****	*****	9.7	9.4	9.1	8.9	8.6	8.3	8.0	7.7	7.0	5.4	3.1
20	******	*****	*****	9.4	9.2	8.9	8.6	8.4	8.1	7.8	7.5	6.8	5.3	3.1
21	******	*****	*****	9.2	8.9	8.7	8.4	8.2	7.9	7.6	7.3	6.7	5.2	3.0
22	******	*****	******	*****	8.7	8.5	8.2	8.0	7.7	7.4	7.1	6.5	5.0	2.9
23	******	*****	******	*****	8.5	8.3	8.1	7.8	7.5	7.3	7.0	6.4	4.9	2.8
24	******	*****	******	*****	8.4	8.1	7.9	7.6	7.4	7.1	6.8	6.2	4.8	2.8
25	*****	*****	******	*****	8.2	8.0	7.7	7.5	7.2	7.0	6.7	6.1	4.7	2.7
30	*****				7.5	7.3	7.1	6.8	6.6	6.4	6.1	5.6	4.3	2.5
35	*****				6.9	6.7	6.5	6.3	6.1	5.9	5.7	5.2	4.0	2.3
40	*****				6.5	6.3	6.1	5.9	5.7	5.5	5.3	4.8	3.7	2.2
45	*****					5.9	5.8	5.6	5.4	5.2	5.0	4.6	3.5	2.0
50	******					5.6	5.5	5.3	5.1	4.9	4.7	4.3	3.3	1.9
55	*****					5.4	5.2	5.0	4.9	4.7	4.5	4.1	3.2	1.8
60	*****					5.1	5.0	4.8	4.7	4.5	4.3	3.9	3.1	1.8
65	*****						4.8	4.6	4.5	4.3	4.1	3.8	2.9	1.7
70	******						4.6	4.5	4.3	4.2	4.0	3.6	2.8	1.6
75	*****						4.5	4.3	4.2	4.0	3.9	3.5	2.7	1.6
80	*****						4.3	4.2	4.0	3.9	3.7	3.4	2.6	1.5
85	*****							4.1	3.9	3.8	3.6	3.3	2.6	1.5
90	******							3.9	3.8	3.7	3.5	3.2	2.5	1.4
95								3.8	3.7	3.6	3.4	3.1	2.4	1.4
100	*****							3.7	3.6	3.5	3.3	3.1	2.4	1.4
125	*****								3.2	3.1	3.0	2.7	2.1	1.2
150	*****										2.7	2.5	1.9	1.1
200	*****											2.2	1.7	1.0
250	******												1.5	0.9
300 350	*******													0.8 0.7
200														0.7

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### $\label{thm:local_prox_state} \mbox{Approximate Sampling Variability Tables for SASKATCHEWAN}$

NUMERATOR O						ESTIMATE	) PERCENT	TAGE						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	******	44.3	44.1	43.4	42.2	41.1	39.8	38.6	37.3	35.9	34.5	31.5	24.4	14.1
2	*****	31.3	31.2	30.7	29.9	29.0	28.2	27.3	26.3	25.4	24.4	22.3	17.2	10.0
3	******	25.6	25.4	25.1	24.4	23.7	23.0	22.3	21.5	20.7	19.9	18.2	14.1	8.1
4	******	*****	22.0	21.7	21.1	20.5	19.9	19.3	18.6	17.9	17.2	15.7	12.2	7.0
5	******	*****	19.7	19.4	18.9	18.4	17.8	17.2	16.7	16.1	15.4	14.1	10.9	6.3
6	******	*****	18.0	17.7	17.2	16.8	16.3	15.7	15.2	14.7	14.1	12.9	10.0	5.7
7	******	*****	16.7	16.4	16.0	15.5	15.1	14.6	14.1	13.6	13.0	11.9	9.2	5.3
8	******	*****	****	15.3	14.9	14.5	14.1	13.6	13.2	12.7	12.2	11.1	8.6	5.0
9	******	*****	*****	14.5	14.1	13.7	13.3	12.9	12.4	12.0	11.5	10.5	8.1	4.7
10	******	*****	****	13.7	13.4	13.0	12.6	12.2	11.8	11.4	10.9	10.0	7.7	4.5
11	******	*****	*****	13.1	12.7	12.4	12.0	11.6	11.2	10.8	10.4	9.5	7.4	4.2
12	******	*****	****	12.5	12.2	11.9	11.5	11.1	10.8	10.4	10.0	9.1	7.0	4.1
13	******	*****	*****	12.0	11.7	11.4	11.0	10.7	10.3	10.0	9.6	8.7	6.8	3.9
14	******	*****	****	11.6	11.3	11.0	10.6	10.3	10.0	9.6	9.2	8.4	6.5	3.8
15	******	*****	*****	11.2	10.9	10.6	10.3	10.0	9.6	9.3	8.9	8.1	6.3	3.6
16	******	*****	*****	10.8	10.6	10.3	10.0	9.6	9.3	9.0	8.6	7.9	6.1	3.5
17	******	*****	*****	10.5	10.2	10.0	9.7	9.4	9.0	8.7	8.4	7.6	5.9	3.4
18	******	*****	*****	10.2	10.0	9.7	9.4	9.1	8.8	8.5	8.1	7.4	5.7	3.3
19	******	*****	****	10.0	9.7	9.4	9.1	8.8	8.5	8.2	7.9	7.2	5.6	3.2
20	******	*****	******	*****	9.4	9.2	8.9	8.6	8.3	8.0	7.7	7.0	5.5	3.1
21	******	*****	*****	*****	9.2	9.0	8.7	8.4	8.1	7.8	7.5	6.9	5.3	3.1
22	******	*****	*****	*****	9.0	8.8	8.5	8.2	7.9	7.7	7.4	6.7	5.2	3.0
23	******	*****	******	*****	8.8	8.6	8.3	8.0	7.8	7.5	7.2	6.6	5.1	2.9
24	******	*****	*****	*****	8.6	8.4	8.1	7.9	7.6	7.3	7.0	6.4	5.0	2.9
25	******	*****	*****	****	8.4	8.2	8.0	7.7	7.5	7.2	6.9	6.3	4.9	2.8
30	******	*****	*****	*****	7.7	7.5	7.3	7.0	6.8	6.6	6.3	5.7	4.5	2.6
35	******	*****	*****	****	7.1	6.9	6.7	6.5	6.3	6.1	5.8	5.3	4.1	2.4
40	******	*****	******	*****	*****	6.5	6.3	6.1	5.9	5.7	5.5	5.0	3.9	2.2
45	******	*****	*****	*****	*****	6.1	5.9	5.7	5.6	5.4	5.1	4.7	3.6	2.1
50	******	*****	******	*****	*****	5.8	5.6	5.5	5.3	5.1	4.9	4.5	3.4	2.0
55	******	*****	******	*****	*****	5.5	5.4	5.2	5.0	4.8	4.7	4.2	3.3	1.9
60	******	*****	*****	****	*****	*****	5.1	5.0	4.8	4.6	4.5	4.1	3.1	1.8
65	******	*****	******	*****	******	*****	4.9	4.8	4.6	4.5	4.3	3.9	3.0	1.7
70	******	*****	******	*****	******	*****	4.8	4.6	4.5	4.3	4.1	3.8	2.9	1.7
75	******	*****	******	*****	******	*****	4.6	4.5	4.3	4.1	4.0	3.6	2.8	1.6
80	******	*****	******	*****	******	*****	*****	4.3	4.2	4.0	3.9	3.5	2.7	1.6
85	******	*****	*****	****	*****	*****	*****	4.2	4.0	3.9	3.7	3.4	2.6	1.5
90	******	*****	******	*****	******	*****	*****	4.1	3.9	3.8	3.6	3.3	2.6	1.5
95	******	*****	******	*****	******	*****	*****	4.0	3.8	3.7	3.5	3.2	2.5	1.4
100	******	*****	******	*****	******	*****	******	*****	3.7	3.6	3.4	3.1	2.4	1.4
125	******	*****	******	****	*****	*****	*****	*****		3.2	3.1	2.8	2.2	1.3
150	******	*****	******	****	*****	*****	*****	*****	*****	*****	2.8	2.6	2.0	1.1
200	******	*****	******	*****	*****	*****	*****	*****	*****	****	*****	*****	1.7	1.0
250	*****	*****	******	*****	*****	*****	*****	*****	*****	****	*****	*****	1.5	0.9
300	*****	*****	******	*****	*****	*****	******	*****	******	*****	******	******	*****	0.8

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for ALBERTA

NUMERATOR O					[	ESTIMATEI	) PERCENT	ГАСЕ						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1 2	60.0	59.8 42.3	59.5 42.1	58.6 41.4	57.0 40.3	55.4 39.2	53.7 38.0	52.0 36.8	50.3 35.5	48.4 34.2	46.5 32.9	42.5 30.0	32.9 23.3	19.0 13.4
3	*****	34.5	34.3	33.8	32.9	32.0	31.0	30.0	29.0	28.0	26.9	24.5	19.0	11.0
4	****	29.9	29.7	29.3	28.5	27.7	26.9	26.0	25.1	24.2	23.3	21.2	16.5	9.5
5	*****	26.7	26.6	26.2	25.5	24.8	24.0	23.3	22.5	21.7	20.8	19.0	14.7	8.5
6	*****	24.4	24.3	23.9	23.3	22.6	21.9	21.2	20.5	19.8	19.0	17.3	13.4	7.8
7	*****	22.6	22.5	22.1	21.5	20.9	20.3	19.7	19.0	18.3	17.6	16.1	12.4	7.0
8	*****	21.1	21.0	20.7	20.1	19.6	19.0	18.4	17.8	17.1	16.5	15.0	11.6	6.7
9	*****	19.9	19.8	19.5	19.0	18.5	17.9	17.3	16.8	16.1	15.5	14.2	11.0	6.3
10	*****	18.9	18.8	18.5	18.0	17.5	17.0	16.5	15.9	15.3	14.7	13.4	10.4	6.0
11	*****		17.9	17.7	17.2	16.7	16.2	15.7	15.2	14.6	14.0	12.8	9.9	5.7
12	*****	****	17.2	16.9	16.5	16.0	15.5	15.0	14.5	14.0	13.4	12.3	9.5	5.5
13	*****	****	16.5	16.2	15.8	15.4	14.9	14.4	13.9	13.4	12.9	11.8	9.1	5.3
14	*****	****	15.9	15.6	15.2	14.8	14.4	13.9	13.4	12.9	12.4	11.4	8.8	5.1
15	*****	****	15.4	15.1	14.7	14.3	13.9	13.4	13.0	12.5	12.0	11.0	8.5	4.9
16	*****	****	14.9	14.6	14.2	13.8	13.4	13.0	12.6	12.1	11.6	10.6	8.2	4.7
17	*****	****	14.4	14.2	13.8	13.4	13.0	12.6	12.2	11.7	11.3	10.3	8.0	4.6
18	*****	****	14.0	13.8	13.4	13.1	12.7	12.3	11.8	11.4	11.0	10.0	7.8	4.5
19	*****	****	13.6	13.4	13.1	12.7	12.3	11.9	11.5	11.1	10.7	9.7	7.5	4.4
20	*****	****	13.3	13.1	12.7	12.4	12.0	11.6	11.2	10.8	10.4	9.5	7.4	4.2
21	*****	****	13.0	12.8	12.4	12.1	11.7	11.4	11.0	10.6	10.2	9.3	7.2	4.1
22	*****	*****	*****	12.5	12.2	11.8	11.5	11.1	10.7	10.3	9.9	9.1	7.0	4.1
23	*****	*****	*****	12.2	11.9	11.5	11.2	10.8	10.5	10.1	9.7	8.9	6.9	4.0
24	*****	*****	*****	12.0	11.6	11.3	11.0	10.6	10.3	9.9	9.5	8.7	6.7	3.9
25	*****	*****	*****	11.7	11.4	11.1	10.7	10.4	10.1	9.7	9.3	8.5	6.6	3.8
30	*****	*****	*****	10.7	10.4	10.1	9.8	9.5	9.2	8.8	8.5	7.8	6.0	3.5
35	*****	*****	*****	9.9	9.6	9.4	9.1	8.8	8.5	8.2	7.9	7.2	5.6	3.2
40	*****	****	*****	9.3	9.0	8.8	8.5	8.2	7.9	7.7	7.4	6.7	5.2	3.0
45	*****			8.7	8.5	8.3	8.0	7.8	7.5	7.2	6.9	6.3	4.9	2.8
50	*****			8.3	8.1	7.8	7.6	7.4	7.1	6.8	6.6	6.0	4.7	2.7
55	*****				7.7	7.5	7.2	7.0	6.8	6.5	6.3	5.7	4.4	2.6
60	*****				7.4	7.2	6.9	6.7	6.5	6.3	6.0	5.5	4.2	2.5
65	*****				7.1	6.9	6.7	6.5	6.2	6.0	5.8	5.3	4.1	2.4
70	*****				6.8	6.6	6.4	6.2	6.0	5.8	5.6	5.1	3.9	2.3
75	******				6.6	6.4	6.2	6.0	5.8	5.6	5.4	4.9	3.8	2.2
80	*****				6.4	6.2	6.0	5.8	5.6	5.4	5.2	4.7	3.7	2.1
85	******				6.2	6.0	5.8	5.6	5.5	5.3	5.0	4.6	3.6	2.1
90	*******				6.0	5.8	5.7	5.5	5.3	5.1	4.9	4.5	3.5	2.0
95	******				5.8	5.7	5.5	5.3	5.2	5.0	4.8	4.4	3.4	1.9
100	******				5.7	5.5	5.4	5.2	5.0	4.8	4.7	4.2	3.3	1.9
125 150	*****					5.0 4.5	4.8 4.4	4.7	4.5	4.3	4.2	3.8	2.9 2.7	1.7
	*****							4.2	4.1	4.0	3.8	3.5		1.6
200 250	******						3.8	3.7	3.6 3.2	3.4 3.1	3.3 2.9	3.0 2.7	2.3 2.1	1.3 1.2
300	*****								2.9	2.8	2.9	2.7	1.9	1.2
350	*****									2.6	2.7	2.3	1.8	1.1
400	*****										2.3	2.3	1.6	0.9
450	*****											2.1	1.6	0.9
500	*****	****	*****	*****	******	*****	*****	*****	*****	*****	*****	1.9	1.5	0.8
750	*****	*****	******	*****	******	*****	*****	*****	******	******	*****		1.2	0.7

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for BRITISH COLUMBIA

NUMERATOR O					[	ESTIMATE	D PERCEN	ΓAGE						
PERCENTAGE	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
( 000)	0.1%	1.0%	2.0%	5.0%	10.0%	13.0%	20.0%	23.0%	30.0%	33.0%	40.0%	30.0%	70.0%	JU. U/A
1	78.3	78.0	77.6	76.4	74.3	72.3	70.1	67.9	65.6	63.2	60.7	55.4	42.9	24.8
2	******	55.1	54.9	54.0	52.6	51.1	49.6	48.0	46.4	44.7	42.9	39.2	30.4	17.5
3	******	45.0	44.8	44.1	42.9	41.7	40.5	39.2	37.9	36.5	35.0	32.0	24.8	14.3
4	******	39.0	38.8	38.2	37.2	36.1	35.0	33.9	32.8	31.6	30.4	27.7	21.5	12.4
5	******	34.9	34.7	34.2	33.2	32.3	31.3	30.4	29.3	28.3	27.1	24.8	19.2	11.1
6	******	31.8	31.7	31.2	30.4	29.5	28.6	27.7	26.8	25.8	24.8	22.6	17.5	10.1
7	******	29.5	29.3	28.9	28.1	27.3	26.5	25.7	24.8	23.9	22.9	20.9	16.2	9.4
8	******	27.6	27.4	27.0	26.3	25.5	24.8	24.0	23.2	22.3	21.5	19.6	15.2	8.8
9	******	26.0	25.9	25.5	24.8	24.1	23.4	22.6	21.9	21.1	20.2	18.5	14.3	8.3
10	******	24.7	24.5	24.2	23.5	22.8	22.2	21.5	20.7	20.0	19.2	17.5	13.6	7.8
11	******	23.5	23.4	23.0	22.4	21.8	21.1	20.5	19.8	19.0	18.3	16.7	12.9	7.5
12	******	22.5	22.4	22.0	21.5	20.9	20.2	19.6	18.9	18.2	17.5	16.0	12.4	7.2
13	******	21.6	21.5	21.2	20.6	20.0	19.4	18.8	18.2	17.5	16.8	15.4	11.9	6.9
14	******	20.8	20.7	20.4	19.9	19.3	18.7	18.1	17.5	16.9	16.2	14.8	11.5	6.6
15	******	20.1	20.0	19.7	19.2	18.7	18.1	17.5	16.9	16.3	15.7	14.3	11.1	6.4
16	******	*****	19.4	19.1	18.6	18.1	17.5	17.0	16.4	15.8	15.2	13.9	10.7	6.2
17	******	*****	18.8	18.5	18.0	17.5	17.0	16.5	15.9	15.3	14.7	13.4	10.4	6.0
18	******	*****	18.3	18.0	17.5	17.0	16.5	16.0	15.5	14.9	14.3	13.1	10.1	5.8
19	******	*****	17.8	17.5	17.1	16.6	16.1	15.6	15.0	14.5	13.9	12.7	9.8	5.7
20	******	*****	17.3	17.1	16.6	16.2	15.7	15.2	14.7	14.1	13.6	12.4	9.6	5.5
21	******	*****	16.9	16.7	16.2	15.8	15.3	14.8	14.3	13.8	13.2	12.1	9.4	5.4
22	******	*****	16.5	16.3	15.9	15.4	14.9	14.5	14.0	13.5	12.9	11.8	9.2	5.3
23	******	*****	16.2	15.9	15.5	15.1	14.6	14.2	13.7	13.2	12.7	11.6	9.0	5.2
24	******	*****	15.8	15.6	15.2	14.7	14.3	13.9	13.4	12.9	12.4	11.3	8.8	5.1
25	******	*****	15.5	15.3	14.9	14.5	14.0	13.6	13.1	12.6	12.1	11.1	8.6	5.0
30	******	*****	14.2	13.9	13.6	13.2	12.8	12.4	12.0	11.5	11.1	10.1	7.8	4.5
35	******	******	*****	12.9	12.6	12.2	11.8	11.5	11.1	10.7	10.3	9.4	7.3	4.2
40	******	******	*****	12.1	11.8	11.4	11.1	10.7	10.4	10.0	9.6	8.8	6.8	3.9
45	******	******	*****	11.4	11.1	10.8	10.4	10.1	9.8	9.4	9.0	8.3	6.4	3.7
50	******	******	*****	10.8	10.5	10.2	9.9	9.6	9.3	8.9	8.6	7.8	6.1	3.5
55	******	******	*****	10.3	10.0	9.7	9.5	9.2	8.8	8.5	8.2	7.5	5.8	3.3
60	******	******	*****	9.9	9.6	9.3	9.0	8.8	8.5	8.2	7.8	7.2	5.5	3.2
65	******	******	*****	9.5	9.2	9.0	8.7	8.4	8.1	7.8	7.5	6.9	5.3	3.1
70	******	******	*****	9.1	8.9	8.6	8.4	8.1	7.8	7.6	7.3	6.6	5.1	3.0
75	******	*****	****	8.8	8.6	8.3	8.1	7.8	7.6	7.3	7.0	6.4	5.0	2.9
80	******	******	******	*****	8.3	8.1	7.8	7.6	7.3	7.1	6.8	6.2	4.8	2.8
85	******	*****	*****	*****	8.1	7.8	7.6	7.4	7.1	6.9	6.6	6.0	4.7	2.7
90	******	*****	*****	*****	7.8	7.6	7.4	7.2	6.9	6.7	6.4	5.8	4.5	2.6
95	******	******	******	*****	7.6	7.4	7.2	7.0	6.7	6.5	6.2	5.7	4.4	2.5
100	******	*****	*****	*****	7.4	7.2	7.0	6.8	6.6	6.3	6.1	5.5	4.3	2.5
125	******	*****	*****	*****	6.6	6.5	6.3	6.1	5.9	5.7	5.4	5.0	3.8	2.2
150	******	******	******	*****	6.1	5.9	5.7	5.5	5.4	5.2	5.0	4.5	3.5	2.0
200	******	*****	*****	*****	*****	5.1	5.0	4.8	4.6	4.5	4.3	3.9	3.0	1.8
250	******	*****	******	*****	******		4.4	4.3	4.1	4.0	3.8	3.5	2.7	1.6
300	******	*****	*****	*****	******	*****	4.0	3.9	3.8	3.6	3.5	3.2	2.5	1.4
350	*****	*****	******	*****	******	*****		3.6	3.5	3.4	3.2	3.0	2.3	1.3
400	*****	*****	******	*****	******	*****	*****		3.3	3.2	3.0	2.8	2.1	1.2
450	*****	*****	******	*****	******	*****	*****	*****	3.1	3.0	2.9	2.6	2.0	1.2
500	*****	*****	******	*****	******	*****	*****	*****		2.8	2.7	2.5	1.9	1.1
750	*****	*****	******	*****	******	*****	*****	*****	*****			2.0	1.6	0.9
1000	*****	******	******	*****	******	*****	*****	*****	******	******	*****		1.4	0.8

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for ATLANTIC

NUMERATOR O	F				I	ESTIMATE	) PERCEN	ΓAGE						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	*****	40.4	40.2	39.6	38.5	37.4	36.3	35.2	34.0	32.7	31.5	28.7	22.2	12.8
2	*****	28.6	28.4	28.0	27.2	26.5	25.7	24.9	24.0	23.1	22.2	20.3	15.7	9.1
3	******	23.3	23.2	22.9	22.2	21.6	21.0	20.3	19.6	18.9	18.2	16.6	12.8	7.4
4	*****	20.2	20.1	19.8	19.3	18.7	18.2	17.6	17.0	16.4	15.7	14.4	11.1	6.4
5	*****	18.1	18.0	17.7	17.2	16.7	16.2	15.7	15.2	14.6	14.1	12.8	9.9	5.7
6	*****	16.5	16.4	16.2	15.7	15.3	14.8	14.4	13.9	13.4	12.8	11.7	9.1	5.2
7	*****	15.3	15.2	15.0	14.6	14.1	13.7	13.3	12.8	12.4	11.9	10.9	8.4	4.9
8	*****	14.3	14.2	14.0	13.6	13.2	12.8	12.4	12.0	11.6	11.1	10.2	7.9	4.5
9	*****		13.4	13.2	12.8	12.5	12.1	11.7	11.3	10.9	10.5	9.6	7.4	4.3
10	*****		12.7	12.5	12.2	11.8	11.5	11.1	10.7	10.4	9.9	9.1	7.0	4.1
11	*****	*****	12.1	11.9	11.6	11.3	11.0	10.6	10.7	9.9	9.5	8.7	6.7	3.9
12	*****	*****	11.6	11.4	11.1	10.8	10.5	10.0	9.8	9.5	9.1	8.3	6.4	3.7
13	*****	*****	11.1	11.0	10.7	10.4	10.3	9.8	9.4	9.1	8.7	8.0	6.2	3.6
14	*****	*****	10.7	10.6	10.7	10.0	9.7	9.4	9.1	8.7	8.4	7.7	5.9	3.4
15	*****		10.7	10.0	9.9	9.7	9.4	9.1	8.8	8.5	8.1	7.4	5.7	3.3
16	*****	*****	10.4	9.9	9.6	9.4	9.1	8.8	8.5	8.2	7.9	7.2	5.6	3.2
17	*****		9.7	9.6	9.3	9.1	8.8	8.5	8.2	7.9	7.6	7.0	5.4	3.1
18	*****			9.3	9.1	8.8	8.6	8.3	8.0	7.7	7.4	6.8	5.2	3.0
19	*****			9.1	8.8	8.6	8.3	8.1	7.8	7.5	7.2	6.6	5.1	2.9
20	*****			8.8	8.6	8.4	8.1	7.9	7.6	7.3	7.0	6.4	5.0	2.9
21	*****	******	****	8.6	8.4	8.2	7.9	7.7	7.4	7.1	6.9	6.3	4.9	2.8
22	*****			8.4	8.2	8.0	7.7	7.5	7.4	7.1	6.7	6.1	4.7	2.7
23	*****			8.3	8.0	7.8	7.6	7.3	7.1	6.8	6.6	6.0	4.6	2.7
24	*****			8.1	7.9	7.6	7.4	7.3	6.9	6.7	6.4	5.9	4.5	2.6
25	*****			7.9	7.7	7.5	7.4	7.2	6.8	6.5	6.3	5.7	4.4	2.6
30	*****	******	*****	7.2	7.0	6.8	6.6	6.4	6.2	6.0	5.7	5.2	4.1	2.3
35	*****			6.7	6.5	6.3	6.1	5.9	5.7	5.5	5.3	4.9	3.8	2.2
40	*****	*****	*****	6.3	6.1	5.9	5.7	5.6	5.4	5.2	5.0	4.5	3.5	2.0
45	*****	******	*****		5.7	5.6	5.4	5.2	5.1	4.9	4.7	4.3	3.3	1.9
50	*****	******	*****	*****	5.4	5.3	5.1	5.0	4.8	4.6	4.4	4.1	3.1	1.8
55	*****				5.2	5.0	4.9	4.7	4.6	4.4	4.2	3.9	3.1	1.7
60	*****				5.0	4.8	4.7	4.5	4.4	4.2	4.1	3.7	2.9	1.7
65	*****	******	*****	*****	4.8	4.6	4.5	4.4	4.2	4.1	3.9	3.6	2.8	1.6
70	*****	******	*****	*****	4.6	4.5	4.3	4.2	4.1	3.9	3.8	3.4	2.7	1.5
75	*****	******	*****	*****	4.4	4.3	4.2	4.1	3.9	3.8	3.6	3.3	2.6	1.5
80	*****	*****	*****	*****	4.3	4.2	4.1	3.9	3.8	3.7	3.5	3.2	2.5	1.4
85	*****	*****	*****	*****	4.2	4.1	3.9	3.8	3.7	3.6	3.4	3.1	2.4	1.4
90	*****	*****	*****	****		3.9	3.8	3.7	3.6	3.5	3.3	3.0	2.3	1.4
95	*****	*****	*****	****	*****	3.8	3.7	3.6	3.5	3.4	3.2	2.9	2.3	1.3
100	*****	*****	*****	****	*****	3.7	3.6	3.5	3.4	3.3	3.1	2.9	2.2	1.3
125	*****	*****	*****	****	*****	3.3	3.2	3.1	3.0	2.9	2.8	2.6	2.0	1.1
150	*****	*****	*****	****	*****		3.0	2.9	2.8	2.7	2.6	2.3	1.8	1.0
200	*****	*****	*****	*****	*****	*****		2.5	2.4	2.3	2.2	2.0	1.6	0.9
250	*****	*****	*****	*****	*****	*****	*****		2.1	2.1	2.0	1.8	1.4	0.8
300	*****	*****	*****	****	*****	*****	*****	*****		1.9	1.8	1.7	1.3	0.7
350	*****	*****	*****	*****	*****	*****	*****	*****	******		1.7	1.5	1.2	0.7
400	*****	*****	*****	****	*****	*****	*****	*****	******	******		1.4	1.1	0.6
450	*****	*****	*****	****	*****	*****	*****	*****	******	******	*****		1.0	0.6
500	*****	*****	*****	*****	*****	*****	*****	*****	******	******	*****	*****	1.0	0.6
750	*****	******	******	*****	*****	*****	*****	******	******	******	*****	*****		0.5

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### Approximate Sampling Variability Tables for PRAIRIES

NUMERATOR O						ESTIMATE	D PERCEN	ГАСЕ						
PERCENTAGE ('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1 2	49.3	49.1 34.7	48.9 34.5	48.1 34.0	46.8 33.1	45.5 32.2	44.1 31.2	42.7 30.2	41.3 29.2	39.8 28.1	38.2 27.0	34.9 24.7	27.0 19.1	15.6 11.0
3	*****	28.4	28.2	27.8	27.0	26.3	25.5	24.7	23.8	23.0	22.1	20.1	15.6	9.0
4	*****	24.6	24.4	24.1	23.4	22.8	22.1	21.4	20.6	19.9	19.1	17.4	13.5	7.8
5	*****	22.0	21.8	21.5	20.9	20.3	19.7	19.1	18.5	17.8	17.1	15.6	12.1	7.0
6	******	20.0	19.9	19.6	19.1	18.6	18.0	17.4	16.9	16.2	15.6	14.2	11.0	6.4
7	******	18.6	18.5	18.2	17.7	17.2	16.7	16.2	15.6	15.0	14.4	13.2	10.2	5.9
8	******	17.4	17.3	17.0	16.6	16.1	15.6	15.1	14.6	14.1	13.5	12.3	9.6	5.5
9	*****	16.4	16.3	16.0	15.6	15.2	14.7	14.2	13.8	13.3	12.7	11.6	9.0	5.2
10	*****	15.5	15.5	15.2	14.8	14.4	14.0	13.5	13.1	12.6	12.1	11.0	8.5	4.9
11 12	*****	14.8 14.2	14.7 14.1	14.5 13.9	14.1 13.5	13.7 13.1	13.3 12.7	12.9 12.3	12.4 11.9	12.0 11.5	11.5 11.0	10.5 10.1	8.2 7.8	4.7
13	*****	14.2	14.1	13.9	13.5	12.6	12.7	11.9	11.9	11.5	10.6	9.7	7.6 7.5	4.5 4.3
14	*****	13.1	13.1	12.9	12.5	12.0	11.8	11.4	11.0	10.6	10.0	9.3	7.3	4.3
15	*****	12.7	12.6	12.4	12.1	11.7	11.4	11.0	10.7	10.3	9.9	9.0	7.2	4.0
16	*****	12.3	12.2	12.0	11.7	11.4	11.0	10.7	10.7	9.9	9.6	8.7	6.8	3.9
17	*****	11.9	11.8	11.7	11.4	11.0	10.7	10.4	10.0	9.7	9.3	8.5	6.6	3.8
18	*****	11.6	11.5	11.3	11.0	10.7	10.4	10.1	9.7	9.4	9.0	8.2	6.4	3.7
19	*****	*****	11.2	11.0	10.7	10.4	10.1	9.8	9.5	9.1	8.8	8.0	6.2	3.6
20	******	*****	10.9	10.8	10.5	10.2	9.9	9.6	9.2	8.9	8.5	7.8	6.0	3.5
21	*****	*****	10.7	10.5	10.2	9.9	9.6	9.3	9.0	8.7	8.3	7.6	5.9	3.4
22	******	*****	10.4	10.3	10.0	9.7	9.4	9.1	8.8	8.5	8.2	7.4	5.8	3.3
23	******	*****	10.2	10.0	9.8	9.5	9.2	8.9	8.6	8.3	8.0	7.3	5.6	3.3
24	******		10.0	9.8	9.6	9.3	9.0	8.7	8.4	8.1	7.8	7.1	5.5	3.2
25	*****	*****	9.8	9.6	9.4	9.1	8.8	8.5	8.3	8.0	7.6	7.0	5.4	3.1
30	*****		8.9	8.8	8.5	8.3	8.1	7.8	7.5	7.3	7.0	6.4	4.9	2.8
35	******		8.3	8.1	7.9	7.7	7.5	7.2	7.0	6.7	6.5	5.9	4.6	2.6
40	*****			7.6	7.4	7.2	7.0	6.8	6.5	6.3	6.0	5.5	4.3	2.5
45	*****			7.2	7.0	6.8	6.6	6.4	6.2	5.9	5.7	5.2	4.0	2.3
50	*****			6.8	6.6	6.4	6.2	6.0	5.8	5.6	5.4	4.9	3.8	2.2
55	******			6.5	6.3	6.1	6.0	5.8	5.6	5.4	5.2	4.7	3.6	2.1
60	******			6.2	6.0	5.9	5.7	5.5	5.3	5.1	4.9	4.5	3.5	2.0
65 70	******			6.0	5.8	5.6	5.5	5.3	5.1	4.9	4.7	4.3	3.4	1.9
70 75	*****			5.7 5.6	5.6 5.4	5.4 5.3	5.3 5.1	5.1 4.9	4.9 4.8	4.8	4.6	4.2 4.0	3.2 3.1	1.9 1.8
80	*****			5.4	5.4	5.3	4.9	4.9	4.6	4.6 4.4	4.4 4.3	3.9	3.1	1.7
85	*****			5.4	5.1	4.9	4.8	4.6	4.5	4.4	4.1	3.8	2.9	1.7
90	*****			5.1	4.9	4.8	4.7	4.5	4.4	4.2	4.1	3.7	2.8	1.6
95	*****	*****	*****		4.8	4.7	4.5	4.4	4.2	4.1	3.9	3.6	2.8	1.6
100	*****	*****	*****	*****	4.7	4.6	4.4	4.3	4.1	4.0	3.8	3.5	2.7	1.6
125	*****	*****	*****	*****	4.2	4.1	3.9	3.8	3.7	3.6	3.4	3.1	2.4	1.4
150	******	******	******	*****	3.8	3.7	3.6	3.5	3.4	3.2	3.1	2.8	2.2	1.3
200	*****	******	******	*****	*****	3.2	3.1	3.0	2.9	2.8	2.7	2.5	1.9	1.1
250	******	*****	******	*****	*****	2.9	2.8	2.7	2.6	2.5	2.4	2.2	1.7	1.0
300	******	******	******	*****	*****	*****	2.5	2.5	2.4	2.3	2.2	2.0	1.6	0.9
350	******						2.4	2.3	2.2	2.1	2.0	1.9	1.4	0.8
400	******							2.1	2.1	2.0	1.9	1.7	1.4	0.8
450	*****							2.0	1.9	1.9	1.8	1.6	1.3	0.7
500	******								1.8	1.8	1.7	1.6	1.2	0.7
750	*****										1.4	1.3	1.0	0.6
1000	******												0.9	0.5
1500	******	*****	******	****	******	*****	*****	*****	*****	****	*****	*****	*****	0.4

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

### RESIDENTIAL TELEPHONE SERVICES SURVEY - 0599 (re-weighted) Approximate Sampling Variability Tables for CANADA

NUMERATOR C					I	ESTIMATE	) PERCEN	ΓAGE						
('000)	0.1%	1.0%	2.0%	5.0%	10.0%	15.0%	20.0%	25.0%	30.0%	35.0%	40.0%	50.0%	70.0%	90.0%
1	74.5	74.1	73.8	72.6	70.7	68.7	66.6	64.5	62.3	60.1	57.7	52.7	40.8	23.6
2	52.7	52.4	52.2	51.3	50.0	48.6	47.1	45.6	44.1	42.5	40.8	37.3	28.9	16.7
3	43.0	42.8	42.6	41.9	40.8	39.7	38.5	37.3	36.0	34.7	33.3	30.4	23.6	13.6
4	37.2	37.1	36.9	36.3	35.3	34.3	33.3	32.3	31.2	30.0	28.9	26.3	20.4	11.8
5	33.3	33.2	33.0	32.5	31.6	30.7	29.8	28.9	27.9	26.9	25.8	23.6	18.2	10.5
6	30.4	30.3	30.1	29.6	28.9	28.0	27.2	26.3	25.4	24.5	23.6	21.5	16.7	9.6
7	28.1	28.0	27.9	27.4	26.7	26.0	25.2	24.4	23.6	22.7	21.8	19.9	15.4	8.9
8	26.3	26.2	26.1	25.7	25.0	24.3	23.6	22.8	22.0	21.2	20.4	18.6	14.4	8.3
9	24.8	24.7	24.6	24.2	23.6	22.9	22.2	21.5	20.8	20.0	19.2	17.6	13.6	7.9
10	23.5	23.4	23.3	23.0	22.4	21.7	21.1	20.4	19.7	19.0	18.2	16.7	12.9	7.5
11	22.5	22.4	22.2	21.9	21.3	20.7	20.1	19.5	18.8	18.1	17.4	15.9	12.3	7.1
12	*****	21.4	21.3	21.0	20.4	19.8	19.2	18.6	18.0	17.3	16.7	15.2	11.8	6.8
13	******	20.6	20.5	20.1	19.6	19.1	18.5	17.9	17.3	16.7	16.0	14.6	11.3	6.5
14	******	19.8	19.7	19.4	18.9	18.4	17.8	17.2	16.7	16.1	15.4	14.1	10.9	6.3
15	******	19.1	19.0	18.7	18.2	17.7	17.2	16.7	16.1	15.5	14.9	13.6	10.5	6.1
16	******	18.5	18.4	18.2	17.7	17.2	16.7	16.1	15.6	15.0	14.4	13.2	10.2	5.9
17	*****	18.0	17.9	17.6	17.1	16.7	16.2	15.6	15.1	14.6	14.0	12.8	9.9	5.7
18	*******	17.5	17.4	17.1	16.7	16.2	15.7	15.2	14.7	14.2	13.6	12.4	9.6	5.6
19		17.0	16.9	16.7	16.2	15.8	15.3	14.8	14.3	13.8	13.2	12.1	9.4	5.4
20	******	16.6	16.5	16.2	15.8	15.4	14.9	14.4	13.9	13.4	12.9	11.8	9.1	5.3
21	*****	16.2	16.1	15.8	15.4	15.0	14.5	14.1	13.6	13.1	12.6	11.5	8.9	5.1
22	*****	15.8	15.7	15.5	15.1	14.6	14.2	13.8	13.3	12.8	12.3	11.2	8.7	5.0
23	*****	15.5	15.4	15.1	14.7	14.3	13.9	13.5	13.0	12.5	12.0	11.0	8.5	4.9
24	*****	15.1	15.1	14.8	14.4	14.0	13.6	13.2	12.7	12.3	11.8	10.8	8.3	4.8
25	*****	14.8	14.8	14.5	14.1	13.7	13.3	12.9	12.5	12.0	11.5	10.5	8.2	4.7
30	*****	13.5	13.5	13.3	12.9	12.5	12.2	11.8	11.4	11.0	10.5	9.6	7.5	4.3
35	*****	12.5	12.5	12.3	11.9	11.6	11.3	10.9	10.5	10.2	9.8	8.9	6.9	4.0
40	*****	11.7	11.7	11.5	11.2	10.9	10.5	10.2	9.9	9.5	9.1	8.3	6.5	3.7
45	*****	11.1	11.0	10.8	10.5	10.2	9.9	9.6	9.3	9.0	8.6	7.9	6.1	3.5
50	*****	10.5	10.4	10.3	10.0	9.7	9.4	9.1	8.8	8.5	8.2	7.5	5.8	3.3
55	*****	10.0	9.9	9.8	9.5	9.3	9.0	8.7	8.4	8.1	7.8	7.1	5.5	3.2
60 65	*****	9.6	9.5	9.4	9.1	8.9	8.6	8.3	8.0	7.8	7.5	6.8	5.3	3.0
70	*****	9.2 8.9	9.1 8.8	9.0 8.7	8.8 8.4	8.5 8.2	8.3 8.0	8.0 7.7	7.7 7.5	7.5 7.2	7.2 6.9	6.5 6.3	5.1 4.9	2.9 2.8
75	*****	8.6	8.5	8.4	8.2	7.9	7.7	7.7	7.3	6.9	6.7	6.1	4.9	2.7
80	*****	8.3	8.2	8.1	7.9	7.9	7.7	7.3	7.2	6.7	6.5	5.9	4.7	2.7
85	*****	8.0	8.0	7.9	7.7	7.7	7.3	7.2	6.8	6.5	6.3	5.7	4.4	2.6
90	*****	7.8	7.8	7.3	7.7	7.3	7.2	6.8	6.6	6.3	6.1	5.6	4.4	2.5
95	*****	7.6	7.6	7.5	7.3	7.0	6.8	6.6	6.4	6.2	5.9	5.4	4.2	2.4
100	*****	7.4	7.4	7.3	7.1	6.9	6.7	6.5	6.2	6.0	5.8	5.3	4.1	2.4
125	******		6.6	6.5	6.3	6.1	6.0	5.8	5.6	5.4	5.2	4.7	3.6	2.1
150	******	*****	6.0	5.9	5.8	5.6	5.4	5.3	5.1	4.9	4.7	4.3	3.3	1.9
200	******	*****	5.2	5.1	5.0	4.9	4.7	4.6	4.4	4.2	4.1	3.7	2.9	1.7
250	******			4.6	4.5	4.3	4.2	4.1	3.9	3.8	3.6	3.3	2.6	1.5
300	*****	*****	*****	4.2	4.1	4.0	3.8	3.7	3.6	3.5	3.3	3.0	2.4	1.4
350	*****	*****	*****	3.9	3.8	3.7	3.6	3.4	3.3	3.2	3.1	2.8	2.2	1.3
400	******	*****	*****	3.6	3.5	3.4	3.3	3.2	3.1	3.0	2.9	2.6	2.0	1.2
450	******	*****	*****	3.4	3.3	3.2	3.1	3.0	2.9	2.8	2.7	2.5	1.9	1.1
500	******	*****	*****	3.2	3.2	3.1	3.0	2.9	2.8	2.7	2.6	2.4	1.8	1.1
750	******	*****	******	*****	2.6	2.5	2.4	2.4	2.3	2.2	2.1	1.9	1.5	0.9
1000	******	*****	*****	*****	2.2	2.2	2.1	2.0	2.0	1.9	1.8	1.7	1.3	0.7
1500	******	*****	*****	****	*****	1.8	1.7	1.7	1.6	1.6	1.5	1.4	1.1	0.6
2000	******						1.5	1.4	1.4	1.3	1.3	1.2	0.9	0.5
3000	******								1.1	1.1	1.1	1.0	0.7	0.4
4000	******									0.9	0.9	0.8	0.6	0.4
5000	******											0.7	0.6	0.3
6000	******												0.5	0.3
7000	******												0.5	0.3
8000	******												0.5	0.3
9000	******													0.2
10000	******	*****	*****	****	*****	*****	******	*****	*****	******	*****	*****	*****	0.2

NOTE: FOR CORRECT USAGE OF THESE TABLES PLEASE REFER TO MICRODATA DOCUMENTATION

## 12.0 Weighting

Since the Residential Telephone Service Survey used a sub-sample of the LFS sample, the derivation of weights for the survey records is clearly tied to the weighting procedure used for the LFS. The LFS weighting procedure is briefly described below.

## 12.1

# Weighting Procedures for the LFS

In the LFS, the final weight attached to each record is the product of the following factors: the basic weight, the cluster sub-weight, the balancing factor for non-response, and the province-age-sex ratio adjustment factor. Each is described below.

#### **Basic Weight**

In a probability sample, the sample design itself determines weights which must be used to produce unbiased estimates of the population. Each record must be weighted by the inverse of the probability of selecting the person to whom the record refers. In the example of a 2% simple random sample, this probability would be .02 for each person and the records must be weighted by 1/.02=50. Because all eligible individuals in a dwelling are interviewed (directly or by proxy), this probability is essentially the same as the probability with which the dwelling is selected.

### Cluster Sub-weight

The cluster delineation is such that the number of dwellings in the sample increases very slightly with moderate growth in the housing stock. Substantial growth can be tolerated in an isolated cluster before the additional sample represents a field collection problem. However, if growth takes place in more than one cluster in an interviewer assignment, the cumulative effect of all increases may create a workload problem. In clusters where substantial growth has taken place, sub-sampling is used as a means of keeping interviewer assignments manageable. The cluster sub-weight represents the inverse of this sub-sampling ratio in clusters where sub-sampling has occurred.

#### Non-response

Notwithstanding the strict controls of the LFS, some non-response is inevitable, despite all the efforts made by the interviewers. The LFS non-response rate is approximately 5%. For certain types of non-response (eg. household temporarily absent, refusal), data from a previous month's interview with the household if any, is brought forward and used as the current month's data for the household.

In other cases, non-response is compensated for by proportionally increasing the weights of responding households. The weight of each responding record is increased by the ratio of the number of households that should have been interviewed, divided by the number that were actually interviewed. This adjustment is done separately for non-response areas, which are defined by employment insurance region, type of area, and rotation group. It is based on the assumption that the households that have been interviewed represent the characteristics of those that should have been interviewed. To the extent that this assumption is not true, the estimates will be somewhat biased.

#### LFS Sub-Weight

The product of the previously described weighting factors is called the LFS sub-weight. All members of the same sampled dwelling have the same sub-weight. Therefore, when calculating a household subweight, we use the subweight of one record (or person) from the household.

#### Sub-provincial and Province-Age-Sex Adjustments

The sub-weight can be used to derive a valid estimate of any characteristic for which information is collected by the LFS. In particular, estimates are produced of the total number of persons 15+ in provincial economic regions and the 24 large metropolitan areas as well as of designated age-sex groups in each of the ten provinces.

Independent estimates are available monthly for various age and sex groups by province. These are population projections based on the most recent Census data, records of births and deaths, and estimates of migration. In the final step, this auxiliary information is used to transform the sub-weight into the final weight. This is done using a calibration method. This method ensures that the final weights it produces sum to the census projections for the auxiliary variables, namely various age-sex groups, economic regions, census metropolitan areas, and rotation groups.

This weighting procedure ensures consistency with external Census counts and that each rotation group is representative of the population, and also ensures that every member of the economic family is assigned the same weight.

### 12.2

# Weighting Procedures for the Residential Telephone Service Survey

The principles behind the calculation of the weights for the Residential Telephone Service Survey are nearly identical to those for the LFS. However, this survey is a household-weighted survey, not a person-weighted survey. Also, further adjustments are made to the LFS weights in order to derive a final weight for the individual records on the Residential Telephone Service Survey microdata file.

- (1) An adjustment to account for the use of a five-sixths sub-sample, instead of the full LFS sample.
- (2) An adjustment to account for the additional non-response to the supplementary survey, i.e., non-response to the Residential Telephone Service Survey for individuals who did respond to the LFS or for which previous month's LFS data was brought forward.
- (3) A readjustment to account for independent province-stratum projections, after the above adjustments are made. These province-stratum totals are simply the final weighted province-stratum totals from the LFS. Note that a stratum roughly corresponds to an EIR-ER region (described in section 5.2.2).

Adjustments (1) and (2) are taken into account by multiplying the LFS subweight for each responding Residential Telephone Service Survey record by:

sum of LFS subweights from each household responding to LFS subweights from each household responding to the RTSS

to obtain a non-response adjusted Residential Telephone Service Survey sub-weight (WEIGHT1).

Adjustment (3) is calculated by multiplying WEIGHT1 for each Residential Telephone Service Survey respondent by:

population total for province-stratum i

sum of WEIGHT1 for survey respondents in province-stratum i

to give the resulting weight (FINWT), which is the final weight which appears on the Residential Telephone Service Survey microdata file.

#### **Calibration Estimation Adjustments**

The weights for each respondent were adjusted in Adjustment 3 by an iterative process using a calibrated estimation procedure. This procedure ensured that estimates produced for a province-stratum group would agree with the population totals for that province-stratum group. This adjustment was made by using a two-stage iterative weighting procedure, each time using the weight obtained from the previous step, until the set of estimates agreed with the LFS population totals (which were created using Census population projections). The final statistical weight can be found in the "WEIGHT" field on the microdata file. Note that this field has a decimal and should be read as (99999V9999) where V represents the location of the decimal place.

## 13.0 Questionnaires and Code Sheets

The Residential Telephone Service Survey questionnaire was used in May 1999 to collect the information for the supplementary survey.

#### **RTS-I01**

This is a voluntary survey on telephone service. Press enter to continue.

Universe: All respondents

#### RTS-Q01

How many different telephone numbers are there for your residence? (Include cellular phone numbers and phone numbers used for business). %RTS->LFSMAILAD%

<-1>	Don't know go to RTS-Q08
<-2>	Refused go to RTS-Q08
<1>	0 go to RTS-Q02
<2>	1 go to RTS-Q08
<3>	2 go to RTS-Q08
<4>	3 or more go to RTS-Q08
Note:	F1 HELP: Include: phone numbers used for businesses even if the business
	is not within the residence or if the employer is paying for the person's phone
	service within that person's household. This includes cell phones from work

that are brought home. Exclude pagers.

Universe: All respondents

#### RTS-Q02

Why don't you have a phone? INTERVIEWER: Do not read the categories. Mark one only. Press F1 for Help.

<-1>	Don't know go to RTS-Q05
<-2>	Refused go to RTS-Q05
<1>	I cancelled it go to RTS-Q03
<2>	Can't afford it go to RTS-Q04
<3>	Moved go to RTS-Q05
<4>	All other responses go to RTS-Q05
Note:	F1 NOTES: The following answers should be coded to "Can't afford it": Too expensive, Can't afford the phone, Unpaid phone bills, Service cancelled by the phone company, Moved and can't afford the installation price, Lost job, Unemployed, I'm on (any form of social assistance), etc. The following answers should be coded to "Moved": Respondent moved, on vacation, going south for the winter, moved and don't want/need the phone anymore etc. The following answers should be coded to "Any other response": Sharing a phone, receiving harassing calls, getting a private number, hard of hearing, dissatisfied with the phone company, don't want other household members to receive calls, etc.

Universe: Respondents who do not have phones RTS-Q01=1

#### **RTS-Q03**

Why did you cancel it? INTERVIEWER: Do not read categories. Press F1 for help.

<-1>	Don't know go to RTS-Q05
<-2>	Refused go to RTS-Q05
<1>	Can't afford it
<2>	Moved go to RTS-Q05
<3>	All other responses go to RTS-Q05
Note:	F1 NOTES: The following answers should be coded to "Can't afford it": Too expensive, Can't afford the phone, Unpaid phone bills, Service cancelled by the phone company, Moved and can't afford the installation price, Lost job, Unemployed, I'm on (any form of social assistance), etc. The following answers should be coded to "Moved": Respondent moved, on vacation, going south for the winter, moved and don't want/need the phone anymore etc. The following answers should be coded to "Any other response": Sharing a phone, receiving harassing calls, getting a private number, hard of hearing, dissatisfied with the phone company, don't want other household members to receive calls, etc.
Universe:	Respondents whose reason for not having a phone is because they cancelled it (RTS-Q02=1)

#### RTS-Q04

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. INTERVIEWER: READ ALL CATEGORIES AND MARK ALL THAT APPLY

<-1>	Don't know
<-2>	Refused

<1> the installation charge <2> the security deposit

<3> monthly charge for your basic phone line which includes local

calls

<4> optional features and/or set charges

<5> long distance charges

<6> other usage charges (i.e. 900 service, \* features, directory

assistance, etc.)

Universe: Respondents who cannot afford a phone RTS-Q02=2 or RTS-Q03=1

#### RTS-Q05

If there were an emergency at home, would members of your household have easy access to a neighbour's phone?

<-1> Don't know <-2> Refused <1> YES <2> NO

Universe: Those who answered 1' (0 telephone number) in RTS-Q01

#### RTS-Q06

If there were an emergency at home, would members of your household have easy access to a payphone near your residence?

<-1> Don't know <-2> Refused <1> YES <2> NO

Universe: Those who answered 1' (0 telephone number ) in RTS-Q01

#### RTS-Q07

If there were an emergency at home, would any member of your household have convenient access to a telephone near your residence, at another location not already mentionned?

<-1> Don't know <-2> Refused <1> YES <2> NO

Universe: Those who answered 1' (0 telephone number ) in RTS-Q01

#### RTS-Q08

In %RTS->RFYEAR%, was your total annual family income before taxes and deductions less or more than \$%RTS->RTS\_LICO%?

<-1> Don't know <-2> Refusal <1> Less <2> More

Universe: All respondents

#### RTS-I08

Thank you for your cooperation.

Note: The application should go back to the LFS component screen after this has

been read out to respondents.

Universe: All respondents

## 14.0 Record Layout and Univariates

## Record Layout - RESIDENTIAL TELEPHONE SERVICES SURVEY (0599 - re-weighted)

Variable:	PROV1	Position:	1	Length:	2	
Province						
					FREQ	WTD
10	Newfour	ndland			1,523	190,297
11	Prince E	dward Island			1,104	51,071
12	Nova Sc	eotia			2,743	361,986
13	New Bru	unswick			2,235	281,149
24	Quebec				7,906	2,980,703
35	Ontario				11,917	4,277,134
46	Manitob	a			2,986	424,627
47	Saskatch	newan			3,092	383,846
48	Alberta				3,103	1,079,912
59	British (	Columbia			3,731	1,551,915
					40,340	11,582,640

Variable:	CMA1	Position:	3	Length:	2		
CMA							
					FREQ	WTD	
00	Non CM	ſΑ			26,840	5,011,525	
01	Halifax	17.1			631	141,070	
02	Quebec				584	288,082	
03	Montrea	1			1,600	1,423,637	
04	Ottawa	•			693	320,344	
05	Toronto				2,144	1,674,866	
06	Kitchene				649	157,228	
07	Hamilto	n			576	254,830	
08	St. Cath	arines			572	142,435	
09	London				631	169,927	
10	Windson	r			506	119,774	
11	Winnipe	eg			1,478	271,612	
12	Calgary				729	347,446	
13	Edmonto	on			820	346,439	
14	Vancouv				1,384	781,490	
15	Victoria				503 =====	131,935	
					40,340	11,582,640	
Variable:	Q01	Position:	5	Length:	1		
How many di business)	fferent telephone	numbers are there	for your res	sidence? (Include o	cellular phone numbe	rs and phone numbers u	sed for
					FREQ	WTD	
1	0 None				759	194,951	
2	1				30,180	8,502,928	
3	2				6,711	2,017,681	
4	3 or mor	re			2,690	867,080	
					40,340	11,582,640	

Variable:	Q02	Position:	6	Length:	1			
Why don't you	Why don't you have a phone?							
1 2 3 4 6 7 8	I cancelled Can't affor Moved All other re Valid skip Don't know Refused	d it esponses				FREQ 80 438 27 192 39,581 0 22 ====== 40,340	WTD 20,788 109,276 5,927 48,846 11,387,689 0 10,114 ===================================	
Variable:	Q03	Position:	7	Length:	1			
Why did you ca	ncel it?							
1 2 3 6 7 8	Can't affor Moved All other r Valid skip Don't knov Refused	esponses				FREQ 40 10 30 40,260 0 0 ====== 40,340	WTD 12,306 2,032 6,450 11,561,852 0 0 =================================	
Variable:	Q041	Position:	8	Length:	1			
I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford the installation charge								
1 6 7 8	Yes Valid skip Don't knov Refused	v				FREQ 352 39,980 4 4 4 ======= 40,340	WTD 86,775 11,493,187 1,559 1,120 ====================================	

Variable: Q042 Position: 9 Length: 1

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. ... the security deposit

		FREQ	WTD
2	Yes	266	64,896
6	Valid skip	40,066	11,515,066
7	Don't know	4	1,559
8	Refused	4	1,120
		======	========
		40,340	11,582,640

Variable: Q043 Position: 10 Length: 1

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. ... monthly charge for your basic phone line which includes local calls

			FREQ	WID
3	3	Yes	326	80,243
6	5	Valid skip	40,006	11,499,718
7	7	Don't know	4	1,559
8	3	Refused	4	1,120
			======	========
			40,340	11,582,640

Variable: Q044 Position: 11 Length: 1

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. ... optional features and/or set charges

		FREQ	WTD
4	Yes	139	33,438
6	Valid skip	40,193	11,546,523
7	Don't know	4	1,559
8	Refused	4	1,120
		======	========
		40,340	11,582,640

Variable: Q045 Position: 12 Length: 1

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. ... long distance charges

		FREQ	WTD
5	Yes	182	39,463
6	Valid skip	40,150	11,540,499
7	Don't know	4	1,559
8	Refused	4	1,120
		40,340	11,582,640

Variable: Q046 Position: 13 Length: 2

I am going to read a list of most common charges which could be on a telephone bill. Please tell me which of these charges you find difficult to afford. ... other usage charges (i.e. 900 service, \* features, directory assistance, etc.)

		FREQ	WTD
06	Yes	107	25,522
96	Valid skip	40,225	11,554,439
97	Don't know	4	1,559
98	Refused	4	1,120
		======	========
		40,340	11,582,640

Variable: Q05 Position: 15 Length: 1

If there were an emergency at home, would members of your household have easy access to a neighbour's phone?

		FREQ	WID
1	Yes	622	149,809
2	No	99	31,497
6	Valid skip	39,581	11,387,689
7	Don't Know	9	2,851
8	Refused	29	10,794
		======	========
		40,340	11,582,640

Variable:	Q06	Position:	16	Length:	1	
If there were	an emergency at hom	ie, would mem	bers of your l	household have e	asy access to a payph	one near your residence?
					FREQ	WTD
1	Yes				417	117,641
2	No				300	61,572
5	Valid skip				39,581	11,387,689
7	Don't Know				6	1,999
3	Refused				36 =====	13,739
					40,340	11,582,640
: -1.1	007	D = -1/d =	17	Landh	1	
Variable:	Q07	Position:	17	Length:	1	
	an emergency at home	•	nember of yo	ur household hav	re convenient access t	o a telephone near your res
					FREQ	WTD
	Yes				381	95,868
	No				329	78,571
	Valid skip				39,581	11,387,689
•	Don't know				10	4,840
3	Refused				39	15,672
					40,340	11,582,640
Variable:	Q08	Position:	18	Length:	1	
	_				1 or more than \$%RTS	->RTS_LICO%?
	_					->RTS_LICO%? WTD
n 1999, was	_				or more than \$%RTS	
n 1999, was	your total annual fam				or more than \$%RTS	WTD
n 1999, was	your total annual fam  Less				or more than \$%RTS FREQ 8,633	WTD 2,537,533
in 1999, was 1 2	your total annual fam Less More				or more than \$%RTS FREQ 8,633 29,570 1,308 829	WTD 2,537,533 8,453,415 355,256 236,437
	your total annual fam  Less  More  Don't know				or more than \$%RTS  FREQ 8,633 29,570 1,308	WTD 2,537,533 8,453,415 355,256

Weight - decimal in 5th byte of the field

Variable:	SZCODE1	Position:	28	Length:	1			
SZCODE1								
1 2 3 4 5 6	Urban, po Urban, po Urban, po	pulation of 500,0 pulation of 100,0 pulation of 30,00 pulation of 15,00 pulation under 13	)00 - 49,999 )0 - 99,999 )0 - 29,999			FREQ 10,008 9,821 3,548 2,189 4,916 9,858 ======= 40,340	WTD 5,708,746 1,822,789 939,563 386,055 937,132 1,788,355 ======== 11,582,640	
Variable:	SZCODE3	Position:	29	Length:	2			
10 20 30 40 51 52 60	Urban, po Urban, po Urban, po Urban, po	pulation of 500,0 pulation of 100,0 pulation of 30,00 pulation of 15,00 pulation of 2,500 pulation under 2,	)00 - 49,999 )0 - 99,999 )0 - 29,999 ) - 14,999			FREQ 10,008 9,821 3,548 2,189 4,116 800 9,858 ======= 40,340	WTD 5,708,746 1,822,789 939,563 386,055 781,498 155,634 1,788,355 ===================================	
Variable:  Case Identifies	REALUKEY r *** suppressed	Position:	31 a file ***	Length:	15			
	REPL	Position:	46	Length:	2			
	for variance estimate							